arXiv:1111.4152v1 [math.NT] 17 Nov 2011

HYPERELLIPTIC CURVES AND THEIR INVARIANTS: GEOMETRIC, ARITHMETIC AND ALGORITHMIC ASPECTS

REYNALD LERCIER AND CHRISTOPHE RITZENTHALER

Abstract. We apply classical invariant theory of binary forms to explicitly characterize isomorphism classes of hyperelliptic curves of small genus and, conversely, propose algorithms for reconstructing hyperelliptic models from given invariants. We focus on genus 3 hyperelliptic curves. Both geometric and arithmetic aspects are considered.

Contents

Introduction 2
1. Invariants and symbolic computations 5
1.1. Terminology 5
1.2. Algebra of invariants and covariants 6
1.3. Hyperelliptic curves and invariants of binary forms 7
1.4. Algorithms in weighted projective spaces 8
1.5. Fundamental invariants and covariants for the binary octics 9
2. Reconstruction of binary forms 11
2.1. Clebsch's identities 11
2.2. A generic reconstruction algorithm 14
2.3. Reconstruction in the hyperelliptic genus 3 case 16
3. Automorphisms and strata of hyperelliptic genus 3 curves 18
3.1. Review on automorphism groups 18
3.2. Identification of loci and reconstruction with non trivial automorphism groups 19
4. Field of definition 32
4.1. Field of definition and field of moduli: general facts 32
4.2. The hyperelliptic case 33
4.3. The hyperelliptic case with no extra-automorphism 35
4.4. The hyperelliptic case with extra-automorphisms 36
4.5. The hyperelliptic case of genus 3 36
4.6. The case of curves over finite fields 39
5. Conclusion 40
References 40
Appendix A. Stratum equations for the automorphism group $\mathbf{D}_4$ 42

Date: November 18, 2011.
2010 *Mathematics Subject Classification.* 14Q05 ; 13A50 ; 14H10 ; 14H37.
Key words and phrases. invariant ; covariant ; binary forms ; field of moduli ; field of definition ; automorphism ; reconstruction ; genus 3 ; moduli space ; weighted projective space ; algorithm.
Both authors acknowledge support by grant ANR-09-BLAN-0020-01.

Introduction

Invariant theory played a central role in 19th century algebra and geometry, yet many of its techniques and algorithms were practically forgotten by the middle of the 20th century and replaced by the abstract and powerful machinery of modern algebraic geometry. However, motivated by computational applications to cryptography, robotics, coding theory, *etc.*, the classical invariant theory has come to a renaissance. Among classical groups, the natural action of SL_2 on binary forms has received most attention. One reason is the remarkable formalism developed by Gordan in 1868 to compute a finite set of generators of invariants. The other reason is the application of hyperelliptic curves in cryptography, especially for the so-called CM methods (see [CFA+06]).

When K is an algebraically closed field of characteristic 0 or a prime $p \neq 2$, hyperelliptic curves of genus $g > 1$ are indeed naturally related to the action of $\mathrm{SL}_2(K)$ on binary forms of even degree $n = 2g+2$. Thus, isomorphisms between curves with equation $y^2 = f(x)$ and $f \in K[x]$ of degree n are globally determined by a $\mathrm{SL}_2(K)$ action on f (see Section 1.3). Thus the K-isomorphism classes of hyperelliptic curves of genus g can be represented by the values of a finite set of invariants for $SL_2(K)$. In this way, exploring properties of the invariants lead to effective results on the geometry and arithmetic of the hyperelliptic moduli space.

On the geometric level, to make explicit the representation of the classes by invariants, we have to tackle a double task: compute a set of generators $\{I_i\}$ of invariants and reciprocally construct a curve from given values of these invariants. We call the latter the *reconstruction* phase. As a by-product, we also want to be able to read some geometric information, in particular the automorphism group of a curve, from the invariants.

The first issue can be addressed thanks to Gordan's method which is based on a differential operator called *transvectant*, see Section 1. Contrary to the genus 2 case which is described by 3 algebraically independent invariants, Shioda [Shi67] gave in genus 3 a basis of 9 invariants (which we call *Shioda invariants*), the first 6 being algebraically independent and the last three related to the others by 5 explicit relations. Besides this, unlike the genus 2 case and the classical Igusa invariants, the discriminant is not an element of this basis. We therefore decided not to use the usual representation based on 'absolute invariants' and to switch to a weighted projective space of invariants for which we rely on some specific algorithms to test equality or to create points. Note that although we restrict to genus 3 in the present article, these algorithms apply to any weighted projective space and are therefore useful for hyperelliptic curves of higher genus too.

The main algorithm for the second issue relies on Mestre's method which he exposes for $g = 2$ in [Mes91]. It is based on computations going back to [Cle72, §103] (see Section 2) and uses a generalization of invariants called *covariants* (see Definition 1.2). Roughly speaking, starting from three order 2 covariants, one constructs a plane conic $\mathcal{Q}$ and a plane degree $g+1$ curve $\mathcal{H}$ whose coefficients are invariants, hence expressible in terms of the generators $\{I_i\}$. After specialization at given values of invariants, if $\mathcal{Q}$ is not singular, the degree 2-cover of $\mathcal{Q}$ ramified at the $2(g+1) = n$ points of intersection of $\mathcal{Q}$ and $\mathcal{H}$ is a hyperelliptic curve with invariants equal to the initial values. In order to make this practical, one of the main computational difficulty is to find the formal expressions of the coefficients in terms of $\{I_i\}$ as the degree and number of variables are for $g = 3$ already quite large. We by-passed the difficulty using an evaluation-interpolation strategy.

For genus 2, this algorithm enabled Mestre to reconstruct hyperelliptic curves C with no *extra-automorphism*, *i.e.* $\mathrm{Aut}(C)$ is generated by the hyperelliptic involution ι. Indeed, it was proved in [Bol87] that the 'classical' $\mathcal{Q}$ is always non-singular in this case. It is however always singular when C has extra-automorphisms and it can not be used anymore. Cardona and Quer in [CQ05] completed the picture by a different choice of order 2 covariants which lead to another conic $\mathcal{Q}$ non-singular in the case $\mathrm{Aut}(C) \simeq (\mathbb{Z}/2\mathbb{Z})^2$ (for bigger automorphism groups, explicit parametrizations were already known). The two authors of the present article have implemented these constructions with the computational algebra software MAGMA and generalized them to the remaining fields of small characteristics (3 and the most difficult case 5) [LR08]. Genus 2 can be considered as solved.

It is thus natural to turn to genus 3. Here also, the issues that we tackle are naturally stratified by the automorphism group of the curve (see Table 3 and Figure 1 for the lattice of automorphism groups in genus

3). In Section 3.2, we find equations for all the strata and show how to reconstruct a curve with given invariants. These two questions are indeed intertwined.

Starting from normal models as in Table 3, we get necessary algebraic conditions for the invariants to define a curve with a certain automorphism group. We determine these equations by evaluations and interpolations, the same approach as the one we followed for finding the expressions of the coefficients of $\mathcal{Q}$ and $\mathcal{H}$. Since we only use Gröbner techniques to reduce bases, we were able to obtain equations for all the strata, even for the dimension 3 stratum $(\mathbb{Z}/2\mathbb{Z})^2$. In order to check that these conditions are also sufficient, we reconstruct a curve from given invariants and check that its Shioda invariants are equal to the original ones. This last step involves calculations in the quotient ring of $\mathbb{Q}[J_2, ..., J_{10}]$ by the ideal defined by the stratum equations. Even if modern computational algebra software can handle them, keeping polynomials reduced to normal forms modulo the ideal of relations yields several hours of computations on a powerful computer, at least for the strata of dimension 2 or 3. A reasonably optimized program written in MAGMA that implements the corresponding computations is available on the web page of the authors for independent checks.

The reconstruction step for most strata is carried out by 'inverting' the expressions of the invariants in terms of the parameters of normal models modulo the stratum equations. For strata of dimension less or equal to 1, we give models whose coefficients are algebraic expressions in terms of the invariants. For the dimension 2 stratum with automorphism group $(\mathbb{Z}/2\mathbb{Z})^3$, we can still work out these computations at the price of a 'cubic extension' (see Lemma 3.14). For the dimension 2 stratum $\mathbb{Z}/4\mathbb{Z}$, we exhibit 5 conics among which at least one is always non-singular and use Mestre's method for this stratum. The dimension 3 stratum $(\mathbb{Z}/2\mathbb{Z})^2$ is more challenging. One can show (Lemma 3.2) that *any* choice of 3 covariants in the set of the 14 fundamental covariants of order 2 (364 possibilities) leads to a singular conic, hence Cardona and Quer's patch is not possible for curves in this stratum. Our computational approach yields, in addition to a set of 24 necessary equations for the invariants to define a curve with automorphism group $(\mathbb{Z}/2\mathbb{Z})^2$, an explicit reconstruction at the price here of a 'degree 8 extension' (Lemma 3.20). We noticed furthermore that the singularity of the 364 conics is equivalent to the nullity of only 19 determinants. Obviously, the locus where these determinants simultaneously vanish contains the stratum $(\mathbb{Z}/2\mathbb{Z})^2$. We show that it is actually *equal* to the 24 stratum equations for $(\mathbb{Z}/2\mathbb{Z})^2$. As a by-product, the reconstruction of curves with no extra-automorphisms can therefore be achieved thanks to Mestre's method by picking one of the non-singular conics $\mathcal{Q}$ among the 19 fixed ones.

So far we have avoided arithmetic issues by working over an algebraically closed field but new challenging and deep issues arrive when one consider over which 'minimal field' these constructions can be achieved. Assume for simplicity that k is a field of characteristic 0 and let C be a curve defined over k (see a more general framework in Section 4.1). One can consider the intersection of all the subfields k' of $K = \bar{k}$ over which there exists a curve K-isomorphic to C (k' is called a *field of definition*). This field is called the *field of moduli* and denoted $\mathbf{M}_C$. If it is a field of definition, it is the minimal field of definition of C. Unfortunately, this is not automatically the case if for instance C has a non-trivial automorphism group. Therefore hyperelliptic curves are highly concerned with this issue: when is $\mathbf{M}_C$ a field of definition ? On top of this, hyperelliptic curves lead to a refined question that we want to address too. Let remember that, in full generality, a hyperelliptic curve C/k is a curve with a degree 2 morphism from C to a non-singular plane conic Q. If Q has a point and, assuming the characteristic of k not 2, one can write $C/k : y^2 = f(x)$ with $f \in k[x]$. We say that C can be *hyperelliptically defined over* k or that C admits a *hyperelliptic equation over* k. This is obviously the case if k is algebraically closed or finite (and therefore people often use this property as a definition) but when k is arbitrary, there might be again an obstruction.

When $g = 2$ (or more generally even), Mestre showed that the two questions are equivalent. Moreover in the case where C has no extra-automorphism, he showed that $\mathbf{M}_C$ is a field of definition if and only if the conic $\mathcal{Q}$ constructed from the covariants has a rational point. When C has extra-automorphisms, Cardona and Quer were able to exhibit hyperelliptic equations over $\mathbf{M}_C$. We implemented and completed their results for fields of characteristic 2, 3 and 5 [LR08].

For $g = 3$, we address both questions, from a theoretical and computational point of view. Let us give some of the results that we have obtained according to the stratification of the moduli space by the automorphism groups (see Section 4.5).

- For dimension 0 and dimension 1 strata, since $\mathrm{Aut}(C)/\langle\iota\rangle$ is not cyclic, Huggins' results in [Hug05] show that there is a hyperelliptic equation over $\mathbf{M}_C$. This is confirmed and made explicit by the computations we performed for the reconstruction. In theses cases we can even exhibit parametrized models over $\mathbf{M}_C$.
- The dimension 2 case $\mathrm{Aut}(C) \simeq (\mathbb{Z}/2\mathbb{Z})^3$ is theoretically covered by [Hug05] and therefore there exists a hyperelliptic equation over $\mathbf{M}_C$ (see Remark 4.16 for the controversy on this subject). In the present article, however we only reconstruct the curve hyperelliptically over at most a cubic extension of $\mathbf{M}_C$. There is a real difficulty to perform an explicit descent over $\mathbf{M}_C$ as geometric isomorphisms between our curve and a model over $\mathbf{M}_C$ might be defined over a degree 24 of $\mathbf{M}_C$. In a forthcoming article, we work out an algorithm based on covariants to obtain a model over $\mathbf{M}_C$ in an efficient way.
- The dimension 2 case $\mathrm{Aut}(C) \simeq \mathbb{Z}/4\mathbb{Z}$ is not covered by the general result of [Hug05]. However using the special form of the ramification signature, we can show that C can always be hyperelliptically defined over $\mathbf{M}_C$. To make this result explicit, we use the fact that there is always a non-singular conic $\mathcal{Q}$. Although this conic has not necessarily a rational point, we can make use of the special shapes of $\mathcal{Q}$ and $\mathcal{H}$ to perform an explicit hyperelliptic descent.
- For the dimension 3 case $\mathrm{Aut}(C) \simeq (\mathbb{Z}/2\mathbb{Z})^2$, Huggins constructed examples of genus 3 curves over $\bar{\mathbb{Q}}$ which cannot be defined over their field of moduli. Hence, we did not explore this case further and our reconstruction takes place over a degree 8 extension at most.
- Finally, for the dimension 5 case $\mathrm{Aut}(C) \simeq \mathbb{Z}/2\mathbb{Z}$, we show that $\mathbf{M}_C$ is always a field of definition. This is more generally true for hyperelliptic curves with no extra-automorphisms of odd genus. However, now C has not automatically a hyperelliptic model over $\mathbf{M}_C$. In genus 3, we show that C has a hyperelliptic equation over $\mathbf{M}_C$ if and only if one (equivalently, all of the) non-singular conic $\mathcal{Q}$ has a rational point over $\mathbf{M}_C$. As we proved that such a conic always exists, the method is explicit as well.

With a view to applications over finite fields, we want hyperelliptic equations over $\mathbf{M}_C$ in all cases. On one hand, the task is made easier as there is never any obstruction for the curve to have a hyperelliptic equation over its field of moduli and we propose an algorithm which makes the reconstruction over $\mathbf{M}_C$ effective. Moreover, starting from the rational parametrizations that we exhibit for most of the strata, we can state the exact number of isomorphism classes of rational curves with a given automorphism group, except for the dimension 3 stratum $\mathrm{Aut}(C) \simeq (\mathbb{Z}/2\mathbb{Z})^2$. On the other hand, as already illustrated by the genus 2 case, strange phenomenon can happen when the characteristic p is too small. This is not so surprising as the stratification itself may be different when $p \leq 2g+1$. We took special care of denominators when computing invariants and covariants and our results are then naturally valid for $p > 7 = 2 \cdot 3 + 1$. We wonder whether the natural bound $p > 2g + 1$ may be reached in this way for general g.

Finally, this article is only the emerged part of the iceberg. A MAGMA code[1] containing the various algorithms to compute invariants and reconstruct the curves is available on the web page of the authors. We tested it over $\mathbb{F}_p$ for $11 \leq p \leq 47$ and checked that we actually obtain the p^5 non-isomorphic curves predicted by the theory. We plan to include soon the computation of twists to obtain a representative of each $\mathbb{F}_q$-isomorphism class as we did for genus 2.

Notation and convention. In the following, the integer p is a prime or 0, k denotes any field of characteristic p and K an algebraically closed field of characteristic p. For a integer g, a *hyperelliptic curve* or a *genus g curve* C over a field k is always assumed to be smooth, projective and absolutely irreducible (if it is given by a singular equation, typically the case for hyperelliptic curves, C is the smooth model associated to this equation). However a 'curve' (like a conic) may be singular. When we speak of a morphism from C to C' we always mean a morphism *defined over k*. However the notation $\mathrm{Aut}(C)$ stands for $\mathrm{Aut}_{\bar{k}}(C)$.

Acknowledgments. This paper has benefit from plenty of discussions with several mathematicians. During the Huggins / Fuertes-González-Diez controversy, the authors wish to thank the help and patient listening

[1] published under the GNU Lesser General Public License

of Nils Bruin, Pierre Dèbes, Yolanda Fuertes, Gabino González-Diez, Ruben Hidalgo, Everett Howe, Bonnie Huggins, David Kohel, Stéphane Louboutin, Enric Nart, Xavier Xarles and finally Alexey Zykin who had the sharpest eye. Proposition 4.14 has been suggested to us by Nils Bruin and we thank Jean-Marc Couveignes for indicating to us the reference [Cou94] which lead to Proposition 4.17 and Jean-François Mestre for the precise reference to [Cle72].

1. Invariants and symbolic computations

1.1. **Terminology.** We let k be an infinite field of characteristic p so we can identify the graded algebra of polynomial functions over a finite k-vector space E with the graded symmetric algebra $S(E^*)$. If $E = \oplus_{i=1}^m E_i$, a function ϕ is homogeneous of degree $(n_1, \dots, n_m)$ if it is homogeneous of degree n_i on E_i for $1 \leq i \leq m$, *i.e.* if $\phi(f_1, \dots, \lambda f_i, \dots, f_m) = \lambda^{n_i} \cdot \phi(f_1, \dots, f_m)$ for all $\lambda \in K$ and all $(f_1, \dots, f_m) \in E$.

Let $V = k^2$ be seen as the space of vectors with coordinates (x, z) and let $E = S^n(V^*)$ be the $(n+1)$-dimension vector space of homogeneous polynomials functions of degree n in $X = (1, 0)^*$ and $Z = (0, 1)^*$. We write a generic element of E as $\sum_{i=0}^n a_i X^i Z^{n-i}$ with $a_i \in k$. We let a group $G \subset \mathrm{GL}_2(k)$ acts on V in the natural left action, *i.e.* if

$$M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in G$$

then

$$M.(x, z) = (ax + bz, cx + dz).$$

Let ρ be the contragredient representation on V^*. We let the group G act with the tensorial representation $\otimes^n \rho$ on E. More concretely, if $f \in E$ then $M.f$ is defined by $(M.f)(x, z) = f(M^{-1}.(x, z))$. This can be extended to $E = \oplus_{i=1}^m S^{n_i}(V^*)$ with $m \geq 1$ and integers $n_i > 0$ using the representation $\bigoplus_i \otimes^{n_i} \rho$. In this setting, we recover the classical definition of invariants and covariants.

Definition 1.1. A homogeneous polynomial function $I : E \to k$ is a *(relative) invariant* if there exists $\omega \in \mathbb{Z}$ such that for all $M \in G$ and all $(f_1, \dots, f_m) \in E$, we have

$$I(M.f_1, \dots, M.f_m) = \det(M)^{-\omega} \cdot I(f_1, \dots, f_m)\,.$$

The space of invariants is classically embedded into a broader framework.

Definition 1.2. A homogeneous polynomial function $C : E \oplus V \to k$ of degree $(d_1, \dots, d_m, r)$ is a *(simultaneous) covariant* if there exists $\omega \in \mathbb{Z}$ such that for all $M \in G$, all $(f_1, \dots, f_m) \in E$ and all $(x, z) \in V$, we have

$$C(M.f_1, \dots, M.f_m, M.(x, z)) = \det(M)^{-\omega} \cdot C(f_1, \dots, f_m, (x, z))\,.$$

Hence, covariants of order 0 are invariants. The integer ω is called the *weight* (or index), the m-uple $(d_1, \dots, d_m)$, the *degree* and the integer r is called the *order* of a covariant.

Example 1.3. When $E = S^n(V^*)$, the polynomial function $(\sum a_i X^i Z^{n-i}, (x, z)) \mapsto \sum_{i=0}^n a_i x^i z^{n-i}$ is a covariant of degree 1 and of order n. We will denote it by $\mathfrak{f}$ in the sequel.

The major operation to generate new covariants from given ones is the transvectant. For $i, j \in \mathbb{N}$ two distinct integers, let (x_i, z_i) and (x_j, z_j) be bases of two copies V_i and V_j of V. Let Ω_{ij} be the differential operator

$$\Omega_{ij} = \frac{\partial}{\partial x_i}\frac{\partial}{\partial z_j} - \frac{\partial}{\partial z_i}\frac{\partial}{\partial x_j}.$$

Let $r_i, r_j > 0$ be integers, $f_i \in S^{r_i}(V) = S^{r_i}(V_i)$ and $f_j \in S^{r_j}(V) = S^{r_j}(V_j)$. We define the following differential operators (where composition is denoted multiplicatively).

Definition 1.4. We define the *h-th transvectant*

$$(\ \ ,\ \ )_h : S^{r_i}(V) \times S^{r_j}(V) \to S^{r_i + r_j - 2h}(V)$$

as

$$(f_i, f_j)_h = \frac{(r_i - h)!\ (r_j - h)!}{r_i!\ r_j!} \cdot \left(\Omega_{ij}^h (f_i(x_i, z_i) \cdot f_j(x_j, z_j))\right)_{(x_i, z_i) = (x_j, z_j) = (x, z)}\,.$$

In the sequel for all $i \in \mathbb{N}$ we denote $\ell_i = \ell_i(x,z) = (\alpha_i x + \beta_i z)$ some linear forms with coefficients $\alpha_i, \beta_i \in k$ and $[\ell_i, \ell_j] = \alpha_i\beta_j - \alpha_j\beta_i$. Using the Clebsch-Gordan formula as in [Pro07, p.565] or by direct computation, one gets the following lemma which will turn to be central for proving the formula of Section 2.

Lemma 1.5. *For any r_i, r_j positive integers,*

$$(\ell_i^{r_i}, \ell_j^{r_j})_h = [\ell_i, \ell_j]^h \cdot \ell_i^{r_i-h}\ell_j^{r_j-h}.$$

If C is a homogeneous covariant of order r, we can see C as an homogeneous polynomial of degree r in the variables x, z, *i.e.* as an element of $S^r(V)$. In particular, $\mathfrak{f} = \sum a_i x^i z^{n-i}$. Since the r-th powers of linear forms span the vector-space $S^r(V)$ and the h-th transvectant is bilinear, we see that the h-th transvectant of two covariants of degree d_1, d_2 and of order r_1, r_2 is a covariant of degree $d_1 + d_2$ and of order $r_1 + r_2 - 2h$ (see also [Pro07, Chap.15] for a conceptual explanation).

1.2. **Algebra of invariants and covariants.** We focus on invariants and covariants for $G = \mathrm{GL}_2(K)$ and $G = \mathrm{SL}_2(K)$. In both groups, using the multiplication by -1 scalar matrix, we first notice that $n\sum d_i - r$ needs to be even in order for a covariant to be defined. Moreover, if C is a covariant for $\mathrm{GL}_2(k)$, then C is a covariant for $\mathrm{SL}_2(K)$. Conversely, let C be a covariant for $\mathrm{SL}_2(K)$ and let $g \in \mathrm{GL}_2(K)$. Since $g' = g/\sqrt{\det(g)} \in \mathrm{SL}_2(K)$, we have that

$$C(g'.f_1, \ldots, g'.f_m, g'.(x,z)) = C(f_1, \ldots, f_m, (x,z)).$$

Since C is homogeneous of degree $(d_1, \ldots, d_m)$ and of order r, for any $\lambda \in k$, we have that $t = \lambda \cdot \mathrm{id}$ acts as

$$C(t.f_1, \ldots, t.f_m, t.(x,z)) = \lambda^{n\sum d_i - r} \cdot C(f_1, \ldots, f_m, (x,z)),$$

and thus

$$C(g.f_1, \ldots, g.f_m, g.(x,z)) = \det(g)^{-(n\sum d_i - r)/2} \cdot C(f_1, \ldots, f_m, x, z).$$

This shows that covariants for $\mathrm{SL}_2(k)$ are equal to the covariants for $\mathrm{GL}_2(k)$, with weight ω equal to $(n\sum d_i - r)/2$.

Classically many results are known on the algebra $\mathcal{C}_n$ generated by the covariants of all degree and order under the action of $G = \mathrm{SL}_2(\mathbb{C})$ on $E = S^n(V^*)$ and on its sub-algebra $\mathcal{I}_n$ generated by invariants. Since Gordan [Gor68], it is known that $\mathcal{I}_n$ and $\mathcal{C}_n$ are finitely generated. Thanks to the so-called Clebsch-Gordan formula, one can even prove that $\mathcal{C}_n$ is generated by a finite number of iterations of transvectants starting from the single covariant $\mathfrak{f}$ (see [GY03]). Thus we can even assume that these generators are defined over $\mathbb{Q}$ and this allows to perform exact arithmetic operations with them. Whereas today this particular case is part of a more general result on reductive groups, Gordan's method is effective and generators for these algebra were given for $n \leq 6$ in the nineteenth century (see [Dix90]). Generators for $\mathcal{I}_7$ were determined by von Gall [VG88] and Dixmier and Lazard [DL86] and for $\mathcal{C}_7$ by Bedratyuk [Bed07]. The case of octics, $n = 8$, which is the case we are interested in this article will be reviewed and developed in Section 1.5. Finally, for $n = 9$ and $n = 10$, only generators for $\mathcal{I}_n$ have been determined (see [Cro02, BP10b],[BP10a]).

We summarize in Tab. 1 for each degree $2 \leq n \leq 10$, the number of generators for $\mathcal{I}_n$ and their degrees. We see that the number and the degree of the generators increase strongly with n, but remains affordable for some small degree n, especially for $n = 8$.

The algebra $\mathcal{I}_n$ is a sufficient tool to grasp the orbits of a generic binary form.

Theorem 1.6 ([MF82, p. 78],[Dix90, p. 47])**.** *Let f, f' be binary forms of even degree n greater than or equal to 3 over an algebraically closed field K with no roots of multiplicity greater than or equal to $n/2$. Let $\{I_i\}$ be a finite set of homogeneous generators of degree d_i for $\mathcal{I}_n$. Then f and f' are in the same orbit under the action of $\mathrm{GL}_2(K)$ (resp. $\mathrm{SL}_2(K)$) if and only if there exists $\lambda \in K$ such that for all i, $I_i(f) = \lambda^{d_i} \cdot I_i(f')$ (resp. $I_i(f) = I_i(f')$).*

Remark 1.7. In the theorem, the exponents are equal to the degrees d_i when one could have expected weights ω_i. Actually the two possibilities are equivalent since the weight is $n/2$ times the degree.

n	# Gen. of $\mathcal{I}_n$	Generator degrees *(exponents mean repetition)*
2	1	2
3	1	4
4	2	2, 3
5	4	4, 8, 12, 18
6	5	2, 4, 6, 10, 15
7	30	4, 8^3, 12^6, 14^4, 16^2, 18^9, 20, 22^2, 26, 30
8	9	2, 3, 4, 5, 6, 7, 8, 9, 10
9	92	4^2, 8^5, 10^5, 12^{14}, 14^{17}, 16^{21}, 18^{25}, 20^2, 22
10	106	2, 4, 6^4, 8^5, 9^5, 10^8, 11^8, 12^{12}, 13^{15}, 14^{13}, 15^{19}, 16^5, 17^5, 18, 19^2, 21^2

TABLE 1. Number and degrees of fundamental invariants for binary forms of degree n

Note that the multiplicity condition in Theorem 1.6 is as good as possible already for $n = 2$ since the only generator of $\mathcal{I}_2$, $a_1^2 - a_0a_2$, is 0 for quadratic forms in the two distinct orbits $f = 0$ and $f = X^2$. Since we will only deal with polynomials defining hyperelliptic curves, and hence having simple roots, this is not a limitation.

1.3. **Hyperelliptic curves and invariants of binary forms.** Let k be a field of characteristic p and $K = \bar{k}$. A curve C of genus $g \geq 2$ defined over k is said *hyperelliptic* if C/K has a separable degree 2 map to $\mathbb{P}^1_K$. Since the extension $K(C)/K(\mathbb{P}^1)$ is Galois, the curve C/K has an involution that we denote ι.

Lemma 1.8 ([Har77, Prop.IV.5.3])**.** *The automorphism ι is the unique involution on C/K such that $C/\langle\iota\rangle$ is of genus 0. It commutes with all automorphisms of C/K and is called the* hyperelliptic involution.

By unicity, ι is defined over k and induces a morphism $\rho : C \to Q = C/\langle\iota\rangle$ where Q/k is a genus 0 curve, not necessarily isomorphic to $\mathbb{P}^1$. However, if the curve Q has a rational point then it is isomorphic to $\mathbb{P}^1$ and $k(C)$ is a degree 2 separable cover of $k(\mathbb{P}^1) \simeq k(x)$. In that case, C is birationally equivalent to an affine curve of the form

$$y^2 + h(x)y = f(x)$$

where $\deg f \leq 2g + 2$ and $\deg h \leq g$. We say that C has a *hyperelliptic equation* if (a curve in the isomorphism class of C (over k)) can be written in the form above. A hyperelliptic curve has automatically a hyperelliptic equation when k is algebraically closed or a finite field and we recover the 'classical' definition of a hyperelliptic curve. However, as we shall see in Section 4.1 for more general fields and odd genus, it is not necessarily the case.

Let us assume from now on that $p \neq 2$. One can assume that $h = 0$ and the polynomial f has then simple roots. We say that f is a *hyperelliptic polynomial*. By homogenizing, $Y^2Z^{2g} = f(X, Z)$, we have $\deg f(X, Z) = 2g + 2$, taking into account a 'root' at infinity when $\deg f = 2g + 1$. With this convention, the roots of f are the ramification points of the cover C/Q and equivalently the Weierstrass points W of C. We will use these conventions for the roots and degree in the sequel when we speak about a hyperelliptic polynomial.

Remark 1.9. One cannot always assume that $\deg f = 2g + 2$ (as a polynomial) if k is not infinite. For instance the genus 2 curve $y^2 = x(x - 1)(x - 2)(x - 3)(x - 4)$ over $\mathbb{F}_5$ has no model with $\deg f = 6$ since all its affine Weierstrass points are rational.

The following proposition will give us the link between hyperelliptic curves and invariants for the action of GL_2.

Proposition 1.10. *Let $C : y^2 = f(x)$ and $C' : y^2 = f'(x)$ two hyperelliptic curves of genus g over a field k. Every isomorphism $\phi : C \to C'$ is given by an expression of the form*

$$(x, y) \mapsto \left(\frac{ax + b}{cx + d}, \frac{ey}{(cx + d)^{g+1}}\right),$$

for some $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{GL}_2(k)$ *and* $e \in k^*$. *The pair* (M, e) *is unique up to replacement by* $(\lambda M, \lambda^{g+1} e)$ *for* $\lambda \in k^*$. *The composition of isomorphisms* (M, e) *and* (M', e') *is* $(M'M, e'e)$.

Remark 1.11. We refer to [CNP05] for the characteristic 2 case.

Let assume now that $k = K$ is algebraically closed. The ramification divisor W of $\rho : C \to Q$ is a positive divisor of degree $2g + 2$. From Proposition 1.10, one easily gets

Corollary 1.12. *Let* C, C' *be two hyperelliptic curves of genus* g *with respective ramification divisor* W *and* W'. *Let* $D = \rho(W) \in \mathrm{div}(Q)$ *and* $D' = \rho'(W') \in \mathrm{div}(Q')$. *Then* C *and* C' *are isomorphic if and only if there exists an isomorphism* $\varphi : Q \to Q'$ *such that* $\varphi(W) = W'$.

Checking an isomorphism between C and C' in this way is often difficult since the points of the ramification divisor can be defined over large degree extensions. This is where invariants enter the game. Indeed, as in the proposition let $C : y^2 = f(x)$ and $C' : y^2 = f'(x)$. Since f and f' have only simple roots, we can use invariants to decide if the homogeneous binary forms $f(X, Z)$ and $f'(X, Z)$ of degree $2g + 2$ are in the same orbits under the action of $\mathrm{GL}_2(K)$. If so, one has $f'(X, Z) = f(aX + bZ, cX + dZ)$ for some $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{GL}_2(K)$ and so $(M, 1)$ defines an isomorphism between C and C'. Conversely, let (M, e) define an isomorphism between C and C'. we can adjust λ in Proposition 1.10 such that $e = 1$ and hence f' and f are in the same orbit under $\mathrm{GL}_2(K)$. Applying Theorem. 1.6 to the situation yields thus the following proposition.

Proposition 1.13. *Let* $\{I_i\}$ *be a set of homogeneous generators of degree* d_i *for* $\mathcal{I}_{2g+2}$. *Two hyperelliptic curves* $C : y^2 = f(x)$ *and* $C' : y^2 = f'(x)$ *of genus* g *are isomorphic if and only if there exists* $\lambda \in K^*$ *such that* $I_i(f') = \lambda^{d_i} \cdot I_i(f)$ *for all* i.

Hence, the possible values of a set of generators for $\mathcal{I}_{2g+2}$ up to this specific equivalence are in bijection with the points of the coarse moduli space H_g of hyperelliptic curves of genus g. We therefore need algorithms to handle such 'weighted sets'.

1.4. **Algorithms in weighted projective spaces.** In the context of curves of genus 1 or 2, one usually prefers to handle absolute invariants instead of homogeneous ones, by calculating ratios of homogeneous invariants of the same degree. In this way, one gets rid of the constant λ in Proposition 1.13. It becomes then easy to span the coarse moduli space or to check that two curves are isomorphic.

But, care has to be taken to ensure that the denominators of absolute invariants do not vanish for some hyperelliptic orbits and a common approach is to choose as denominator some powers of the discriminant invariant. We give up this strategy for higher genus curves, because the degree of the discriminant is too large (this degree is already equal to 14 for octics) and selecting invariants of smaller degree as denominators yields too many technical cases to consider. We choose instead to work with a (kind of) weighted projective space, given by m-uples $(I_1 : I_2 : \ldots : I_m)$ of weights d_1, d_2, $\ldots$, d_m. More precisely

Definition 1.14. Let k be a field and $d_1, \ldots, d_m$ positive integers wit $m \geq 2$. We call $\mathbb{W}$ a *weighted projective space* over k of dimension $m - 1$ and weights $(d_1, \ldots, d_m)$ the set of elements denoted $(\iota_1 : \ldots : \iota_m)$ which are equivalence classes of m-uples $(\iota_1, \ldots, \iota_m) \in k^m \setminus (0, \ldots, 0)$ for the relation

$$(\iota_1, \ldots, \iota_m) \sim (\iota'_1, \ldots, \iota'_m) \iff \exists\, \lambda \in \bar{k}^* \text{ such that } \iota_j = \lambda^{d_j} \cdot \iota'_j \; \forall\, 1 \leq j \leq m.$$

As a first tool, we need an algorithm for testing the equality of two points in a weighted projective space.

Proposition 1.15. *Let* k *be a field and* $\mathbb{W}$ *a* k*-weighted projective space of dimension* $m - 1$ *and weights* $(d_1, d_2, \ldots, d_m)$, *then Algorithm 1 tests if two elements of* k^m *are in the same class of* $\mathbb{W}$.

If k *is a field which admits operations with quasi-linear complexity in time and space (multiplications, inverses, tests), then Algorithm 1 has quasi linear complexity in time and space.*

Algorithm 1: Equality in a weighted projective space.

Input : Two elements $(U_1, U_2, \ldots, U_m)$ and $(V_1, V_2, \ldots, V_m)$ in k^m and $\mathbb{W}$ a k-weighted projective space of dimension m with weights $(d_1, d_2, \ldots, d_m)$.

Output: The boolean "`true`" if $(U_1 : U_2 : \ldots : U_m) = (V_1 : V_2 : \ldots : V_m)$, "`false`" otherwise.

$\mathcal{S}_U \leftarrow \{i \in \{1, \ldots, m\}$ s.t. $U_i \neq 0\}$; $\mathcal{S}_V \leftarrow \{i \in \{1, \ldots, m\}$ s.t. $V_i \neq 0\}$;
if $\mathcal{S}_U \neq \mathcal{S}_V$ **then**
  return `false`
d, $(c_i : i \in \mathcal{S}_U) \leftarrow$ `ExtentedGCD`$(d_i : i \in \mathcal{S}_U)$;
$\Lambda \leftarrow \prod_{i \in \mathcal{S}_U} (V_i/U_i)^{c_i}$;
return `true` *if* $V_i/U_i = \Lambda^{d_i/d}$ *for all* $i \in \mathcal{S}_U$, `false` *otherwise.*

Proof. If $(U_1, U_2, \ldots, U_m)$ and $(V_1, V_2, \ldots, V_m)$ are in the same class of $\mathbb{W}$, then there exists some $\lambda \in \bar{k}^*$ such that $V_i = \lambda^{d_i} \cdot U_i$ for all i. So, Λ is equal to $\prod (V_i/U_i)^{c_i} = \lambda^{\sum c_i d_i} = \lambda^d$, and thus we have $V_i/U_i = \Lambda^{d_i/d} = \lambda^{d_i}$ for $i \in \mathcal{S}_U$. Conversely, if $\mathcal{S}_U = \mathcal{S}_V$ and $V_i/U_i = \Lambda^{d_i/d}$ for $i \in \mathcal{S}_U$, then let λ be a d-th root of Λ in $\bar{k}^*$ and we can easily check that $V_i = \lambda^{d_i} \cdot U_i$. □

We deduce from Algorithm 1 how to associate to a class $(U_1 : U_2 : \ldots : U_m)$ a unique vector of k^m representing the class. Let $\mathcal{S}_U$ be the support of U, *i.e.* $\mathcal{S}_U = \{i \in \{1, \ldots, m\}$ s.t. $U_i \neq 0\}$, let c_i be integers such that $\sum_{i \in \mathcal{S}_U} c_i d_i = d$ with $d = \gcd(d_i : i \in \mathcal{S}_U)$ and let $\Lambda = \prod_{i \in \mathcal{S}_U} U_i^{c_i}$. Set $u_i = U_i/\Lambda^{d_i/d}$ for $i \in \mathcal{S}_U$ and $u_i = 0$ otherwise. Then $(u_1, u_2, \ldots, u_m)$ is the unique representative of $(U_1 : U_2 : \ldots : U_m)$ such that $\prod_{i \in \mathcal{S}_U} u_i^{c_i} = 1$.

When k is a finite field, enumerating elements in $\mathbb{W}$ is another feature that is needed to span $\mathsf{H}_g(k)$. In weighted projective spaces, this can be easily done by enumerating representatives. Typically, for each support $\mathcal{S}_U$, considered in turn, fix, for once, integers c_i such that $\sum_{i \in \mathcal{S}_U} c_i d_i = \gcd(d_i : i \in \mathcal{S}_U)$. Then enumerate all the vectors $(u_1, u_2, \ldots, u_m)$ of k^m with support $\mathcal{S}_U$ such that $\prod_{i \in \mathcal{S}_U} u_i^{c_i} = 1$.

Example 1.16. Let $k = \mathbb{F}_7$ and $\mathbb{W}$ be a k-weighted projective space of dimension 2 and weights (5, 7). To enumerate elements in $\mathbb{W}$, we consider supports in turn. The supports $\mathcal{S} = \{1\}$ and $\mathcal{S} = \{2\}$, yields the classes $(1 : 0)$ and $(0 : 1)$. Let us now consider the support $\mathcal{S} = \{1, 2\}$. We fix $c_1 = 3$, $c_2 = -2$, so that $5c_1 + 7c_2 = \gcd(5, 7)$. This yields 6 representatives $(U_1 : U_2)$ such that $\prod_{i=1,2} U_i^{c_i} = 1$, that is $(1 : 1)$, $(1 : 6)$, $(2 : 1)$, $(2 : 6)$, $(4 : 1)$, $(4 : 6)$. Another choice of c_1 and c_2 would lead to different representatives: for instance $c_1 = 24$ and $c_2 = -17$ gives $(1 : i)$ for $1 \leq i \leq 6$.

Finally, enumerating points on a coarse moduli space can be translated into enumerating points on a variety defined inside a weighted projective space. This is much more intricate than enumerating the full space, at least for curves of genus larger than 2. A naive strategy consists in enumerating all the points in the ambient space and for each point check if it is defined on the projective moduli variety. It is often inefficient. More sophisticated methods may be possible, based on a nice description of the variety, especially when the variety is rational and one has an explicit parametrization for it. In this direction, we give in Section 1.5 an efficient method for the moduli space of genus 3 hyperelliptic curves.

1.5. **Fundamental invariants and covariants for the binary octics.** According to the syzygy theorem of Hilbert, the graded ring $\mathcal{I}_8$ of invariants of binary octics fits into a finite exact sequence of $\mathbb{C}[\mathsf{X}]$-module where $\mathbb{C}[\mathsf{X}] := \mathbb{C}[X_2, \ldots, X_{10}]$. In what Mumford called a 'tour de force', Shioda managed to find an explicit description.

Theorem 1.17 ([Shi67], Th. 3, p. 1042)**.** *The graded ring $\mathcal{I}_8$ of invariants of binary octics is generated by 9 elements J_2, J_3, …, J_{10} of degree 2, 3, …, 10. There exists 5 generating relations, $\mathfrak{R}_i(J)$, of degree $15 + i$, $(i = 1, \ldots, 5)$, which, in turn, are connected by 5 fundamental first syzygies $\mathfrak{T}_i(R)$ of degree $24 + i$, $(i = 1, \ldots, 5)$. The second syzygy $\mathfrak{F}$ is unique up to constants and of degree 45. The syzygy sequence of $\mathcal{I}_8$ (as $\mathbb{C}[\mathsf{X}]$-module) is given by*

$$0 \to \mathbb{C}[\mathsf{X}]\mathfrak{F} \to \sum_{i=1}^{5} \mathbb{C}[\mathsf{X}]\mathfrak{T}_i \to \sum_{i=1}^{5} \mathbb{C}[\mathsf{X}]\mathfrak{R}_i \to \mathbb{C}[\mathsf{X}] \to \mathcal{I}_8 \to 0\,.$$

Using transvectants and the covariant $\mathfrak{f}$ from Example 1.3, Shioda defined the invariants J_i's of Theorem 1.17 in the following way,

$$J_2 = (\mathfrak{f},\mathfrak{f})_4,\ J_3 = (f,\mathfrak{g})_8,\ J_4 = (\mathfrak{k},\mathfrak{k})_4,\ J_5 = (\mathfrak{m},\mathfrak{k})_4,\ J_6 = (\mathfrak{k},\mathfrak{h})_4,$$
$$J_7 = (\mathfrak{m},\mathfrak{h})_4, J_8 = (\mathfrak{p},\mathfrak{h})_4,\ J_9 = (\mathfrak{n},\mathfrak{h})_4,\ J_{10} = (\mathfrak{q},\mathfrak{h})_4$$

where

$$\mathfrak{g} = (\mathfrak{f},\mathfrak{f})_4,\ \mathfrak{k} = (\mathfrak{f},\mathfrak{f})_4,\ \mathfrak{h} = (\mathfrak{k},\mathfrak{k})_2,\ \mathfrak{m} = (\mathfrak{f},\mathfrak{k})_4,\ \mathfrak{n} = (\mathfrak{f},\mathfrak{h})_4,\ \mathfrak{p} = (\mathfrak{g},\mathfrak{k})_4,\ \mathfrak{q} = (\mathfrak{g},\mathfrak{h})_4\,.$$

One may remark that none of these transvectant calculations involves covariants of order greater than 10. These formulae are thus valid, in addition to fields of characteristic 0, over any field of or characteristic greater than or equal to 11. We found for a generic form $f = a_8x^8 + a_7x^7 + \ldots + a_0$,

$$\begin{aligned}
J_2 &= 1/140\,(280\,a_0\,a_8 - 35\,a_1\,a_7 + 10\,a_2\,a_6 - 5\,a_3\,a_5 + 2\,{a_4}^2)\,,\\
J_3 &= 1/137200\,(11760\,a_0\,a_4\,a_8 - 7350\,a_0\,a_5\,a_7 + 3150\,a_0\,{a_6}^2 - 7350\,a_1\,a_3\,a_8 + 2205\,a_1\,a_4\,a_7 - 525\,a_1\,a_5\,a_6\\
&\quad + 3150\,{a_2}^2a_8 - 525\,a_2\,a_3\,a_7 - 330\,a_2\,a_4\,a_6 + 225\,a_2\,{a_5}^2 + 225\,{a_3}^2a_6 - 135\,a_3\,a_4\,a_5 + 36\,{a_4}^3)\,,\\
J_4 &= 1/3687936\,(8064\,a_0\,a_4\,{a_6}^2 - 614656\,a_0\,a_1\,a_7\,a_8 - 12544\,a_0\,a_2\,a_6\,a_8 + 53312\,a_0\,a_3\,a_5\,a_8 - 35280\,a_0\,a_3\,a_6\,a_7\\
&\quad + 4704\,a_0\,a_4\,a_5\,a_7 - 35280\,a_1\,a_2\,a_5\,a_8 - 13132\,a_1\,a_2\,a_6\,a_7 + 4704\,a_1\,a_3\,a_4\,a_8 + 3626\,a_1\,a_3\,a_5\,a_7\\
&\quad - 3864\,a_1\,a_4\,a_5\,a_6 - 3864\,a_2\,a_3\,a_4\,a_7 - 1516\,a_2\,a_3\,a_5\,a_6 - 320\,a_3\,{a_4}^2a_5 + 82320\,a_0\,a_2\,{a_7}^2\\
&\quad - 3360\,a_0\,{a_5}^2a_6 + 2458624\,{a_0}^2{a_8}^2 + 2401\,{a_1}^2{a_7}^2 + 1260\,a_1\,{a_5}^3 + 256\,{a_2}^2{a_6}^2 + 1260\,{a_3}^3a_7\\
&\quad + 589\,{a_3}^2{a_5}^2 - 25088\,a_0\,{a_4}^2a_8 + 82320\,{a_1}^2a_6\,a_8 + 3780\,a_1\,a_3\,{a_6}^2 + 784\,a_1\,{a_4}^2a_7 + 8064\,{a_2}^2a_4\,a_8\\
&\quad + 3780\,{a_2}^2a_5\,a_7 - 3360\,a_2\,{a_3}^2a_8 + 1984\,a_2\,{a_4}^2a_6 - 504\,a_2\,a_4\,{a_5}^2 - 504\,{a_3}^2a_4\,a_6 + 64\,{a_4}^4)\,,
\end{aligned}$$

etc.

The invariants $J_2, \ldots, J_7$ are algebraically independent (see [Shi67, Lemma 4, p. 1037]). We have moreover 5 relations between J_8, J_9 and J_{10}, of the form

$$\begin{array}{lcl}
\mathfrak{R}_1 & : & J_8^2 + A_6\,J_{10} + A_7\,J_9 + A_8\,J_8 + A_{16} = 0\\
\mathfrak{R}_2 & : & J_8\,J_9 + B_7\,J_{10} + B_8\,J_9 + B_9\,J_8 + B_{17} = 0\\
\mathfrak{R}_3 & : & J_8\,J_{10} + C_0\,J_9^2 + C_8\,J_{10} + C_9\,J_9 + C_{10}\,J_8 + C_{18} = 0\\
\mathfrak{R}_4 & : & J_9\,J_{10} + D_9\,J_{10} + D_{10}\,J_9 + D_{11}\,J_8 + D_{19} = 0\\
\mathfrak{R}_5 & : & J_{10}^2 + E_0\,J_2\,J_9^2 + E_{10}\,J_{10} + E_{11}\,J_9 + E_{12}\,J_8 + E_{20} = 0
\end{array} \tag{1}$$

where the A_i's, B_i's, C_i's and D_i's are invariants of degree i, that can be explicitly determine as functions of J_2, J_3, $\ldots$, J_7 (see [Shi67, Th. 1, p. 1030]). Moreover, following Shioda [Shi67, p. 1043]), we may consider $\mathfrak{R}_1$, $\mathfrak{R}_2$, $\mathfrak{R}_3$ and $(J_9 - B_9)\mathfrak{R}_2 - B_7\mathfrak{R}_4$ as linear equations in 1, J_9, J_9^2 and J_{10} and we obtain that J_8 always satisfies an equation of degree 5,

$$\begin{aligned}
&{J_8}^5 + (A_8 + 2\,B_8 + C_8){J_8}^4\\
&+ (-A_6\,C_{10} - A_7\,B_9 + 2\,A_8\,B_8 + A_8\,C_8 + A_{16} + B_7\,B_9\,C_0 + B_7\,C_0D_9 - B_7\,C_9 + {B_8}^2 + 2\,B_8\,C_8){J_8}^3\\
&+ (-A_6\,B_7\,C_0D_{11} - 2\,A_6\,B_8\,C_{10} - A_6\,{B_9}^2C_0 + A_6\,B_9\,C_9 - A_6\,C_{18} + A_7\,B_7\,C_{10} - A_7\,B_8\,B_9 - A_7\,B_9\,C_8 - A_7\,B_{17}\\
&+ A_8\,B_7\,B_9\,C_0 + A_8\,B_7\,C_0D_9 - A_8\,B_7\,C_9 + A_8\,{B_8}^2 + 2\,A_8\,B_8\,C_8 + 2\,A_{16}\,B_8 + A_{16}\,C_8 - {B_7}^2C_0D_{10} + B_7\,B_8\,C_0D_9\\
&- B_7\,B_8\,C_9 + B_7\,B_{17}\,C_0 + {B_8}^2C_8){J_8}^2 + (-A_6\,B_7\,C_0D_{19} + A_6\,B_8\,B_9\,C_9 + A_7\,{B_7}^2C_0D_{11} + A_7\,B_7\,B_8\,C_{10}\\
&- A_7\,B_8\,B_9\,C_8 - A_8\,{B_7}^2C_0D_{10} - A_8\,B_7\,B_8\,C_9 + A_8\,B_7\,B_{17}\,C_0 + A_{16}\,B_7\,B_9\,C_0 + A_{16}\,B_7\,C_0D_9 + A_{16}\,{B_8}^2 - A_6\,{B_8}^2C_{10}\\
&+ A_6\,B_{17}\,C_9 + A_7\,B_7\,C_{18} - A_7\,B_8\,B_{17} - A_7\,B_{17}\,C_8 + A_8\,{B_8}^2C_8 - A_{16}\,B_7\,C_9 - A_6\,B_7\,B_8\,C_0D_{11} + A_6\,B_7\,B_9\,C_0D_{10}\\
&- A_7\,B_7\,B_9\,C_0D_9 + A_8\,B_7\,B_8\,C_0D_9 - 2\,A_6\,B_9\,B_{17}\,C_0 - 2\,A_6\,B_8\,C_{18} + 2\,A_{16}\,B_8\,C_8)J_8 - A_6\,B_7\,B_8\,C_0D_{19}\\
&+ A_6\,B_7\,B_{17}\,C_0D_{10} - A_6\,{B_8}^2C_{18} + A_6\,B_8\,B_{17}\,C_9 - A_6\,{B_{17}}^2C_0 + A_7\,{B_7}^2C_0D_{19} + A_7\,B_7\,B_8\,C_{18} - A_7\,B_7\,B_{17}\,C_0D_9\\
&- A_7\,B_8\,B_{17}\,C_8 - A_{16}\,{B_7}^2C_0D_{10} + A_{16}\,B_7\,B_8\,C_0D_9 - A_{16}\,B_7\,B_8\,C_9 + A_{16}\,B_7\,B_{17}\,C_0 + A_{16}\,{B_8}^2C_8\,.
\end{aligned} \tag{2}$$

Generically, *i.e.* when $\delta = A_6\,J_8 + A_6\,B_8 - A_7\,B_7 \neq 0$, $\mathfrak{R}_1$ and $\mathfrak{R}_2$ yields

$$\begin{aligned}
J_9 &= (B_7\,{J_8}^2 - A_6\,B_9\,J_8 - A_6\,B_{17} + B_7\,A_8\,J_8 + A_{16}\,B_7)/\delta\,,\\
J_{10} &= -({J_8}^3 + {J_8}^2B_8 - A_7\,B_9\,J_8 - A_7\,B_{17} + A_8\,{J_8}^2 + A_8\,J_8\,B_8 + A_{16}\,J_8 + A_{16}\,B_8)/\delta\,.
\end{aligned}$$

Otherwise, *i.e.* when $\delta = 0$, one might have no rational solution for the system $(\mathfrak{R}_i)_{i=1\ldots5}$ in J_9 and J_{10} (for instance $J_2 = 1$, $J_3 = 0$, $J_4 = 0$, $J_5 = 0$, $J_6 = 8$, $J_7 = 0$, $J_8 = 7$ modulo 11) or several (for instance $J_2 = 9$, $J_3 = 0$, $J_4 = 0$, $J_5 = 0$, $J_6 = 2$, $J_7 = 0$, $J_8 = 0$ which has 3 solutions modulo 11 for (J_9, J_{10}): $(0,4)$, $(2,9)$ or $(9,9)$).

Remark 1.18. Instead of $\mathcal{I}_8$, one may look at $\mathrm{Frac}(\mathcal{I}_8)$ and consider absolute invariants. It is known that $\mathrm{Frac}(\mathcal{I}_n)$ is a rational function field (see [BK86]) and in the case $n = 8$, Maeda worked out 6 algebraically independent absolute invariants [Mae90, Th. B, p. 631] which generate $\mathrm{Frac}(\mathcal{I}_8)$. Unfortunately their degrees are too large for practical computations.

Now, following Section 1.4, we represent the coarse moduli space of genus 3 hyperelliptic curves by the projective variety given by Eq. (1) defined in a weighted projective space of dimension 9 with weights 2, 3,..., 10 the points of which are of the form $(J_2 : J_3 : \ldots : J_{10})$. In this setting, enumerating points on the moduli space over a finite field can be done as follows.

(1) Enumerate representatives for all the points of the weighted projective subspace of dimension 6 defined by the algebraically independent invariants J_2, J_3, ... J_7 (*cf.* Section 1.4).
(2) For each such representative, denoted $(j_2, j_3, \ldots, j_7)$, compute the gcd δ of its support $\{d : j_d \neq 0\}$.
(3) For each representative π of the quotient $k^*/(k^*)^\delta$ (we choose $\pi = 1$ when $\delta = 1$), compute the roots in J_8 of Eq. (2) specialized at $J_d = \pi^{(d/\delta)} j_d$ for $2 \leq d \leq 7$.
(4) For each root j_8, solve Eq. (1) in j_9 and j_{10}.
(5) Return representatives for $\{(\pi^{(2/\delta)} j_2 : \pi^{(3/\delta)} j_3 : \ldots : \pi^{(7/\delta)} j_7 : j_8 : j_9 : j_{10})\}$.

We consider in this algorithm several representatives $(\pi^{(2/\delta)} j_2 : \pi^{(3/\delta)} j_3 : \ldots : \pi^{(7/\delta)} j_7)$ starting from the same class $(j_2 : j_3 : \ldots : j_7)$ when $\delta \neq 1$, even if we may, so, encounter several times the same class at the very end of the enumeration. We do so because such collisions are straightforward to filter, and otherwise we may miss points on the coarse moduli space.

For instance, modulo 11, the representative $j_2 = -1$, $j_3 = j_4 = j_5 = j_6 = j_7 = 0$ yields only one point in the moduli space, *i.e* $(-1 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 0)$, while another choice of representative for the class $(-1 : 0 : \ldots : 0)$, as for instance $j_2 = 1$, $j_3 = j_4 = j_5 = j_6 = j_7 = 0$, yields two points in the moduli space, among which the (new) point $(1 : 0 : 0 : 0 : 0 : 0 : 8 : 2 : 7)$. This is mostly due to the fact that there exists equivalent representatives, here $(-1, 0, 0, 0, 0, 0)$ and $(1, 0, 0, 0, 0, 0)$, for some classes $(j_2 : j_3 : \ldots : j_7)$, here $(-1 : 0 : 0 : 0 : 0 : 0)$, linked by a projective constant λ which may only be defined in a non-trivial extension of $\mathbb{F}_{11}$ of degree δ (in the example $\delta = 2$). In this situation, specializing Eq. (1) with such representatives yield solutions which are no more necessarily equivalent.

Some of the classes encountered while enumerating may be specializations of Shioda invariants at forms f which do not have simple roots. These classes are not points of the coarse moduli space. To discriminate them, we may check that the discriminant is non-zero for forms in the class $(j_2 : j_3 : \ldots : j_{10})$. The discriminant is an invariant Δ, of degree 14, and the test can thus be easily done once one has expressed Δ as a polynomial in the J_is which is

$$\begin{aligned}
&351157512403353600\, J_2\, J_3\, J_4\, J_5 - 106232524760678400\, {J_7}^2 + 369994358063104\, {J_2}^7 \\
&- 66189456176578560\, {J_2}^3 J_3\, J_5 + 33474799101542400\, {J_2}^2 {J_3}^2 J_4 - 1292776756346880000\, {J_2}^2 J_3\, J_7 \\
&+ 1267160326927810560\, {J_2}^2 J_4\, J_6 - 1143872189796188160\, J_2\, {J_3}^2 J_6 - 16397790463918080\, {J_2}^5 J_4 \\
&+ 10245839137013760\, {J_2}^4 {J_3}^2 - 93857730923593728\, {J_2}^4 J_6 + 313290675380551680\, {J_2}^3 {J_4}^2 \\
&+ 2345327219866337280\, {J_2}^3 J_8 - 895451656628551680\, {J_2}^2 {J_5}^2 + 1834618840311398400 0\, {J_2}^2 J_{10} \\
&+ 31476303632793600\, J_2\, {J_3}^4 + 968390964336918528\, J_2\, {J_4}^3 + 4879526536497070080\, J_2\, {J_6}^2 \\
&+ 212465049521356800\, {J_3}^3 J_5 + 169419330714009600\, {J_3}^2 {J_4}^2 + 2549580594256281600\, {J_3}^2 J_8 \\
&+ 2751208802098348032\, {J_4}^2 J_6 + 29008561427982581760\, J_4\, J_{10} - 21756421070986936320\, J_5\, J_9 \\
&- 3021725148748185600\, J_6\, J_8 - 7554312871870464000\, J_2\, J_3\, J_9 + 31803657190574653440\, J_2\, J_4\, J_8 \\
&- 8849169312564510720\, J_2\, J_5\, J_7 - 5139293475644375040\, J_3\, J_4\, J_7 + 4582949808934748160\, J_3\, J_5\, J_6\,.
\end{aligned} \tag{3}$$

2. Reconstruction of binary forms

Let $n > 2$ be an even positive integer, k be a field of characteristic 0 or greater than n and $K = \bar{k}$.

2.1. **Clebsch's identities.** In the following section we give proofs of the results in the Example 3 of [Mes91]. They are the fundamental tools to reconstruct a generic binary form of even degree n from its invariants as it is explained in *loc. cit.* The classical reference for this section is [Cle72, §103] but we wish to give full proofs using the material of Section 1. Note that all formulae but the last one could also be checked by a computer algebra software.

2.1.1. *Useful identities for three quadratic forms.* Let q_1, q_2, q_3 be three elements of $S^2(V)$, *i.e.* binary homogeneous quadratic forms in x, z over k. We denote

$$q_1^* = (q_2, q_3)_1, \quad q_2^* = (q_3, q_1)_1, \quad q_3^* = (q_1, q_2)_1.$$

Lemma 2.1. *We have the relation*

$$q_1 q_1^* + q_2 q_2^* + q_3 q_3^* = 0.$$

Proof. Let us consider the multilinear map $S^2(V^*)^3 \to S^4(V^*)$, $(q_1, q_2, q_3) \mapsto \sum_{i=1}^3 q_i q_i^*$. By multilinearity, to check the equality to zero, it is enough to verify it on linear powers $q_i = \ell_i^2$. Since $(\ell_j^2, \ell_k^2)_1 = [\ell_j, \ell_k]\ell_j\ell_k$, we get

$$\sum q_i q_i^* = \ell_1\ell_2\ell_3 \cdot ([\ell_2, \ell_3]\ell_1 + [\ell_3, \ell_1]\ell_2 + [\ell_1, \ell_2]\ell_3).$$

Of course $w := {}^t([\ell_2, \ell_3], [\ell_3, \ell_1], [\ell_1, \ell_2]) = {}^t(\alpha_1, \alpha_2, \alpha_3) \wedge {}^t(\beta_1, \beta_2, \beta_3)$, so

$$w \cdot {}^t(\alpha_1 x + \beta_1 z, \alpha_2 x + \beta_2 z, \alpha_3 x + \beta_3 z) = 0.$$

This proves the equality to 0 (this is called the first fundamental identity in [GY03, p.19]). □

Lemma 2.2. *Let $R(q_1, q_2, q_3)$ be the determinant of q_1, q_2, q_3 in the basis x^2, xz, z^2. Then*

$$-4 \cdot R(q_1, q_2, q_3) = (\Omega_{12} \cdot \Omega_{23} \cdot \Omega_{31})(q_1(x_1, z_1) \cdot q_2(x_2, z_2) \cdot q_3(x_3, z_3)).$$

Proof. The operator $\Omega_{12} \cdot \Omega_{23} \cdot \Omega_{31} : S^2(V^*)^3 \to k$ is multi-linear and anti-symmetric. Hence it is equal to $R(q_1, q_2, q_3)$ up to a constant. □

Lemma 2.3. *Let $A_{ij} = (q_i, q_j)_2$ for $i, j \in \{1, 2, 3\}$. Then*

$$2 \cdot \det((A_{ij})_{i,j=1,2,3}) = R^2(q_1, q_2, q_3).$$

Proof. We follow [GY03, p.79]. Let us consider the multilinear form $\phi : S^2(V^*)^6 \to k$ given by

$$(q_1, q_2, q_3, q_4, q_5, q_6) \mapsto \det \begin{bmatrix} (q_1, q_4)_2 & (q_1, q_5)_2 & (q_1, q_6)_2 \\ (q_2, q_4)_2 & (q_2, q_5)_2 & (q_2, q_6)_2 \\ (q_3, q_4)_2 & (q_3, q_5)_2 & (q_3, q_6)_2 \end{bmatrix}.$$

If we evaluate ϕ on the elements of the basis $q_i = \ell_i^2$, since $(q_i, q_j)_2 = [\ell_i, \ell_j]^2 = (\alpha_i\beta_j - \alpha_j\beta_i)^2$, we get

$$\begin{aligned} \phi(\ell_1^2, \ldots, \ell_6^2) &= \det \begin{bmatrix} \alpha_1^2 & \alpha_1\beta_1 & \beta_1^2 \\ \alpha_2^2 & \alpha_2\beta_2 & \beta_2^2 \\ \alpha_3^2 & \alpha_3\beta_3 & \beta_3^2 \end{bmatrix} \cdot \det \begin{bmatrix} \beta_4^2 & \beta_5^2 & \beta_6^2 \\ -2\alpha_4\beta_4 & -2\alpha_5\beta_5 & -2\alpha_6\beta_6 \\ \alpha_4^2 & \alpha_5^2 & \alpha_6^2 \end{bmatrix} \\ &= \frac{1}{2} \cdot R(q_1, q_2, q_3) \cdot R(q_4, q_5, q_6) \end{aligned}$$

We conclude by letting $q_4 = q_1, q_5 = q_2$ and $q_6 = q_3$. □

Lemma 2.4. *Let f be any quadratic form. Then*

$$\frac{1}{2} \cdot R(q_1, q_2, q_3) \cdot f = (f, q_1^*)_2 \cdot q_1 + (f, q_2^*)_2 \cdot q_2 + (f, q_3^*)_2 \cdot q_3 \,.$$

Proof. Let us consider the multi-linear map $\psi : S^2(V^*)^4 \to S^2(V^*)$ given by the difference of the left and right member of the equality above. We want to prove that ψ is the zero map and again it is enough to do this on the basis $q_i = \ell_i^2$ and $f = \ell^2 = (\mu x + \nu z)^2$. We proceed like in [Cle72, p.414]. Obviously

$$\det \begin{bmatrix} \ell_1^2 & \alpha_1^2 & \alpha_1\beta_1 & \beta_1^2 \\ \ell_2^2 & \alpha_2^2 & \alpha_2\beta_2 & \beta_2^2 \\ \ell_3^2 & \alpha_3^2 & \alpha_3\beta_3 & \beta_3^2 \\ f & \mu^2 & \mu\nu & \nu^2 \end{bmatrix} = 0.$$

Expanding the determinant with respect to the first column, we get

$$-fR(q_1, q_2, q_3) + q_3R(q_1, q_2, f) - q_2R(q_1, q_3, f) + q_1R(q_2, q_3, f) = 0.$$

To get our result, we only need to prove that $R(q_i, q_j, f) = 2 \cdot (f, (q_i, q_j)_1)_2$. By a direct computation, $R(q_i, q_j, f) = 2[\ell_i, \ell_j] \cdot [\ell, \ell_j] \cdot [\ell, \ell_i]$. On the other hand,

$$\begin{aligned} (f, (q_i, q_j)_1)_2 &= (f, [\ell_i, \ell_j] \cdot \ell_i\ell_j)_2 = [\ell_i, \ell_j] \cdot (\ell^2, \ell_i\ell_j)_2 \,, \\ &= \frac{1}{4}[\ell_i, \ell_j] \cdot \Omega_{12}^2 \left[\ell(x_1, z_1)^2 \ell_i(x_2, z_2)\ell_j(x_2, z_2)\right] \,, \\ &= [\ell_i, \ell_j] \cdot [\ell, \ell_j] \cdot [\ell, \ell_i] \,. \end{aligned}$$

□

Mestre's formula is somehow dual to this result and can be checked using the same arguments.

Lemma 2.5. *Let f be any quadratic form. Then*

$$R(q_1,q_2,q_3)\cdot f = 2\left((f,q_1)_2\cdot q_1^* + (f,q_2)_2\cdot q_2^* + (f,q_3)_2\cdot q_3^*\right).$$

2.1.2. *Relations among a binary form and three quadratic forms.* We now come to the two fundamental results.

Proposition 2.6. *We have $\sum_{i,j} A_{ij}\cdot q_i^*\cdot q_j^* = 0$.*

Proof. For $i\in\{1,2,3\}$ using Lemma 2.5, we get

$$R(q_1,q_2,q_3)q_i = (q_i,q_1)_2 q_1^* + (q_i,q_2)_2 q_2^* + (q_i,q_3)_2 q_3^*.$$

We multiply the previous equality by q_i^*,

$$R(q_1,q_2,q_3)q_iq_i^* = A_{i1}q_i^*q_1^* + A_{i2}q_i^*q_2^* + A_{i3}q_i^*q_3^*,$$

and we sum on $1\le i\le 3$,

$$R(q_1,q_2,q_3)\cdot\sum_{i=1}^3 q_iq_i^* = \sum_{i=1}^3 A_{i1}q_i^*q_1^* + A_{i2}q_i^*q_2^* + A_{i3}q_i^*q_3^* = \sum_{1\le i,j\le 3} A_{ij}q_i^*q_j^*,$$

which is zero by Lemma 2.1. □

Proposition 2.7. *Let f be a binary form of even degree n. Then*

$$R(q_1,q_2,q_3)^{n/2}\cdot f(x,z) = \frac{1}{n!}\cdot\left(\sum_{i=1}^3 q_i^*(x,z)\delta_i\right)^{n/2} f(x_1,z_1)$$

where δ_i is the differential operator $\phi(x_1,z_1)\mapsto\Omega_{12}^2(\phi(x_1,z_1)\cdot q_i(x_2,z_2))$.

Proof. As the right member is linear in f we can assume that $f=\ell^n=(\mu x_1+\nu z_1)^n$. To make it linear in the q_i we consider $n/2$ triplets of quadratic forms (q_{1j},q_{2j},q_{3j}) and define accordingly

$$q_{1j}^* = (q_{2j},q_{3j})_1,\quad q_{2j}^* = (q_{3j},q_{1j})_1,\quad q_{3j}^* = (q_{1j},q_{2j})_1.$$

We replace the previous operator by

$$\Psi := \frac{1}{n!}\cdot\prod_{j=1}^{n/2}\sum_{i=1}^3 q_{ij}^*(x,z)\,\delta_{ij}$$

where

$$\delta_{ij}(\phi(x_1,z_1)) = \Omega_{12}^2(\phi(x_1,z_1)q_{ij}(x_2,z_2)).$$

As Ψ is linear in each q_{ij} we can assume that

$$q_{ij} = \ell_{ij}^2 = (\alpha_{ij}x+\beta_{ij}z)^2.$$

For $1\le j\le n/2$, let $i_j\in\{1,2,3\}$. Observe that for all m

$$\delta_{i_jj}(\ell^m) = \Omega_{12}^2(\ell^m(x_1,z_1)q_{i_jj}(x_2,z_2)) = \frac{2!\;m!}{(m-2)!}\cdot[\ell,\ell_{i_jj}]\cdot\ell^{m-2}.$$

Hence

$$\frac{1}{n!}\left(\prod_{j=1}^{n/2}\delta_{i_jj}\right)(f) = \prod_{j=1}^{n/2}2[\ell,\ell_{i_jj}]^2 = \prod_{j=1}^{n/2}2(\ell^2,q_{i_j,j})_2.$$

Therefore, if we develop the expression of Ψ, we can replace each product of the δ_{i_jj} operators on f by the right expression. Re-factoring the new expression, we get that

$$\Psi(f) = \prod_{j=1}^{n/2}2\sum_{i=1}^3 q_{ij}^*\cdot(\ell^2,q_{ij})_2.$$

Using Lemma 2.5,

$$\Psi(f) = \prod_{j=1}^{n/2} \left(R(q_{1j}, q_{2j}, q_{3j}) \cdot \ell^2\right) = \left(\prod_{j=1}^{n/2} R(q_{1j}, q_{2j}, q_{3j})\right) \cdot \ell^n.$$

To conclude, we let $q_{ij} = q_i$ for all j.

□

2.2. **A generic reconstruction algorithm.** Starting from a weighted projective point $(\iota_1 : \iota_2 : \ldots)$ of values in k for a finite set of generators $\{I_i\}$ of $\mathcal{I}_{2g+2}$, we aim at recovering a hyperelliptic curve $C/K : y^2 = f(x)$ such that $(\iota_1 : \iota_2 : \ldots) = (I_1(f) : I_2(f) : \ldots)$. Inverting the polynomial system giving the invariants in terms of a generic polynomial f can be efficiently done only in very specific cases (see Section 3.2). However, Mestre explained in his Example 3 and Remark p. 321 [Mes91] that for a generic hyperelliptic curves, one can use the background developed in Section 2.1 in the following way.

Let $f(X, Z)$ be a binary form of even degree $2g+2$ defined over K. Using transvectants one can compute three covariants $q_i(x, z) = q_i(f, (x, z))$ of order 2 defined over K. With the notation of Section 2.1, we denote

$$q_1^* = (q_2, q_3)_1, \quad q_2^* = (q_3, q_1)_1, \quad q_3^* = (q_1, q_2)_1.$$

From Propositions 2.6 and 2.7, we see that the point $P = (x_1^* : x_2^* : x_3^*) = (q_1^*(x, z) : q_2^*(x, z) : q_3^*(x, z))$ is solution of

$$\begin{cases} \sum A_{ij} x_i x_j = 0, \\ \sum_I h_I x_I - R(q_1, q_2, q_3)^{n/2} f(x, z) = 0, \end{cases}$$

where I is a multi-index of cardinality $n/2$.

Clearly the $A_{ij} = (q_i, q_j)_2$ are invariants. In Proposition 2.7, each time the operator Ω_{ij}^2 is applied on a covariant of order r and on a covariant q_i of order 2, we get a covariant of order $r + 2 - 4 = r - 2$. As we started with f which is a covariant of order n and we applied the operators $n/2$ times, the coefficients h_I are invariants. Hence, both A_{ij} and h_I being invariants, they can be express as polynomials in the I_is with coefficients in the prime field of K. Thus if the invariants of a given $f(X, Z)$ lives inside a field $k \subset K$, the specializations of the A_{ij} and h_I belongs to k as well. We refer to Section 2.3 for a strategy to find the formal expressions when $n = 8$.

Now, assume that $R(q_1, q_2, q_3) \neq 0$. Note that this is an invariant as well, and so it can be computed as a polynomial in the I_is. Then P lies on the curve $\mathcal{H}/k : \sum_I h_I x_I = 0$ if and only if $(x : z)$ is a root of $f(x, z)$. Moreover, by Lemma 2.3, the conic $\mathcal{Q}/k : \sum A_{ij} x_i x_j = 0$ is non singular, hence the morphism $(x : z) \mapsto (x_1^* : x_2^* : x_3^*) \in \mathcal{Q}$ is a parametrization. As there a quadratic extension of k, we get a parametrization $\phi : (x, z) \mapsto (\chi_1 : \chi_2 : \chi_3)$. All parametrizations being GL_2-equivalent, the binary polynomial $\sum_I h_I \chi_I$ is GL_2-equivalent to f. Hence we get the following proposition.

Proposition 2.8. *Let (q_1, q_2, q_3) be three covariants of order 2 of a binary form f of even degree n defined over K. If $R(q_1, q_2, q_3) \neq 0$, there exists a non-singular conic $\mathcal{Q}$ and a plane curve $\mathcal{H}$ of degree $n/2$ defined over a subfield k of K such that there is a K-isomorphism $\mathcal{Q} \to \mathbb{P}^1$ mapping the intersections points of $\mathcal{Q} \cap \mathcal{H}$ on the roots of $f(X, Z)$. Moreover, this isomorphism is defined at most over a quadratic extension of k and is defined over k as soon as $\mathcal{Q}$ has a k-rational point.*

This is Mestre's main result in [Mes91, Sec.2.4], that for a generic hyperelliptic curve, the obstruction to reconstruct the curve from its invariant over k is equivalent to $\mathcal{Q}(k) = \emptyset$. We will come back on these questions in Section 4.1 and relate k' to the field of moduli.

To conclude, given values $(\iota_1 : \iota_2 : \ldots)$ defined in k, the general method is as follows.

(1) Find a triple (q_1, q_2, q_3) of covariants of order 2 such that that the expression of $R(q_1, q_2, q_3)$ as a polynomial in the I_is evaluated at the $\iota_i s$ is non-zero;
(2) Find a point on the conic $\mathcal{Q}$ defined by the expression of the coefficients $A_{ij}(q_1, q_2, q_3)$ as polynomials in the I_is evaluated at the $\iota_i s$, and get a parametrization for the conic.
(3) Let $\mathcal{H}$ be the curve defined by the expression of the coefficients $h_I(q_1, q_2, q_3)$ as polynomials in the I_i evaluated at the ι_is. After a parametrization of the conic $\mathcal{Q}$ return the intersection divisor $\mathcal{Q} \cdot \mathcal{H}$ as a polynomial $f(X, Z)$ over at most a quadratic extension of k.

It might happen that all possible $R(q_1, q_2, q_3)$ evaluated at the $\iota_i s$ are zero. Even if it is not the case and for a general field K, $\mathcal{Q}$ might have no k-rational point although we know, for theoretical reasons, we can reconstruct the curve over k. Section 3.2 and Section 4 will deal with these issues.

Ord. Deg.	0	2	4	6	8	10	12	14	18	Tot
1	–	–	–	–	f	–	–	–	–	1
2	$(\mathrm{f},\mathrm{f})_8$	–	$(\mathrm{f},\mathrm{f})_6$	–	$(\mathrm{f},\mathrm{f})_4$	–	$(\mathrm{f},\mathrm{f})_2$	–	–	4
3	$(C_{2,8},\mathrm{f})_8$	–	$(C_{2,8},\mathrm{f})_6$	$(C_{2,8},\mathrm{f})_5$	$(C_{2,8},\mathrm{f})_4$	$(C_{2,8},\mathrm{f})_3$	$(C_{2,8},\mathrm{f})_2$	$(C_{2,8},\mathrm{f})_1$	$(C_{2,12},\mathrm{f})_1$	8
4	$(C_{3,8},\mathrm{f})_8$	–	$(C_{3,4},\mathrm{f})_4$ $(C_{3,8},\mathrm{f})_6$	$(C_{3,4},\mathrm{f})_3$	$(C_{3,4},\mathrm{f})_2$	$(C_{3,4},\mathrm{f})_1$ $(C_{3,8},\mathrm{f})_3$	$(C_{3,8},\mathrm{f})_2$	$(C_{3,8},\mathrm{f})_1$	$(C_{3,12},\mathrm{f})_1$	10
5	$(C_{4,8},\mathrm{f})_8$	$(C_{4,10},\mathrm{f})_8$	$(C_{4,10},\mathrm{f})_7$ $(C_{4,8},\mathrm{f})_6$	$(C_{4,10},\mathrm{f})_6$ $(C_{4,8},\mathrm{f})_5$	$(C_{4,10},\mathrm{f})_5$	$(C_{4,8},\mathrm{f})_3$ $(C_{4,10},\mathrm{f})_4$ $(C'_{4,10},\mathrm{f})_4$	–	$(C_{4,10},\mathrm{f})_2$	–	11
6	$(C_{3,4}\,C_{2,4},\mathrm{f})_8$	$(C_{5,8},\mathrm{f})_7$	$(C_{5,8},\mathrm{f})_6$ $(C'_{5,4},\mathrm{f})_4$	$(C_{5,8},\mathrm{f})_5$ $(C'_{5,4},\mathrm{f})_3$ $(C'_{5,10},\mathrm{f})_6$	$(C'_{5,4},\mathrm{f})_2$	$(C'_{5,4},\mathrm{f})_1$	–	–	–	9
7	$(C_{2,4}\,C'_{4,4},\mathrm{f})_8$	$(C_{2,4}\,C_{4,6},\mathrm{f})_8$ $(C''_{6,6},\mathrm{f})_6$	$(C_{2,4}\,C_{4,6},\mathrm{f})_7$ $(C''_{6,6},\mathrm{f})_5$	$(C''_{6,6},\mathrm{f})_4$ $(C_{6,2},\mathrm{f})_2$ $(C_{2,4}\,C_{4,6},\mathrm{f})_6$	–	–	–	–	–	8
8	$(C_{3,4}\,C_{4,4},\mathrm{f})_8$	$(C_{2,8}\,C_{5,2},\mathrm{f})_8$ $(C_{3,6}\,C_{4,4},\mathrm{f})_8$	$(C_{3,6}\,C_{4,4},\mathrm{f})_7$ $(C_{3,4}\,C_{4,6},\mathrm{f})_7$	$(C_{3,6}\,C_{4,4},\mathrm{f})_6$ $(C_{3,4}\,C_{4,6},\mathrm{f})_6$	–	–	–	–	–	7
9	$(C_{2,4}\,C_{6,4},\mathrm{f})_8$	$(C_{4,6}\,C'_{4,4},\mathrm{f})_8$ $(C_{2,4}\,C_{6,4},\mathrm{f})_7$ $(C_{2,4}\,C'_{6,6},\mathrm{f})_8$	$(C_{2,4}\,C_{6,4},\mathrm{f})_6$	–	–	–	–	–	–	5
10	$(C_{4,4}\,C'_{5,4},\mathrm{f})_8$	$(C'_{7,2}\,C_{2,4},\mathrm{f})_6$ $(C_{4,6}\,C_{5,4},\mathrm{f})_8$	–	–	–	–	–	–	–	3
11	–	$(C'_{8,4}\,C_{2,4},\mathrm{f})_7$ $(C'_{5,6}\,C'_{5,4},\mathrm{f})_8$	–	–	–	–	–	–	–	2
12	–	$(C'_{6,6}\,C'_{5,4},\mathrm{f})_8$	–	–	–	–	–	–	–	1
Tot	9	14	13	12	6	7	3	3	2	69

TABLE 2. Fundamental invariants and covariants for a binary form of degree 8

2.3. **Reconstruction in the hyperelliptic genus 3 case.** We have computed a system of fundamental generators for invariants and covariants of octics, using Gordan's algorithm. Results are given in Table 2. In this table, generators are all defined by the mean of transvectants of the form $(\prod_{d,r} C_{d,r}, \mathfrak{f})_h$ where we denote recursively by $C_{d,r}$ generators of degree d and order r given at the intersection of the row d and the column r of the table. When we have two or three generators of degree d and r, we denote them by $C'_{d,r}$ and $C''_{d,r}$. For instance, $C_{5,10} = (C_{4,8}, \mathfrak{f})_3$, $C'_{5,10} = (C_{4,10}, \mathfrak{f})_4$ and $C''_{5,10} = (C'_{4,10}, \mathfrak{f})_4$. Covariants $C_{2,0}$, $C_{3,0}$, …, $C_{10,0}$ given in column 0 are invariants and so, they are related to Shioda invariants J_2, J_3, …, J_{10} by

$$
\begin{aligned}
&C_{2,0} = J_2\,,\ C_{3,0} = J_3\,,\ C_{4,0} = \frac{1}{30}\,{J_2}^2 - \frac{4}{35}\,J_4\,,\ C_{5,0} = \frac{3}{14}\,J_5\,,\\
&C_{6,0} = -\frac{1}{1890}\,{J_2}^3 + \frac{1}{15}\,J_2\,J_4 + \frac{1}{63}\,{J_3}^2 + \frac{8}{21}\,J_6\,,\\
&C_{7,0} = \frac{1}{42}\,J_2\,J_5 - \frac{3}{245}\,J_3\,J_4 + \frac{1}{49}\,J_7\,,\\
&C_{8,0} = -\frac{11}{264600}\,{J_2}^4 + \frac{59}{14700}\,{J_2}^2 J_4 + \frac{11}{8820}\,J_2\,{J_3}^2 + \frac{22}{735}\,J_2\,J_6 + \frac{3}{980}\,J_3\,J_5 + \frac{198}{8575}\,{J_4}^2 + \frac{9}{49}\,J_8\,,\\
&C_{9,0} = -\frac{1}{45360}\,{J_2}^3 J_3 + \frac{1}{1260}\,{J_2}^2 J_5 + \frac{61}{17640}\,J_2\,J_3\,J_4 - \frac{17}{4704}\,J_2\,J_7 + \frac{1}{1512}\,{J_3}^3 + \frac{1}{144}\,J_3\,J_6 - \frac{1}{112}\,J_4\,J_5 - \frac{1}{28}\,J_9\,,\\
&C_{10,0} = -\frac{1}{595350}\,{J_2}^5 + \frac{107}{411600}\,{J_2}^3 J_4 + \frac{1}{19845}\,{J_2}^2 {J_3}^2 + \frac{317}{105840}\,{J_2}^2 J_6 + \frac{5}{2352}\,J_2\,J_3\,J_5\\
&\qquad + \frac{293}{180075}\,J_2\,{J_4}^2 + \frac{25}{5488}\,J_2\,J_8 + \frac{5}{6174}\,{J_3}^2 J_4 + \frac{17}{2744}\,J_3\,J_7 + \frac{19}{5145}\,J_4\,J_6 - \frac{153}{27440}\,{J_5}^2 + \frac{3}{49}\,J_{10}\,.
\end{aligned}
$$

Our main motivation for computing Table 2 is to determine fundamental covariants of order 2 which can be used as quadratic forms in Proposition 2.8 to reconstruct a generic binary octic from its Shioda invariants. As with Shioda invariants, we paid attention that none of the transvectant computations in Table 2 involves covariants of order greater than 10 (except the 2 covariants of order 18, but it does not matter since these 2 covariants are not useful to compute other invariants or covariants of order 2). These formulas are thus also valid, in addition to fields of characteristic 0, over any field of or characteristic greater or equal than 11.

The main computational difficulty in the reconstruction method of Section 2.2 is to write the invariants A_{ij} and h_I as polynomials in the J_i's, since their degree may be large (close to 40 in our cases). Writing them as a polynomial with 9 unknowns a_i's for a generic form $f = a_8X^8 + a_7X^7 + \ldots + a_0$ is hopeless. We follow instead a "black-box" approach.

Algorithm 2: Write an invariant as a polynomial in the J_i's.

Input : An invariant I of degree d (given as an evaluation program).
Output: A polynomial P in $\mathbb{Q}[J_2, \ldots, J_{10}]$ such that $I(f) = P(J_2(f), J_3(f), \ldots, J_{10}(f))$.

`// A basis for the polynomials of degree` d `in the weighted graded algebra` $\mathbb{Q}[J_2, J_3, \ldots, J_{10}]$
$\mathcal{B} \leftarrow \left[\prod_w J_w^{e_w}$ `s.t.` $\sum_w w\,e_w = d\right]$;

`// Choose at random` $\#\mathcal{B} + O(1)$ `octics over` $\mathbb{Q}$
$\mathcal{F} \leftarrow \left[a_8X^8 + \ldots + a_1X\,Z^7 + a_0Z^8$ `for` $\#\mathcal{B} + O(1)$ $9-$`uples` $(a_0, a_1, \ldots, a_8)$ `randomly chosen in` $\mathbb{Q}^9\right]$;

`// Evaluate the invariant` I `and the basis` $\mathcal{B}$ `at each form of` $\mathcal{F}$
for $i = 1$ **to** $\#\mathcal{F}$ **do**
    $V_i \leftarrow I(\mathcal{F}_i)$;
    for $j = 1$ **to** $\#\mathcal{B}$ **do**
        $M_{i,j} \leftarrow \mathcal{B}_j(\mathcal{F}_i)$

`// Invert the linear system defined by the matrix` M `and the vector` V
Find the vectors U such that $M \times U = V$;

return $\sum_i U_i\,\mathcal{B}_i$ for each U

More precisely, since invariants are computed through sequences of covariants which are the result of transvectant or differential operations, we represent a covariant no more by a formal expression in the a_i's but as an algorithm which performs the corresponding sequence of operations. If one inputs a generic form $f \in \mathbb{Q}[a_0, \ldots, a_8][X, Z]$, such an algorithm returns the formal expression of the covariant as a multivariate homogeneous polynomial in the a_is, X and Z. But, if one inputs a form $f \in \mathbb{Q}[X, Z]$, the algorithm returns

the covariant as a homogeneous polynomial in $\mathbb{Q}[x,z]$, without the use of the formal expression in the a_is. In other words, we consider that a covariant is given by an evaluation program. Note that it is immediate to determine the degree and the order of a covariant from the sequence of operations which compose its evaluation program.

Now coming back to the question of writing a homogeneous invariant as a polynomial in the J_is, we propose to construct a basis $\mathcal{B} = \{ \prod_w J_w^{e_w} \ : \ \sum_w w\, e_w = d \}$ for the polynomials of degree d in the weighted graded algebra $\mathbb{Q}[J_2, J_3, \ldots, J_{10}]$ and we evaluate this basis (with some given evaluation programs for the J_i's), and I (given as an evaluation program too), at $\#\mathcal{B} + O(1)$ octics chosen at random over $\mathbb{Q}$. It remains to invert the corresponding linear system to find I as a polynomial in the J_i's. Algorithm 2 summarizes this method.

Remark 2.9. Working with $\mathbb{Q}[J_2, J_3, \ldots, J_{10}]$ instead of $\mathbb{Q}[a_0, a_1, \ldots, a_8]$ in Algorithm 2 makes a real difference in practice. For instance, an invariant of degree 20 yields a basis $\mathcal{B}$ with only 107 monomials. This must be compared to the 61731 monomials that otherwise we would have to deal with in the weighted projective points $(a_0 : a_1 : \ldots, a_8)$ where a_0, a_1, …, a_8 are of weight 0, 1, …, 8. Note that an invariant of degree d is of weight $4d$ in the last algebra. For instance, $J_2 = 2\,a_0\,a_8 - a_1\,a_7/4 + a_2\,a_6/14 - a_3\,a_5/28 + {a_4}^2/70$ is of degree 2 and weight 8.

Example 2.10. Consider the covariants of order 2 of smallest degree in Table 2, that is $q_1 = C_{5,2}$, $q_2 = C_{6,2}$ and $q_3 = C_{7,2}$. A first call to Algorithm 2 yields that $R = R(q_1, q_2, q_3)$ is equal, up to a constant, to

$$
\begin{aligned}
R = {} & -4937630140800\ {J_9}^2 + 6172588800000\ J_8\ J_{10} + 1016336160000\ {J_6}^3 - 1646487542700\ J_5\ J_6\ J_7 + 475344450\ {J_5}^2 J_8 - 13778100\ J_4\ {J_7}^2 \\
& + 6154254741600\ J_4\ J_6\ J_8 + 2469123699840\ J_4\ J_5\ J_9 - 3175414824960\ {J_4}^2 J_{10} - 1028718873000\ J_3\ J_7\ J_8 - 1555231104000\ J_3\ J_6\ J_9 \\
& + 514676332800\ J_3\ J_5\ J_{10} - 579162433500\ J_2\ {J_8}^2 + 231655788000\ J_2\ J_7\ J_9 + 47632860\ {J_4}^2 {J_5}^2 - 201602675520\ {J_4}^3 J_6 \\
& - 264617457390\ J_3\ J_4\ J_5\ J_6 + 529262244990\ J_3\ {J_4}^2 J_7 + 4618063800\ {J_3}^2 {J_6}^2 - 35210700\ {J_3}^2 J_5\ J_7 - 228766979700\ {J_3}^2 J_4\ J_8 \\
& + 38124172800\ {J_3}^3 J_9 + 77149935135\ J_2\ {J_5}^2 J_6 - 40603006080\ J_2\ J_4\ {J_6}^2 - 115812049185\ J_2\ J_4\ J_5\ J_7 - 330859026540\ J_2\ {J_4}^2 J_8 \\
& + 145802916000\ J_2\ J_3\ J_6\ J_7 - 15715198800\ J_2\ J_3\ J_4\ J_9 + 42877447200\ J_2\ {J_3}^2 J_{10} + 53596043550\ {J_2}^2 {J_7}^2 - 145802916000\ {J_2}^2 J_6\ J_8 \\
& - 53606606760\ {J_2}^2 J_5\ J_9 + 137217628800\ {J_2}^2 J_4\ J_{10} - 36737464140\ {J_3}^2 {J_4}^3 - 7824600\ {J_3}^3 J_4\ J_5 + 11300902200\ {J_3}^4 J_6 - 47249726760\ J_2\ {J_4}^4 \\
& - 12161979900\ J_2\ J_3\ {J_4}^2 J_5 + 33446455740\ J_2\ {J_3}^2 J_4\ J_6 + 1760535\ {J_2}^2 J_4\ {J_5}^2 + 25514097660\ {J_2}^2 {J_4}^2 J_6 - 153935460\ {J_2}^3 {J_6}^2 + 1173690\ {J_2}^3 J_5\ J_7 \\
& + 7625565990\ {J_2}^3 J_4\ J_8 - 1270805760\ {J_2}^3 J_3\ J_9 - 1429248240\ {J_2}^4 J_{10} + 289800\ J_2\ {J_3}^4 J_4 + 900887400\ {J_2}^2 {J_3}^2 {J_4}^2 + 2575261188\ {J_2}^3 {J_4}^3 \\
& + 260820\ {J_2}^3 J_3\ J_4\ J_5 - 753393480\ {J_2}^3 {J_3}^2 J_6 - 1114881858\ {J_2}^4 J_4\ J_6 - 19320\ {J_2}^4 {J_3}^2 J_4 - 30029580\ {J_2}^5 {J_4}^2 + 12556558\ {J_2}^6 J_6 + 322\ {J_2}^7 J_4\,.
\end{aligned}
$$

Similarly, we find that the equation of the conic, $\sum_{i,j} A_{ij}\, x_i x_j = 0$, is equal to

$$
\begin{aligned}
& (9217732608000\ J_{10} - 1422489600\ {J_2}^3 J_4 + 1814283878400\ J_4\ J_6 - 384072192000\ J_3\ J_7 \\
& \quad + 42674688000\ {J_3}^2 J_4 - 1152216576000\ {J_5}^2 + 212154163200\ J_2\ {J_4}^2 + 384072192000\ J_2\ J_8)\ {x_1}^2 + \\
& (-80015040000\ {J_3}^2 J_5 + 2667168000\ {J_2}^3 J_5 - 12002256000\ {J_2}^2 J_7 + 288054144000\ J_2\ J_9 + 216040608000\ J_4\ J_7 \\
& \quad - 102019176000\ J_2\ J_4\ J_5 + 138883248000\ J_3\ {J_4}^2 - 48009024000\ J_5\ J_6 + 360067680000\ J_3\ J_8)\ x_1\, x_2 + \\
& (-12040358400\ {J_2}^2 J_8 - 902039040\ {J_2}^3 J_6 - 24768737280\ {J_4}^3 + 27061171200\ {J_3}^2 J_6 + 18627840\ {J_2}^4 J_4 - 424308326400\ J_2\ J_{10} \\
& \quad - 5482391040\ {J_2}^2 {J_4}^2 - 43481733120\ J_2\ J_4\ J_6 + 216726451200\ J_5\ J_7 + 12040358400\ J_2\ J_3\ J_7 - 762657638400\ J_4\ J_8 \\
& \quad + 36121075200\ J_2\ {J_5}^2 + 135339724800\ J_3\ J_9 - 162570240000\ {J_6}^2 - 10516262400\ J_3\ J_4\ J_5 - 558835200\ J_2\ {J_3}^2 J_4)\ x_1\, x_3 + \\
& (135025380000\ J_3\ J_9 + 55566000\ {J_2}^3 J_6 - 15788682000\ J_2\ J_4\ J_6 + 2813028750\ {J_2}^2 J_8 - 2813028750\ J_2\ J_3\ J_7 + 149333625\ {J_2}^4 J_4 \\
& \quad + 8439086250\ J_3\ J_4\ J_5 - 151903552500\ J_4\ J_8 - 2509400250\ {J_2}^2 {J_4}^2 + 75951776250\ J_5\ J_7 - 1666980000\ {J_3}^2 J_6 - 4480008750\ J_2\ {J_3}^2 J_4 \\
& \quad + 92610000\ {J_2}^3 {J_3}^2 - 1543500\ {J_2}^6 - 1389150000\ {J_3}^4 - 2893401000\ {J_4}^3 - 50009400000\ {J_6}^2 - 67512690000\ J_2\ J_{10})\ {x_2}^2 + \\
& (1434793500\ {J_2}^2 J_4\ J_5 - 1629217800\ J_2\ J_3\ {J_4}^2 + 6460738200\ J_2\ J_5\ J_6 + 365148000\ {J_3}^3 J_4 - 41806800\ {J_2}^4 J_5 - 12748654800\ J_3\ J_4\ J_6 \\
& \quad + 1254204000\ J_2\ {J_3}^2 J_5 + 914457600\ J_2\ J_4\ J_7 - 12171600\ {J_2}^3 J_3\ J_4 - 172254600\ {J_2}^3 J_7 - 2400451200\ {J_2}^2 J_9 - 714420000\ J_6\ J_7 \\
& \quad - 5643918000\ J_2\ J_3\ J_8 - 4445733600\ {J_4}^2 J_5 + 14402707200\ J_4\ J_9 + 10811556000\ {J_3}^2 J_7 - 63440496000\ J_3\ J_{10} - 44365482000\ J_5\ J_8)\ x_2\, x_3 + \\
& (94363920\ {J_2}^3 J_8 + 2592705024\ {J_4}^2 J_6 - 32568480\ {J_3}^2 {J_4}^2 + 57512\ {J_2}^5 J_4 - 283091760\ {J_2}^2 {J_5}^2 + 4386130560\ {J_2}^2 J_{10} - 1905120000\ J_6\ J_8 \\
& \quad - 40824000\ {J_7}^2 + 34895088\ {J_2}^3 {J_4}^2 + 1886976000\ J_2\ {J_6}^2 - 10150479360\ J_5\ J_9 - 109801152\ {J_2}^2 J_4\ J_6 + 23227223040\ J_4\ J_{10} \\
& \quad + 21819168\ {J_2}^4 J_6 + 3110425920\ J_3\ J_5\ J_6 + 15630965280\ J_2\ J_4\ J_8 + 164838240\ J_2\ J_3\ J_4\ J_5 + 635065920\ J_2\ {J_4}^3 - 3676609440\ J_3\ J_4\ J_7 \\
& \quad - 1725360\ {J_2}^2 {J_3}^2 J_4 - 3397101120\ J_2\ J_5\ J_7 - 2121396480\ J_2\ J_3\ J_9 - 94363920\ {J_2}^2 J_3\ J_7 - 654575040\ J_2\ {J_3}^2 J_6)\ {x_3}^2 = 0\,.
\end{aligned}
$$

The beginning of the quartic $\sum_{i,j,k,l} h_{i,j,k,l} x_i x_j x_k x_l$ (its coefficients are too large to be all written here) is then

$$
\begin{aligned}
& (20832487200\ {J_7}^3 - 98761420800\ J_6\ J_7\ J_8 - 14814213120\ {J_6}^2 J_9 + 140619288600\ J_5\ {J_8}^2 + 21526903440\ J_5\ J_7\ J_9 + 192584770560\ J_4\ J_8\ J_9 \\
& \quad - 29628426240\ J_3\ {J_9}^2 + 6351593875200000\ J_2\ J_9\ J_{10} - 231472080\ {J_5}^3 J_6 + 17310682368\ J_4\ J_5\ {J_6}^2 - 24651776520\ J_4\ {J_5}^2 J_7 \\
& \ldots \\
& + 653457959280\ {J_2}^5 J_5\ J_6 - 653460684660\ {J_2}^5 J_4\ J_7 + 108909756900\ {J_2}^6 J_9 + 47040\ {J_2}^4 {J_3}^3 J_4 + 56723695560\ {J_2}^5 J_3\ {J_4}^2 \\
& \quad + 141120\ {J_2}^5 {J_3}^2 J_5 + 222264\ {J_2}^6 J_4\ J_5 + 117600\ {J_2}^6 J_3\ J_6 + 7056\ {J_2}^7 J_7 - 784\ {J_2}^7 J_3\ J_4 - 2352\ {J_2}^8 J_5)\ {x_1}^4 + \cdots
\end{aligned}
$$

These precomputations done, let us look now for an octic f defined over $\mathbb{F}_{11}$ such that for instance $J_2 = 1$, $J_3 = J_4 = J_5 = J_6 = J_7 = 0$, $J_8 = 8$, $J_9 = 2$ and $J_{10} = 7$. We first check that $R \neq 0$ and that the conic equation is equal to

$$x_1\, x_2 + 3\, x_1\, x_3 + 6\, x_2^2 + x_2\, x_3 + 8\, x_3^2 = 0\,.$$

Then, since the point $(1 : 0 : 1)$ is on this conic, we have the parametrization

$$(X, Z) \mapsto (8\, X^2 + 10\, X\, Z + 6\, Z^2 : 8\, X\, Z + 9\, Z^2 : X\, Z + 6\, Z^2)\,.$$

In this case, the quartic equation of $\mathcal{H}$ is equal to

$$6\,x_1^4 + 5\,x_1^3\,x_2 + 9\,x_1^3\,x_3 + 5\,x_1^{*2}\,x_2^{*2} + x_1^2\,x_2\,x_3 + 7\,x_1^2\,x_3^2 + 8\,x_1\,x_2^3 \\ + 10\,x_1\,x_2^2\,x_3 + 3\,x_1\,x_2\,x_3^2 + 3\,x_1\,x_3^3 + 7\,x_2^4 + 7\,x_2^3\,x_3 + 9\,x_2\,x_3^3 + 5\,x_3^4$$

and we finally find that, up to a constant,

$$f(X,Z) = 2\,X^8 + 7\,X^7\,Z^1 + 9\,X^6\,Z^2 + 9\,X^5\,Z^3 + 8\,X^4\,Z^4 + 3\,X^3\,Z^5 + 2\,X^2\,Z^6 + 4\,X\,Z^7 + 8\,Z^8\,.$$

3. Automorphisms and strata of hyperelliptic genus 3 curves

In the sequel, K is an algebraically closed field of characteristic p where p is a prime or 0.

3.1. **Review on automorphism groups.** Let C be a hyperelliptic curve of genus $g \geq 2$ over K and ι be its hyperelliptic involution. We denote $\mathrm{Aut}(C)$ the automorphism group of C (by convention if C is defined over an arbitrary field k then $\mathrm{Aut}(C)$ is $\mathrm{Aut}(C/\bar{k})$). Since ι is in the center of $\mathrm{Aut}(C)$ we can defined the *reduced automorphism group* $\overline{\mathrm{Aut}}(C) = \mathrm{Aut}(C)/\langle\iota\rangle$. We say that C has *extra-automorphism* when $\#\overline{\mathrm{Aut}}(C) > 1$. The reduced automorphism group acts on $C/\langle\iota\rangle \simeq \mathbb{P}^1$. The list of possible finite groups $\overline{\mathcal{G}}$ acting on $\mathbb{P}^1$ was given in [Web99, § 71-74]. In his PhD thesis, Brandt [Bra88] gave the full list of polynomial orbits under any $\overline{\mathcal{G}}$ which in turn gives the *normal models* of hyperelliptic curves which automorphism group contains a group $\mathcal{G}$ such that $\mathcal{G}/\langle\iota\rangle = \overline{\mathcal{G}}$. The structure of $\mathcal{G}$ itself then depends on the behavior of the exact sequence

$$1 \to \langle\iota\rangle \to \mathcal{G} \to \overline{\mathcal{G}} \to 1$$

which is measured by $H^2(\overline{\mathcal{G}}, \mathbb{Z}/2\mathbb{Z})$. When $p = 0$, the structure of $\mathcal{G}$, depending on its signature, has then been worked out in [BS86]. Finally among the groups $\overline{\mathcal{G}}$, one has to determine the ones which appear as full automorphism group of C. When $p = 0$, this can be done using Fuchsian groups [Sin72], Teichmüller theory [Rie93] or Hurwitz spaces [MSSV02].

For $g = 2$, a complete list of automorphism groups and models can be found in [Car03] for $p \neq 2$ and in [CNP05] for $p = 2$.

For $g = 3$ and $p = 0$, this work has been achieved explicitly in several papers. We refer to [MSSV02] (or [GSS05]) for a historical viewpoint and we gather their results in Table 3 (see also Remark 3.1). Using the signature [MSSV02, Sec.4] (or the shape of the equations), we can deduce the relations between the stratum in the moduli space (*cf.* Figure 1).

For sake of completeness, we show how these results extend to all p. By [Roq70], we know that when $p > 3 + 1 = 4$ and $p \neq 2 \cdot 3 + 1 = 7$, then $C \mapsto C/\mathrm{Aut}(C)$ is tamely ramified and so by [Gro71, XIII.2.12], it can be lifted to characteristic 0. Hence, Table 3 is also valid in these characteristics. When $p = 2$, the possible automorphism groups and models are in [NS04]. When $p = 7$, by [Roq70] the curve $C : y^2 = x^7 - x$ which has a group of order $2^5 \cdot 3 \cdot 7$ is the only exceptional case. Finally for $p = 3$, going through the list of [Bra88, Satz.2.3], it seems that there is no new automorphism group and moreover the cases for which 3 divides $\#\,\mathrm{Aut}(C)$ in Table 3 do not appear anymore.

Remark 3.1. Some remarks on the notation and convention for Table 3 and Figure 1.

- We have the following notation for the groups:
  - $\mathbf{C}_n = \mathbb{Z}/n\mathbb{Z}$;
  - $\mathbf{D}_n$ is the dihedral group with n elements;
  - $\mathbf{U}_6$ is a group with 24 elements defined by $\langle S, T\rangle$ with $S^{12} = T^2 = 1$ and $TST = S^5$;
  - $\mathbf{V}_8$ is a group with 32 elements defined by $V^8 = \langle S, T\rangle$ with $S^4 = T^8 = (ST)^2 = (S^{-1}T)^2 = 1$;
  - $\mathbf{S}_n$ is the symmetric group over n letters.
- An exponent in the signature must be understood as repetition: for instance $(2^6) = (2, 2, 2, 2, 2, 2)$. The dimension δ of the stratum in H_3 is easily computed as $-3 + \#S$ where S is the signature. The column Id. refers to the GAP or MAGMA library of small groups.
- The order chosen for Table 3 (resp. for Figure 1) is by decreasing dimension δ (resp. from top to bottom) then by increasing order of the automorphism group (resp. from left to right).
- The equation of the normal model is valid for the stratum: for some special values of the parameters, the curve can have more automorphisms.

- In both references [MSSV02] and [GSS05], Case 11 of Table 3 is wrongly written as x^8+14x^2+1.
- In [GSS05, Fig.1] and in [BV05, Tab.3], the organization of the strata is wrong.

#	Aut(C)	$\overline{\text{Aut}}(C)$	signature	δ	equation $y^2=f(x)$	Id.
1	$\mathbf{C}_2$	$\{1\}$	(2^8)	5	$x(x-1)(x^5+ax^4+bx^3+cx^2+dx+e)$	$(2,1)$
2	$\mathbf{D}_4$	$\mathbf{C}_2$	(2^6)	3	$\begin{cases} x^8+ax^6+bx^4+cx^2+1 & \text{or} \\ (x^2-1)(x^6+ax^4+bx^2+c) \end{cases}$	$(4,2)$
3	$\mathbf{C}_4$	$\mathbf{C}_2$	$(2^3,4^2)$	2	$x(x^2-1)(x^4+ax^2+b)$	$(4,1)$
4	$\mathbf{C}_2^3$	$\mathbf{D}_4$	(2^5)	2	$(x^4+ax^2+1)(x^4+bx^2+1)$	$(8,5)$
5	$\mathbf{C}_2\times\mathbf{C}_4$	$\mathbf{D}_4$	$(2^2,4^2)$	1	$\begin{cases} (x^4-1)(x^4+ax^2+1) & \text{or} \\ x(x^2-1)(x^4+ax^2+1) \end{cases}$	$(8,2)$
6	$\mathbf{D}_{12}$	$\mathbf{D}_6$	$(2^3,6)$	1	$x(x^6+ax^3+1)$	$(12,4)$
7	$\mathbf{C}_2\times\mathbf{D}_8$	$\mathbf{D}_8$	$(2^3,4)$	1	x^8+ax^4+1	$(16,11)$
8	$\mathbf{C}_{14}$	$\mathbf{C}_7$	$(2,7,14)$	0	x^7-1	$(14,2)$
9	$\mathbf{U}_6$	$\mathbf{D}_{12}$	$(2,4,12)$	0	$x(x^6-1)$	$(24,5)$
10	$\mathbf{V}_8$	$\mathbf{D}_{16}$	$(2,4,8)$	0	x^8-1	$(32,9)$
11	$\mathbf{C}_2\times\mathbf{S}_4$	$\mathbf{S}_4$	$(2,4,6)$	0	x^8+14x^4+1	$(48,48)$

TABLE 3. Automorphism groups for genus 3 hyperelliptic curves in characteristic 0 (see Remark 3.1)



FIGURE 1. Organization of the strata according to inclusion and dimension (see Remark 3.1)

3.2. **Identification of loci and reconstruction with non trivial automorphism groups.** Let C be a hyperelliptic curve of genus 3 defined over a field k of characteristic $p\neq 2,3,5,7$ and let $K=\bar{k}$. Let $(j_2:j_3:\ldots:j_{10})$ be the Shioda invariants of C, our aim is to retrieve from them a hyperelliptic model $y^2=f(x)$.

The conic and quartic method presented in Section 2.3, based on Prop. 2.8 is our main tool for solving this problem, but unfortunately Lemma 3.2 shows that this method can not work for most of the curves with a non trivial automorphism group.

Lemma 3.2. *When the automorphism group of an hyperelliptic curve $C: y^2=f(x)$ contains $\mathbf{D}_4$, i.e. in cases 2,4,5,6,7,9,10,11 of Table 3, then for any choice of 3 quadratic covariants $q_i=q_i(f,(x,z))$ of $f=x^8+ax^6+bx^4+cx^2+1$, we have that $R(q_1,q_2,q_3)=0$.*

Proof. Let d_i be the degree of the covariants q_i. Their weight is then $(8d_i-2)/2 = 4d_i - 1$ which is always odd. Hence, for $g \in \mathrm{Aut}(C)$ acting as $g.(x,z) = (-x,z)$,

$$q_i(g.f, g.(x,z)) = -q_i(f,(x,z)) = q_i(f,(-x,z)).$$

Hence for all i, q_i has only an xz term. In particular the determinant of the q_is in the bases x^2, xz, z^2 is zero. □

We thus have to develop more specific methods for reconstructing models for curves in this case. Given invariants $(j_2 : j_3 : \ldots : j_{10})$, a prerequisite is to determine what is the automorphism group of the corresponding curve so that we can select the right normal model to reconstruct. To this aim, we determine for each automorphism group equations for the corresponding stratum in H_3. There are very few cases where determining some of these equations is straightforward, for instance automorphism groups which contains $\mathbf{C}_4$.

Lemma 3.3. *When the automorphism group of an hyperelliptic curve $C: y^2 = f(x)$ contains $\mathbf{C}_4$, i.e. in cases 3,5,9 and 10, then $J_i(f) = 0$ for all odd index i.*

Proof. The weight of an invariant of degree d is $8d/2 = 4d$. If $\mathrm{Aut}(C)$ contains $\mathbf{C}_4$, the curve C admits a model of the form $y^2 = f(x)$ with $f(x) = x(x^2-1)(x^4+ax^2+1)$. For $g \in \mathrm{Aut}(C)$ acting as $g.(x,z) = (-x,z)$, we get that

$$J_i(g.f) = \det(g)^{4i} J_i(f) = J_i(f).$$

On the other hand, $g.f = -f$, so if the degree of J_i is odd, then $J_i(g.f) = J_i(-f) = -J_i(f)$. Hence we get that $J_i(f) = -J_i(f)$. □

In order to exhibit necessary conditions for all the strata, we have applied Algorithm 2. We choose for I the constant invariant 0, seen as an invariant of increasing positive degree d, and choose for $\mathcal{F}$ (at line 2) a set of random octics over $\mathbb{Q}$ of the form the normal model for the automorphism group that we consider. Increasing one by one d from 0 to 30 yields generators for the ideal of relations which defines the stratum. The shape of these generators is very close to the one of a Gröbner basis for the graded reverse lexicographical (or 'grevlex') order $J_2 < J_3 < \ldots < J_{10}$ with weights $2, 3, \ldots, 10$. For this order, it is thus possible to deduce a reduced Gröbner basis. In the easier cases, note that it is feasible to apply a 'change of order algorithm' and to deduce a reduced Gröbner basis for the lexical order $J_2 < J_3 < \ldots < J_{10}$ too.

Now, in order to exhibit a model from given invariants, we proceed as above, except that we slightly modify Algorithm 2 to add in the basis $\mathcal{B}$ (at line 1) the coefficients a, b, *etc.* of the normal model in Tab. 3 for the considered stratum. We still choose for $\mathcal{F}$ (at line 2) a set of random octics in the shape of the normal model. The lowest degree equations found in this way are then enough to reconstruct a model.

All in all, we give below reconstruction lemmas, one for each automorphism group in Table 3. We precisely state the stratum equations that we have obtained and a model in terms of the j_is for the curve C which may be defined over a non trivial extension of k. In order to make this extension as small as possible, we introduce models which have more non-zero coefficients than the ones in Table 3. We refer the reader to Section 4 for the existence of a model over the field of moduli.

The proofs of these lemmas do not depend on the way we have obtained the equations. They follow essentially all the same principle.

- We check that the normal models of Tab. 3 have Shioda invariants that satisfy the stratum equations, so that we are convinced that these models are a subset of all the models which satisfy the stratum equations.
- Conversely, we check that the reconstructed model has Shioda invariants in the same weighted projective class as the 9-uple $(j_2 : j_3 : \ldots : j_{10})$ provided in input. Since these models are of normal form, we have proved that only normal models satisfy the stratum equations. If we can perform this step, we then have checked that the equations we found describe the stratum.

Most of these proofs need heavy computations, far too complex to be written down here, and so we must skip them. But a program written in the MAGMA computational algebra system that implements the corresponding computations is available on the web page of the authors for independent checks.

Incidentally, we succeed in parameterizing the projective variety defined by the stratum equations of dimension ≤ 2 (it is often a pencil of rational curves). As a first consequence, we give parametrized models over the field of moduli for all these strata, except the cases $\mathbf{C}_2^3$ where we have to deal with algebraic extensions. As a second consequence, we give the exact number of isomorphic classes of curves over a finite field $\mathbb{F}_q$ in these strata.

Algorithm 3: Reconstruct a hyperelliptic polynomial from its Shioda invariants

Input : Shioda invariants $(j_2 : j_3 : \ldots : j_{10})$.
Output: A hyperelliptic polynomial f

1 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (4), then reconstruct f with Lemma 3.4 (case $\mathbf{C}_2 \times \mathbf{S}_4$);
2 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (5), then reconstruct f with Lemma 3.5 (case $\mathbf{V}_8$);
3 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (6), then reconstruct f with Lemma 3.6 (case $\mathbf{U}_6$);
4 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (7), then reconstruct f with Lemma 3.7 (case $\mathbf{C}_{14}$);
5 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (8), then reconstruct f with Lemma 3.8 (case $\mathbf{C}_2 \times \mathbf{D}_8$);
6 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (9), then reconstruct f with Lemma 3.10 (case $\mathbf{D}_{12}$);
7 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (10), then reconstruct f with Lemma 3.12 (case $\mathbf{C}_2 \times \mathbf{C}_4$);
8 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (12), then reconstruct f with Lemma 3.14 (case $\mathbf{C}_2^3$);
9 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (15), then reconstruct f with Lemma 3.16 (case $\mathbf{C}_4$);
10 If $(j_2 : j_3 : \ldots : j_{10})$ satisfies Eq. (18), then reconstruct f with Lemma 3.20 (case $\mathbf{D}_4$);
11 Otherwise, reconstruct f with Lemma 3.24 (case $\mathbf{C}_2$);
12 **return** f;

Algorithm 3 summarizes how one can apply the lemmas below to reconstruct a model in any case. Actually, this algorithm returns more generally a binary form f of degree 8, even in the cases where f does not have simple roots. But, when the order of one of these roots is greater than 4, only one among all the possible orbits, is returned. Typically only $f = x^8$ is returned for $j_2 = j_3 = \ldots = j_{10} = 0$. Moreover the automorphism group of such a form may in general be bigger than the one indicated in the algorithm.

We state now, by increasing dimension, the reconstruction lemmas.

3.2.1. *Strata of dimension 0.*

STRATUM $\mathrm{Aut}(C) = \mathbf{C}_2 \times \mathbf{S}_4$.

Lemma 3.4. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C is $\mathbf{C}_2 \times \mathbf{S}_4$ if and only if*

$$\begin{aligned} 0 &= 30\, {j_3}^2 - {j_2}^3, \\ 0 &= j_4 = j_5 = \cdots = j_{10}\,. \end{aligned} \tag{4}$$

Furthermore, a curve C with automorphism group $\mathbf{C}_2 \times \mathbf{S}_4$ is K-isomorphic to the curve $y^2 = x^8 + 14\,x^4 + 1$.

STRATUM $\mathrm{Aut}(C) = \mathbf{V}_8$.

Lemma 3.5. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C is $\mathbf{V}_8$ if and only*

$$\begin{aligned} 0 &= 6\, j_4 - {j_2}^2\,, \\ 0 &= 36\, j_6 + {j_2}^3\,, \\ 0 &= 420\, j_8 + {j_2}^4\,, \\ 0 &= 2520\, j_{10} - {j_2}^5\,, \\ 0 &= j_3 = j_5 = j_7 = j_9\,. \end{aligned} \tag{5}$$

Furthermore, a curve C with automorphism group $\mathbf{V}_8$ is K-isomorphic to the curve $y^2 = x^8 - 1$.

STRATUM $\mathrm{Aut}(C) = \mathbf{U}_6$.

Lemma 3.6. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C is $\mathbf{U}_6$ if and only if*

$$\begin{aligned} 0 &= 96\, j_4 - {j_2}^2\,, \\ 0 &= 2304\, j_6 + {j_2}^3\,, \\ 0 &= 17920\, j_8 - {j_2}^4\,, \\ 0 &= 430080\, j_{10} + {j_2}^5\,, \\ 0 &= j_3 = j_5 = j_7 = j_9\,. \end{aligned} \tag{6}$$

Furthermore, a curve C with automorphism group $\mathbf{U}_6$ is K-isomorphic to the curve $y^2 = x^7 - x$.

STRATUM $\mathrm{Aut}(C) = \mathbf{C}_{14}$.

Lemma 3.7. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C is $\mathbf{C}_{14}$ if and only if*

$$j_2 \;=\; j_3 \;=\; j_4 \;=\; j_5 \;=\; j_6 \;=\; j_8 \;=\; j_9 \;=\; j_{10} \;=\; 0 \qquad (j_7 \neq 0)\,. \tag{7}$$

Furthermore, a curve C with automorphism group $\mathbf{C}_{14}$ is K-isomorphic to the curve $y^2 = x^7 - 1$.

3.2.2. *Strata of dimension 1.*

STRATUM $\mathrm{Aut}(C) = \mathbf{C}_2 \times \mathbf{D}_8$.

Lemma 3.8. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C contains $\mathbf{C}_2 \times \mathbf{D}_8$ if and only if*

$$\begin{aligned} 0 &= 54\, j_4^3 - 81\, j_4^2\, j_2^2 - 1080\, j_4\, j_3^2\, j_2 + 36\, j_4\, j_2^4 - 3600\, j_3^4 + 240\, j_3^2\, j_2^3 - 4\, j_2^6\,, \\ 0 &= 60\, j_5\, j_3 + 18\, j_4^2 - 15\, j_4\, j_2^2 - 60\, j_3^2\, j_2 + 2\, j_2^4\,, \\ 0 &= 3\, j_5\, j_4 - 2\, j_5\, j_2^2 + 15\, j_4\, j_3\, j_2 + 60\, j_3^3 - 2\, j_3\, j_2^3\,, \\ 0 &= 5\, j_5^2 - 6\, j_4^2\, j_2 - 30\, j_4\, j_3^2 + j_4\, j_2^3\,, \\ 0 &= 18\, j_6 - 9\, j_4\, j_2 - 60\, j_3^2 + 2\, j_2^3\,, \\ 0 &= 3\, j_7 - j_5\, j_2 + 3\, j_4\, j_3\,, \\ 0 &= 1260\, j_8 - 18\, j_4^2 - 63\, j_4\, j_2^2 - 420\, j_3^2\, j_2 + 14\, j_2^4\,, \\ 0 &= 18\, j_9 - 2\, j_5\, j_2^2 + 15\, j_4\, j_3\, j_2 + 60\, j_3^3 - 2\, j_3\, j_2^3\,, \\ 0 &= 630\, j_{10} - 15\, j_4^2\, j_2 - 30\, j_4\, j_3^2 + j_4\, j_2^3\,. \end{aligned} \tag{8}$$

Furthermore, a curve C with automorphism group $\mathbf{C}_2 \times \mathbf{D}_8$ is K-isomorphic to the curve $y^2 = x^8 + a_4\, x^4 + a_0$ where $a_0 = -a_4^2/140 + j_2/2$ and

$$a_4 = \begin{cases} 35(j_5\, j_2 + 6\, j_4\, j_3)/(-60\, {j_3}^2 + 2{j_2}^3) & \text{if } -30\, {j_3}^2 + {j_2}^3 \neq 0\,, \\ 35 j_5 \,/\, 3 j_4 & \text{otherwise}\,. \end{cases}$$

Remark 3.9. When $-30\, {j_3}^2 + {j_2}^3$ and j_4 are both equal to 0, Eq. (8) can be reduced to Eq. (4). This means that C has a larger automorphism group, specifically $\mathbf{C}_2 \times \mathbf{S}_4$. So Lemma 3.4 can be used, instead, for reconstructing a model.

Actually the quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (8) is obtained by adjoining j_4 to $k[j_2, j_3]$ and j_4 satisfies an irreducible monic equation of degree 3. The invariants j_5, j_6, …, j_{10} are then rational in j_2, j_3 and j_4.

From a more geometric viewpoint, the projective variety defined by Eq. (8) has two singularities, the $\mathbf{C}_2 \times \mathbf{S}_4$ and $\mathbf{V}_8$ points. Moreover, it is birationally equivalent to the conic ${X_2}^2 - 400\, X_1\, X_3 + (4000/3)\, {X_3}^2$ of discriminant $-2^8 \cdot 5^4$. We can classically parameterized it by the slope of lines which all intersect at the $\mathbf{V}_8$ point. This yields

$$t \mapsto \left(\frac{30\,t+1}{30} : \frac{45\,t+1}{900} : \frac{t^2}{6} : \frac{t^2}{60} : \frac{-t^3}{36} : \frac{-t^3}{360} : \frac{-t^3\,(180\,t+7)}{2^4 \cdot 3^3 \cdot 5^2 \cdot 7} : \frac{t^4}{2^4 \cdot 3^3 \cdot 5} : \frac{t^4\,(180\,t+7)}{2^5 \cdot 3^4 \cdot 5^2 \cdot 7} \right).$$

The corresponding curves are of the form

$$y^2 = x^8 + 7/6\, x^4 + 1/2\, t + 1/144\,.$$

The values $t = -1/72$ and $t = 2/3$ give polynomials with multiple roots, the curve with automorphism group $\mathbf{C}_2 \times \mathbf{S}_4$ is obtained with $t = 0$, otherwise the automorphism group is always $\mathbf{C}_2 \times \mathbf{D}_8$ since the point $\mathbf{V}_8$ is not reachable with this conic parametrization we chose. Incidentally, this shows that over a finite field $\mathbb{F}_q$ of characteristic at least 11 there are $q - 3$ isomorphic classes of curves with automorphism group $\mathbf{C}_2 \times \mathbf{S}_4$.

STRATUM $\mathrm{Aut}(C) = \mathbf{D}_{12}$.

Lemma 3.10. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C contains $\mathbf{D}_{12}$ if and only if*

$$\begin{aligned}
0 &= 55296\, j_4^3 - 5184\, j_4^2\, j_2^2 - 4320\, j_4\, j_3^2\, j_2 + 144\, j_4\, j_2^4 - 900\, j_3^4 + 60\, j_3^2\, j_2^3 - j_2^6\,,\\
0 &= 720\, j_5\, j_3 - 2304\, j_4^2 + 120\, j_4\, j_2^2 + 30\, j_3^2\, j_2 - j_2^4\,,\\
0 &= 576\, j_5\, j_4 - 24\, j_5\, j_2^2 - 120\, j_4\, j_3\, j_2 - 30\, j_3^3 + j_3\, j_2^3\,,\\
0 &= 180\, j_5^2 - 96\, j_4^2\, j_2 - 30\, j_4\, j_3^2 + j_4\, j_2^3\,,\\
0 &= 576\, j_6 - 72\, j_4\, j_2 - 30\, j_3^2 + j_2^3\,,\\
0 &= 12\, j_7 - j_5\, j_2 - 2\, j_4\, j_3\,,\\
0 &= 60480\, j_8 - 14976\, j_4^2 + 504\, j_4\, j_2^2 + 210\, j_3^2\, j_2 - 7\, j_2^4\,,\\
0 &= 3456\, j_9 - 24\, j_5\, j_2^2 - 120\, j_4\, j_3\, j_2 - 30\, j_3^3 + j_3\, j_2^3\,,\\
0 &= 30240\, j_{10} - 600\, j_4^2\, j_2 - 390\, j_4\, j_3^2 + 13\, j_4\, j_2^3\,.
\end{aligned} \tag{9}$$

Furthermore, a curve C with automorphism group $\mathbf{D}_{12}$ is K-isomorphic to the curve $y^2 = x\,(x^6 + a_4\, x^3 + a_1)$ where $a_1 = 2\, {a_4}^2 \,/\, 35 - 4\, j_2$ and

$$a_4 = \begin{cases} 280\,(-j_5\, j_2 + 4\, j_4\, j_3)/(30\, {j_3}^2 - {j_2}^3) & \text{if } -30\, {j_3}^2 + {j_2}^3 \neq 0\,,\\ 35 j_5 \,/\, 3 j_4 & \text{otherwise}\,. \end{cases}$$

Remark 3.11. When $-30\, {j_3}^2 + {j_2}^3$ and j_4 are both equal to 0, Eq. (9) can be reduced to Eq. (4). Again, this means that C has a larger automorphism group, specifically $\mathbf{C}_2 \times \mathbf{S}_4$. So, Lemma 3.4 can be used, instead, for reconstructing a model.

Similarly to $\mathbf{C}_2 \times \mathbf{D}_8$, the quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (9) is obtained by adjoining j_4 to $k[j_2, j_3]$ and j_4 satisfies an irreducible monic equation of degree 3. The invariants j_5, j_6, …, j_{10} are then rational in j_2, j_3 and j_4.

More geometrically, the projective variety defined by Eq. (9) has two singularities, the $\mathbf{C}_2 \times \mathbf{S}_4$ and $\mathbf{U}_6$ points. Moreover, it is birationally equivalent to the conic ${X_2}^2 - (25/16)\, X_1\, X_3 + (125/6144)\, {X_3}^2$ of discriminant $-5^4/2^8$. We can classically parameterized it by the slope of lines which all intersect at the $\mathbf{U}_6$ point, this yields

$$t \mapsto \left(\frac{360\, t + 1}{30} : \frac{540\, t + 1}{900} : \frac{3\, t^2}{2} : \frac{-t^2}{10} : \frac{-3\, t^3}{4} : \frac{t^3}{20} : \frac{t^3\,(4860\, t + 7)}{2^3 \cdot 3 \cdot 5^2 \cdot 7} : \frac{-t^4}{40} : \frac{-t^4\,(4860\, t + 7)}{2^4 \cdot 3 \cdot 5^2 \cdot 7} \right).$$

The corresponding curves are of the form

$$y^2 = x\,(x^6 - 7/9\, x^3 - 48\, t - 8/81)\,.$$

The values $t = -1/(2 \cdot 3^5)$ and $t = -1/(2^6 \cdot 3)$ give polynomials with multiple roots, the curve with automorphism group $\mathbf{C}_2 \times \mathbf{S}_4$ is obtained with $t = 0$, otherwise the automorphism group is always $\mathbf{D}_{12}$ since the point $\mathbf{U}_6$ is not reachable with this conic parametrization we chose. Incidentally, this shows that over a finite field $\mathbb{F}_q$ of characteristic at least 11 there are $q - 3$ isomorphic classes of curves with automorphism group $\mathbf{D}_{12}$.

STRATUM $\mathrm{Aut}(C) = \mathbf{C}_2 \times \mathbf{C}_4$.

Lemma 3.12. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C contains $\mathbf{C}_2 \times \mathbf{C}_4$ if and only if*

$$\begin{aligned} 0 &= 648000\, j_6^2 - 356400\, j_6\, j_4\, j_2 + 4275\, j_6\, j_2^3 - 839808\, j_4^3 + 75249\, j_4^2\, j_2^2 - 1449\, j_4\, j_2^4 + 8\, j_2^6\,, \\ 0 &= 85050\, j_8 + 13860\, j_6\, j_2 + 37908\, j_4^2 - 2961\, j_4\, j_2^2 + 28\, j_2^4\,, \\ 0 &= 2041200\, j_{10} + 560520\, j_6\, j_4 + 2205\, j_6\, j_2^2 + 98091\, j_4^2\, j_2 - 5607\, j_4\, j_2^3 + 56\, j_2^5\,, \\ 0 &= j_3 = j_5 = j_7 = j_9\,. \end{aligned} \tag{10}$$

Furthermore, a curve C with automorphism group $\mathbf{C}_2 \times \mathbf{C}_4$ is K-isomorphic to the curve $y^2 = a^2 x^8 + 2\, a^2 x^6 + 8\, a x^2 - 16$ with

$$a = \begin{cases} 196/3 & \text{if } 6\, j_4 - {j_2}^2 = 0\,, \\ -84 & \text{if } 147\, j_4 - 2\, {j_2}^2 = 0\,, \\ 98\, \dfrac{36288\, {j_4}^2 - 3906\, j_4\, {j_2}^2 + 14400\, j_6\, j_2 + 43\, {j_2}^4}{9(96\, j_4 - {j_2}^2)(147\, j_4 - 2\, {j_2}^2)} & \text{otherwise}\,. \end{cases} \tag{11}$$

Remark 3.13. There is no difficulty to check that curves that annihilate denominators in Eq. (11) have a larger automorphism group than $\mathbf{C}_2 \times \mathbf{C}_4$.

- When $6\, j_4 - {j_2}^2$ and $36\, j_6 + {j_2}^3$ are both equal to 0, Eq. (10) can be reduced to Eq. (5) and Lemma 3.5 can be used, instead, for reconstructing a model. Otherwise, *i.e.* when $6\, j_4 - {j_2}^2 = 0$ but $36\, j_6 + {j_2}^3 \neq 0$, then now $576\, j_6 - 65\, {j_2}^3 = 0$ but this lemma can still be applied.
- When $96\, j_4 - {j_2}^2 = 0$, Eq. (10) can be reduced to Eq. (6) and Lemma 3.6 can now be used for reconstructing a model.
- When $147\, j_4 - 2\, {j_2}^2 = 0$ and $3087\, j_6 - 2\, {j_2}^3 = 0$, then we can choose for $f(x)$ the polynomial $x(x-1)(x+1)(x^2+1)^2$. Its discriminant is obviously equal to zero and no hyperelliptic curve exists with such invariants. Otherwise, *i.e.* when $147\, j_4 - 2\, {j_2}^2 = 0$ but $3087\, j_6 - 2\, {j_2}^3 \neq 0$, then now $197568\, j_6 - 47\, {j_2}^3 = 0$ but this lemma can still be applied.

Here, the quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (10) is obtained by adjoining j_6 to $k[j_2, j_4]$ and j_6 satisfies an irreducible monic equation of degree 2. The invariants j_8 and j_{10} are then rational in j_2, j_4 and j_6 (the invariants j_3, j_5, j_7 and j_9 are trivial).

Geometrically, the projective variety defined by Eq. (10) has two singularities, the $\mathbf{V}_8$ and $\mathbf{U}_6$ points. Moreover, it is birationally equivalent to the conic ${X_1}^2 - 11/20\; X_1\, X_3 - 162/125\; X_2\, X_3 + 713/8000\, {X_3}^2$ of discriminant $-2^2 \cdot 3^8/5^6$. We can classically parameterized it by the slope of lines which all intersect at the $\mathbf{V}_8$ point, this yields

$$\begin{aligned} t \mapsto \Bigg(t \;:\; 0 \;:\; \frac{t^2}{6} - \frac{3\,t}{2} + \frac{18}{5} \;:\; 0 \;:\; -\frac{(5\,t-27)\left(25\,t^2 - 405\,t + 1296\right)}{4500} \;:\; 0 \;:\; \\ -\frac{t^4}{420} + \frac{51\,t^3}{700} - \frac{684\,t^2}{875} + \frac{15516\,t}{4375} - \frac{25272}{4375} \;:\; 0 \;:\; \frac{t^5}{2520} - \frac{11\,t^4}{560} + \frac{3\,t^3}{10} - \frac{813\,t^2}{400} + \frac{28026\,t}{4375} - \frac{168156}{21875} \Bigg)\,. \end{aligned}$$

The corresponding curves are of the form

$$y^2 = (x^4 + 2\,x^2 + 5/147\,t - 1/7)(x^4 - 5/147\,t + 1/7)\,.$$

The values $t = 21/5$ and $t = 168/5$ give polynomials with multiple roots, the curve with automorphism group $\mathbf{U}_6$ is obtained with $t = 24/5$, otherwise the automorphism group is always $\mathbf{C}_2 \times \mathbf{C}_4$ since the point $\mathbf{V}_8$ is not reachable with this conic parametrization. Incidentally, this shows again that over a finite field $\mathbb{F}_q$ of characteristic at least 11 there are $q - 3$ isomorphic classes of curves with automorphism group $\mathbf{C}_2 \times \mathbf{C}_4$.

3.2.3. *Strata of dimension 2.*

STRATUM $\mathrm{Aut}(C) = \mathbf{C}_2^3$.

Lemma 3.14. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. Then the automorphism group of C contains $\mathbf{C}_2^3$ if and only if*

$$
\begin{aligned}
0 &= 97200\,j_5^4 - 656100\,j_5^3\,j_3\,j_2 - 437400\,j_5^2\,j_4^2\,j_2 + 2916000\,j_5^2\,j_4\,j_3^2 + 212625\,j_5^2\,j_4\,j_2^3 + 753300\,j_5^2\,j_3^2\,j_2^2 - 25110\,j_5^2\,j_2^5 \\
&\quad + 2494800\,j_5\,j_4^3\,j_3 - 1336500\,j_5\,j_4^2\,j_3\,j_2^2 - 5427000\,j_5\,j_4\,j_3^3\,j_2 + 180900\,j_5\,j_4\,j_3\,j_2^4 - 3024000\,j_5\,j_3^5 + 201600\,j_5\,j_3^3\,j_2^3 \\
&\quad - 3360\,j_5\,j_3\,j_2^6 + 381024\,j_4^5 - 259200\,j_4^4\,j_2^2 - 186300\,j_4^3\,j_3^2\,j_2 + 66960\,j_4^3\,j_2^4 + 2592000\,j_4^2\,j_3^4 + 151200\,j_4^2\,j_3^2\,j_2^3 \\
&\quad - 7920\,j_4^2\,j_2^6 + 351000\,j_4\,j_3^4\,j_2^2 - 23400\,j_4\,j_3^2\,j_2^5 + 390\,j_4\,j_2^8 + 108000\,j_3^6\,j_2 - 10800\,j_3^4\,j_2^4 + 360\,j_3^2\,j_2^7 - 4\,j_2^{10}\,, \\
0 &= 7620480\,j_6\,j_4^3 - 4626720\,j_6\,j_4^2\,j_2^2 - 6609600\,j_6\,j_4\,j_3^2\,j_2 + 929070\,j_6\,j_4\,j_2^4 + 10368000\,j_6\,j_3^4 + 1502550\,j_6\,j_3^2\,j_2^3 - 61605\,j_6\,j_2^6 \\
&\quad + 874800\,j_5^3\,j_3 - 2449440\,j_5^2\,j_4^2 + 947700\,j_5^2\,j_4\,j_2^2 - 3207600\,j_5^2\,j_3^2\,j_2 - 89910\,j_5^2\,j_2^4 + 3440880\,j_5\,j_4^2\,j_3\,j_2 + 16912800\,j_5\,j_4\,j_3^3 \\
&\quad - 982935\,j_5\,j_4\,j_3\,j_2^3 - 1555200\,j_5\,j_3^3\,j_2^2 + 51840\,j_5\,j_3\,j_2^5 + 843696\,j_4^4\,j_2 - 4581360\,j_4^3\,j_3^2 - 559278\,j_4^3\,j_2^3 - 168480\,j_4^2\,j_3^2\,j_2^2 \\
&\quad + 125091\,j_4^2\,j_2^5 - 64800\,j_4\,j_3^4\,j_2 + 305370\,j_4\,j_3^2\,j_2^4 - 10107\,j_4\,j_2^7 + 432000\,j_3^6 + 104400\,j_3^4\,j_2^3 - 8400\,j_3^2\,j_2^6 + 148\,j_2^9\,, \\
0 &= 16200\,j_6\,j_5\,j_3 + 45360\,j_6\,j_4^2 - 17550\,j_6\,j_4\,j_2^2 - 49950\,j_6\,j_3^2\,j_2 + 1665\,j_6\,j_2^4 - 14580\,j_5^2\,j_4 + 2430\,j_5^2\,j_2^2 + 22275\,j_5\,j_4\,j_3\,j_2 \\
&\quad + 43200\,j_5\,j_3^3 - 1440\,j_5\,j_3\,j_2^3 + 5022\,j_4^3\,j_2 - 34560\,j_4^2\,j_3^2 - 2223\,j_4^2\,j_2^3 - 7650\,j_4\,j_3^2\,j_2^2 + 255\,j_4\,j_2^5 - 3600\,j_3^4\,j_2 + 240\,j_3^2\,j_2^4 - 4\,j_2^7\,, \\
0 &= 22680\,j_6\,j_5\,j_4 - 4995\,j_6\,j_5\,j_2^2 - 14850\,j_6\,j_4\,j_3\,j_2 - 86400\,j_6\,j_3^3 + 2880\,j_6\,j_3\,j_2^3 - 7290\,j_5^3 \\
&\quad + 26730\,j_5^2\,j_3\,j_2 + 2511\,j_5\,j_4^2\,j_2 - 80460\,j_5\,j_4\,j_3^2 - 693\,j_5\,j_4\,j_2^3 - 360\,j_5\,j_3^2\,j_2^2 + 12\,j_5\,j_2^5 \\
&\quad - 10206\,j_4^3\,j_3 + 1350\,j_4^2\,j_3\,j_2^2 - 4500\,j_4\,j_3^3\,j_2 + 150\,j_4\,j_3\,j_2^4 - 3600\,j_3^5 + 240\,j_3^3\,j_2^3 - 4\,j_3\,j_2^6\,, \\
0 &= 194400\,j_6\,j_5^2 + 2034720\,j_6\,j_4^2\,j_2 + 2073600\,j_6\,j_4\,j_3^2 - 777870\,j_6\,j_4\,j_2^3 - 2193750\,j_6\,j_3^2\,j_2^2 + 73125\,j_6\,j_2^5 - 913680\,j_5^2\,j_4\,j_2 \\
&\quad + 518400\,j_5^2\,j_3^2 + 179550\,j_5^2\,j_2^3 + 1516320\,j_5\,j_4^2\,j_3 + 419175\,j_5\,j_4\,j_3\,j_2^2 + 1555200\,j_5\,j_3^3\,j_2 - 51840\,j_5\,j_3\,j_2^4 + 244944\,j_4^4 \\
&\quad + 112590\,j_4^3\,j_2^2 - 1170720\,j_4^2\,j_3^2\,j_2 - 80451\,j_4^2\,j_2^4 + 86400\,j_4\,j_3^4 - 306810\,j_4\,j_3^2\,j_2^3 + 10131\,j_4\,j_2^6 - 147600\,j_3^4\,j_2^2 + 9840\,j_3^2\,j_2^5 - 164\,j_2^8\,,
\end{aligned}
$$

$$
\begin{aligned}
0 &= 32400\,j_6^2 + 28350\,j_6\,j_4\,j_2 + 87750\,j_6\,j_3^2 - 2925\,j_6\,j_2^3 - 21870\,j_5^2\,j_2 + 76545\,j_5\,j_4\,j_3 \\
&\quad + 13122\,j_4^3 - 405\,j_4^2\,j_2^2 + 5130\,j_4\,j_3^2\,j_2 - 171\,j_4\,j_2^4 + 3600\,j_3^4 - 240\,j_3^2\,j_2^3 + 4\,j_2^6\,,
\end{aligned}
\tag{12}
$$

$$
\begin{aligned}
0 &= 12150\,j_7\,j_3 + 7560\,j_6\,j_4 - 1665\,j_6\,j_2^2 - 2430\,j_5^2 + 1620\,j_5\,j_3\,j_2 + 837\,j_4^2\,j_2 + 1530\,j_4\,j_3^2 - 231\,j_4\,j_2^3 - 120\,j_3^2\,j_2^2 + 4\,j_2^5\,, \\
0 &= 4860\,j_7\,j_4 - 810\,j_7\,j_2^2 - 1080\,j_6\,j_5 + 3330\,j_6\,j_3\,j_2 - 837\,j_5\,j_4\,j_2 - 2880\,j_5\,j_3^2 - 12\,j_5\,j_2^3 + 2916\,j_4^2\,j_3 + 360\,j_4\,j_3\,j_2^2 + 240\,j_3^3\,j_2 - 8\,j_3\,j_2^4\,, \\
0 &= 18225\,j_7\,j_5 + 7425\,j_6\,j_4\,j_2 + 43200\,j_6\,j_3^2 - 1440\,j_6\,j_2^3 - 10935\,j_5^2\,j_2 + 42525\,j_5\,j_4\,j_3 \\
&\quad + 5103\,j_4^3 - 675\,j_4^2\,j_2^2 + 2250\,j_4\,j_3^2\,j_2 - 75\,j_4\,j_2^4 + 1800\,j_3^4 - 120\,j_3^2\,j_2^3 + 2\,j_2^6\,, \\
0 &= 145800\,j_7\,j_6 + 4050\,j_7\,j_2^3 - 64800\,j_6\,j_5\,j_2 + 83700\,j_6\,j_4\,j_3 - 16650\,j_6\,j_3\,j_2^2 + 77760\,j_5^2\,j_3 + 13122\,j_5\,j_4^2 \\
&\quad - 5751\,j_5\,j_4\,j_2^2 + 8640\,j_5\,j_3^2\,j_2 + 252\,j_5\,j_2^4 - 10260\,j_4^2\,j_3\,j_2 + 2880\,j_4\,j_3^3 - 1896\,j_4\,j_3\,j_2^3 - 1200\,j_3^3\,j_2^2 + 40\,j_3\,j_2^5\,, \\
0 &= 546750\,j_7^2 - 963900\,j_6\,j_4^2 + 453600\,j_6\,j_4\,j_2^2 + 1404000\,j_6\,j_3^2\,j_2 - 46800\,j_6\,j_2^4 + 320760\,j_5^2\,j_4 \\
&\quad - 196830\,j_5^2\,j_2^2 + 204120\,j_5\,j_4\,j_3\,j_2 - 680400\,j_5\,j_3^3 + 22680\,j_5\,j_3\,j_2^3 - 12312\,j_4^3\,j_2 \\
&\quad + 537570\,j_4^2\,j_3^2 + 32301\,j_4^2\,j_2^3 + 157680\,j_4\,j_3^2\,j_2^2 - 5256\,j_4\,j_2^5 + 82800\,j_3^4\,j_2 - 5520\,j_3^2\,j_2^4 + 92\,j_2^7\,, \\
0 &= 85050\,j_8 - 11655\,j_6\,j_2 + 11340\,j_5\,j_3 + 2187\,j_4^2 - 1260\,j_4\,j_2^2 - 840\,j_3^2\,j_2 + 28\,j_2^4\,, \\
0 &= 9720\,j_9 + 810\,j_7\,j_2 - 3330\,j_6\,j_3 - 2187\,j_5\,j_4 + 108\,j_5\,j_2^2 - 360\,j_4\,j_3\,j_2 - 240\,j_3^3 + 8\,j_3\,j_2^3\,, \\
0 &= 170100\,j_{10} + 37890\,j_6\,j_4 - 22680\,j_5^2 + 4221\,j_4^2\,j_2 + 1680\,j_4\,j_3^2 - 56\,j_4\,j_2^3\,.
\end{aligned}
$$

Furthermore, a curve C with automorphism group $\mathbf{C}_2^3$ is K-isomorphic to the curve $y^2 = a_8\,x^8 + a_6\,x^6 + a_4\,x^4 + \lambda a_6\,x^2 + \lambda^2 a_8$ where a_4 is any root of the degree 3 equation

$$
\begin{aligned}
0 &= 192\,(-60\,{j_3}^2 + 2\,{j_2}^3 + 18\,j_6 - 9\,j_4\,j_2)\,x^3 - 90720\,(3\,j_4\,j_3 + 3\,j_7 - j_5\,j_2)\,x^2 \\
&\quad + 294\,(765\,j_4\,{j_2}^2 + 1440\,j_6\,j_2 - 5940\,j_5\,j_3 - 1782\,{j_4}^2 + 1140\,{j_3}^2 j_2 - 38\,{j_2}^4)\,x \\
&\quad + 6637050\,j_6\,j_3 + 1250235\,j_5\,j_4 - 833490\,j_5\,{j_2}^2 + 2932650\,j_4\,j_3\,j_2 + 2881200\,{j_3}^3 - 96040\,j_3\,{j_2}^3
\end{aligned}
\tag{13}
$$

and $a_6 = -28\,\nu^2 - {a_4}^2/5 + 14\,j_2$, $a_8 = \nu\,a_6$ and $\lambda = 1/a_6$ with

$$
\nu = \frac{18\,j_6\,a_4 - 9\,a_4\,j_4\,j_2 - 60\,a_4\,{j_3}^2 + 2\,a_4\,{j_2}^3 - 810\,j_7 + 270\,j_5\,j_2 - 810\,j_4\,j_3}{10\,(-18\,j_6 + 9\,j_4\,j_2 + 60\,{j_3}^2 - 2\,{j_2}^3)}\,.
$$

Eq. (13) may not have a root in the base field k. In this case, Lemma 3.14 yields a model over an extension k' of degree 3 (see Section 3 for the rationality issue).

Remark 3.15. When $-18\,j_6 + 9\,j_4\,j_2 + 60\,{j_3}^2 - 2\,{j_2}^3 = 0$, Eq. (12) can be reduced to Eq. (8) and Lemma 3.8 can be used, instead, for reconstructing a model.

The quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (12) is obtained by adjoining j_5 to $k[j_2, j_3, j_4]$ and j_5 satisfies an irreducible monic equation of degree 4. The invariants j_6, j_7 , $\dots$ j_{10} are then rational in j_2, j_3, j_4 and j_5.

More geometrically, singularities of the projective variety defined by Eq. (12) are on the curve $\mathbf{C}_2 \times \mathbf{D}_8$. Moreover, for non-zero J_2, J_3, we can set $J_3 = J_2 = t$ for some parameter t and the intersection of the stratum with this hyperplane is a pencil of rational curves that we can parametrize by an other parameter u. So, we find for $t \neq 0$ the parametrization $(t, u) \mapsto (j_2(t, u) : j_3(t, u) : \cdots : j_{10}(t, u))$ where

$$
\begin{aligned}
&j_2(t,u) = t\,,\ \ j_3(t,u) = t\,, j_4(t,u) = -9800\,u^4 + 85\,tu^2 - \frac{35}{3}\,tu + \frac{2}{147}\,t^2\,,\\
&j_5(t,u) = -137200\,u^5 + 700\,tu^3 + \frac{350}{3}\,tu^2 - \frac{25}{28}\,t^2u + \frac{1}{28}\,t^2,\\
&j_6(t,u) = -617400\,u^6 + 2030\,tu^4 + 1470\,tu^3 - \frac{689}{56}\,t^2u^2 + \frac{11}{12}\,t^2u - \frac{1}{24}\,t^2 + \frac{2}{3087}\,t^3,\\
&j_7(t,u) = -1920800\,u^7 - \frac{72520}{3}\,tu^5 + \frac{43120}{3}\,tu^4 + \frac{377}{6}\,t^2u^3 - \frac{226}{9}\,t^2u^2 + \frac{1}{588}\,t^2(17\,t+686)u - \frac{1}{588}\,t^3,\\
&j_8(t,u) = -2469600\,u^8 - \frac{125300}{3}\,tu^6 + \frac{37240}{3}\,tu^5 - \frac{643}{14}\,t^2u^4 + \frac{1432}{9}\,t^2u^3\\
&\qquad - \frac{1}{246960}\,t^2(4705960 + 120087\,t)u^2 + \frac{1411}{17640}\,t^3u - \frac{11}{252105}\,t^4 - \frac{1}{1680}\,t^3,\\
&j_9(t,u) = 302526000\,u^9 - \frac{12022150}{3}\,tu^7 - \frac{325850}{3}\,tu^6 + \frac{665105}{36}\,t^2u^5 - \frac{9905}{18}\,t^2u^4\\
&\qquad - \frac{1}{2016}\,t^2(-397880 + 56341\,t)u^3 + \frac{18829}{6048}\,t^3u^2 + \frac{1}{98784}\,t^3(472\,t - 30527)u + \frac{1}{96}\,t^3 - \frac{1}{4116}\,t^4,\\
&j_{10}(t,u) = 1162084000\,u^{10} - 11872700\,tu^8 - \frac{7991900}{3}\,tu^7 + \frac{1366535}{18}\,t^2u^6 - \frac{51835}{9}\,t^2u^5 - \frac{1}{2352}\,t^2(-13267240 + 268943\,t)u^4\\
&\qquad - \frac{77657}{3024}\,t^3u^3 + \frac{1}{1037232}\,t^3(67571 + 105696\,t)u^2 - \frac{1}{148176}\,t^3(824\,t - 1029)u - \frac{11}{5294205}\,t^5 + \frac{1}{6174}\,t^4\,.
\end{aligned}
\tag{14}
$$

The lines $280\,u^2 - t = 0$ and $14280\,u^3 - 33\,tu - 7\,t = 0$ are the loci of forms with multiple roots (discriminant equal to 0). The rational curve $-564715200\,u^6 + 6338640\,u^4t + 1344560\,u^3t - 50127\,u^2t^2 + 6174\,ut^2 + 96\,t^3 - 3087\,t^2 = 0$ is the locus of forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{D}_8$.

In the case $J_2 = 0$ and non-zero J_3, J_4, setting $J_4 = J_3$ yields a rational curve that we parametrize as $t \mapsto (j_2(t) : j_3(t) : \ldots : j_{10}(t))$, with $t \neq -105/1260$, where

$$
\begin{aligned}
&j_2(t) = 0\,,\ \ j_3(t) = 1260\,t + 105\,,\ \ j_4(t) = 7350\,t\,,\ \ j_5(t) = 36750\,t + 7350\,,\ \ j_6(t) = -66150\,t^2 - 242550\,t - 29400\,,\\
&j_7(t) = -926100\,t^2 + 977550\,t + 102900\,,\ \ j_8(t) = -7563150\,t^2 - 1749300\,t - 102900\,,\\
&j_9(t) = 20837250\,t^3 - 33957000\,t^2 - 8232000\,t - 1029000\,,\ \ j_{10}(t) = -6945750\,t^3 + 557975250\,t^2 + 119364000\,t + 7203000\,.
\end{aligned}
$$

The point $t = -2/7$ yields a form with multiple roots, the roots of $81\,t^2 + 17\,t + 1 = 0$ yield forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{D}_8$.

Similarly, in the case $J_3 = 0$ and non-zero J_2, J_5, setting $J_5 = J_2^2$ yields the rational curve

$$
\begin{aligned}
&j_2(t) = \frac{1010}{3}\,(52\,t+59)(t+1)\,,\ \ j_3(t) = 0\,,\ \ j_4(t) = \frac{50}{27}\,(52\,t+59)^2(11272\,t^2 + 23147\,t + 11767)\,,\\
&j_5(t) = \frac{25}{9}\,(52\,t+59)^3(17\,t+29)^2\,,\ \ j_6(t) = -\frac{25}{486}\,(52\,t+59)^3(27670948\,t^3 + 88650255\,t^2 + 94796346\,t + 33852031)\,,\\
&j_7(t) = \frac{50}{81}\,(52\,t+59)^4(17\,t+29)(38117\,t^2 + 81691\,t + 43682)\,,\\
&j_8(t) = -\frac{25}{3402}\,(52\,t+59)^4(6266813468\,t^4 + 26795826605\,t^3 + 42971724291\,t^2 + 30632960351\,t + 8190389165)\,,\\
&j_9(t) = -\frac{125}{2916}\,(52\,t+59)^5(17\,t+29)(11612756\,t^3 + 34093287\,t^2 + 33246138\,t + 10800599)\,,\\
&j_{10}(t) = \frac{125}{30618}\,(52\,t+59)^5(794827443664\,t^5 + 4302413461544\,t^4 + 9310923963145\,t^3 + 10069939679311\,t^2 + 5442845374787\,t + 1176260576869)\,.
\end{aligned}
$$

The point $\left(\frac{1010}{39} : 0 : \frac{563600}{4563} : \frac{14450}{1521} : -\frac{345886850}{533871} : \frac{64798900}{177957} : -\frac{78335168350}{48582261} : -\frac{12338553250}{20820969} : \frac{49676715229000}{5684124537}\right)$, when defined, is the only point which is not reached by the parametrization. Moreover, the points $t = -4259/2831$, $t = -59/52$ and $t = 1/83$ (when defined) yield forms with multiple roots. The points $t = -59/52$, $t = -29/17$ (when defined) and the roots of $9844\,t^2 + 19487\,t + 9679 = 0$ yield forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{D}_8$.

Modulo 101, the latter degenerates to only one projective point. In this case, we thus consider instead the rational curve

$$j_2(t) = 32\,(t+44)(t+12)\,,\ \ j_3(t) = 0\,,\ \ j_4(t) = 57\,(t^2+84\,t+59)(t+12)^2\,,\ \ j_5(t) = 14\,(t+12)^3(t+84)^2\,,$$
$$j_6(t) = 22\,\frac{(t+62)(t+12)^3(t+22)(t^2+73\,t+23)(t+54)}{2\,t^2+45\,t+46}\,,\ \ j_7(t) = \frac{97}{51}\,(t+12)^4(t^2+7\,t+86)(t+84)\,,$$
$$j_8(t) = 81\,\frac{(t+12)^4(t^2+51\,t+54)(t^2+73\,t+23)(t^2+64\,t+62)}{2\,t^2+45\,t+46}\,,\ \ j_9(t) = \frac{2}{51}\,(t+12)^5(t^2+3\,t+19)(t+73)(t+84)\,,$$
$$j_{10}(t) = 3\,\frac{(t^2+3\,t+75)(t^2+26\,t+79)(t+12)^5(t^2+73\,t+23)(t+68)}{2\,t^2+45\,t+46}\,.$$

The point $(32:0:57:14:11:93:91:4:52)$ is the only point which is not reached by the parametrization. Moreover, the points $t=20$, $t=56$ and $t=89$ yield forms with multiple roots. The points $t=17$, $t=89$ and the roots of $t^2+14\,t+87=0$ yield forms with automorphism groups which contain $\mathbf{C}_2\times\mathbf{D}_8$.

We can now easily check that the rational projective points defined on this variety are none other than the ones given by these parametrizations. Incidentally, this shows that over a finite field $\mathbb{F}_q$ of characteristic at least 11 there are $q^2-2\,q+2$ isomorphic classes of curves with automorphism group $\mathbf{C}_2^3$.

STRATUM $\mathrm{Aut}(C)=\mathbf{C}_4$.

Unlike the previous strata, curves with automorphism group $\mathbf{C}_4$ can be reconstructed with the conic and quartic method of Section 2.3.

Lemma 3.16. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2:j_3:\ldots:j_{10})$ be its Shioda invariants. Then the automorphism group of C contains $\mathbf{C}_4$ if and only if*

$$\begin{aligned}
0 &= 1093705578000\,j_8^3 + 291654820800\,j_8^2\,j_6\,j_2 - 1104121821600\,j_8^2\,j_4^2 + 6076142100\,j_8^2\,j_4\,j_2^2 - 1469076134400\,j_8\,j_6^2\,j_4\\
&\quad + 36726903360\,j_8\,j_6^2\,j_2^2 + 231472080000\,j_8\,j_6\,j_4^2\,j_2 - 6631246440\,j_8\,j_6\,j_4\,j_2^3 + 30005640\,j_8\,j_6\,j_2^5\\
&\quad + 267846264000\,j_8\,j_4^4 - 23803045560\,j_8\,j_4^3\,j_2^2 + 540101520\,j_8\,j_4^2\,j_2^4 - 3333960\,j_8\,j_4\,j_2^6 - 256048128000\,j_6^4\\
&\quad + 42247941120\,j_6^3\,j_4\,j_2 - 405409536\,j_6^3\,j_2^3 + 8888527872\,j_6^2\,j_4^3 - 5893679232\,j_6^2\,j_4^2\,j_2^2 + 274718304\,j_6^2\,j_4\,j_2^4\\
&\quad - 3093174\,j_6^2\,j_2^6 - 9523422720\,j_6\,j_4^4\,j_2 + 1237946976\,j_6\,j_4^3\,j_2^3 - 60328800\,j_6\,j_4^2\,j_2^5 + 1037232\,j_6\,j_4\,j_2^7\\
&\quad - 5488\,j_6\,j_2^9 + 5509980288\,j_4^6 - 1031704128\,j_4^5\,j_2^2 + 58796766\,j_4^4\,j_2^4 - 1091475\,j_4^3\,j_2^6 + 6174\,j_4^2\,j_2^8,
\end{aligned}$$

$$\begin{aligned}
0 &= 2540160\,j_{10}\,j_6 - 317520\,j_{10}\,j_4\,j_2 + 4410\,j_{10}\,j_2^3 + 1786050\,j_8^2 + 238140\,j_8\,j_6\,j_2 - 884520\,j_8\,j_4^2\\
&\quad + 133056\,j_6^2\,j_4 + 47880\,j_6\,j_4^2\,j_2 - 1092\,j_6\,j_4\,j_2^3 - 17496\,j_4^4 - 2205\,j_4^3\,j_2^2 + 49\,j_4^2\,j_2^4,
\end{aligned} \tag{15}$$

$$\begin{aligned}
0 &= 1543147200\,j_{10}\,j_8 - 793618560\,j_{10}\,j_4^2 + 34292160\,j_{10}\,j_4\,j_2^2 - 357210\,j_{10}\,j_2^4 - 144670050\,j_8^2\,j_2 + 1538248320\,j_8\,j_6\,j_4\\
&\quad - 36435420\,j_8\,j_6\,j_2^2 - 82668600\,j_8\,j_4^2\,j_2 + 1905120\,j_8\,j_4\,j_2^3 + 254016000\,j_6^3 - 10160640\,j_6^2\,j_4\,j_2 - 38808\,j_6^2\,j_2^3\\
&\quad - 50388480\,j_6\,j_4^3 + 6373080\,j_6\,j_4^2\,j_2^2 - 278460\,j_6\,j_4\,j_2^4 + 3136\,j_6\,j_2^6 - 11809800\,j_4^4\,j_2 + 643545\,j_4^3\,j_2^3 - 7497\,j_4^2\,j_2^5,\\
0 &= 4800902400\,j_{10}^2 + 262906560\,j_{10}\,j_4^2\,j_2 - 9485910\,j_{10}\,j_4\,j_2^3 + 92610\,j_{10}\,j_2^5 - 3541737150\,j_8^2\,j_4 + 37507050\,j_8^2\,j_2^2\\
&\quad - 555660000\,j_8\,j_6^2 - 94382820\,j_8\,j_6\,j_4\,j_2 + 1049580\,j_8\,j_6\,j_2^3 + 133698600\,j_8\,j_4^3 - 793800\,j_8\,j_4^2\,j_2^2 - 23756544\,j_6^2\,j_4^2\\
&\quad + 3087000\,j_6^2\,j_4\,j_2^2 + 98280\,j_6\,j_4^3\,j_2 + 47460\,j_6\,j_4^2\,j_2^3 - 980\,j_6\,j_4\,j_2^5 + 3831624\,j_4^5 + 2010015\,j_4^4\,j_2^2 - 102312\,j_4^3\,j_2^4 + 1029\,j_4^2\,j_2^6,\\
0 &= j_3 = j_5 = j_7 = j_9.
\end{aligned}$$

Furthermore, at least one among the five determinants

$$R(C_{5,2},C_{6,2},C_{7,2}),\ \ R(C_{5,2},C_{6,2},C'_{7,2}),\ \ R(C_{5,2},C_{7,2},C'_{8,2}),\ \ R(C_{5,2},C_{8,2},C_{9,2})\ \text{ and }\ R(C_{5,2},C_{6,2},C''_{9,2}) \tag{16}$$

is non-zero for a curve C with automorphism group $\mathbf{C}_4$ and if the corresponding conic has a k-rational point, then C is K-isomorphic to a curve with a k-rational model constructed with the conic and quartic method of Prop. 2.7.

Remark 3.17. To check that at least one among the five determinants of Eq. (16) is non-zero for a curve with automorphism group $\mathbf{C}_4$, we solve in a and b (over the integers to ensure that the result is still true modulo any positive prime p) the system obtained by evaluating these determinants at normal forms $x\,(x^2-1)\,(x^4+a\,x^2+b)$. We found a finite number of irreducible components, possibly defined in a k-extension, but all with an automorphism group larger than $\mathbf{C}_4$ (*cf.* Tab. 4).

Remark 3.18. In fields of characteristic 0, we can conclude with only the 4 determinants $R(C_{5,2},C_{6,2},C_{7,2})$, $R(C_{5,2},C_{7,2},C'_{8,2})$, $R(C_{5,2},C_{8,2},C_{9,2})$ and $R(C_{5,2},C_{6,2},C''_{9,2})$. These four determinants are enough in positive characteristic p too, except for a finite number of primes the smallest of which is $p=47$ (modulo 47, the point $(1:0:1:0:3:0:43:0:18)$ cancel all the determinants of Eq. (16) except $R(C_{5,2},C_{6,2},C'_{7,2})$).

$f(x) = x\,(x^2-1)\,(x^4 + a\,x^2 + b)$		Aut(C)
$a = 0$	$b = 0$	$\mathbf{U}_6, \mathbf{V}_8, \mathbf{C}_2 \times \mathbf{S}_4 \subset$ Aut(C)
any a	$b = 1$	$\mathbf{C}_2 \times \mathbf{C}_4 \subset$ Aut(C)
any a	$b^2 + (-3\,a + 1)\,b + a^3 = 0$	$\mathbf{C}_2 \times \mathbf{C}_4 \subset$ Aut(C)
$a = -6$	$b = -27$	$\mathbf{U}_6 \subset$ Aut(C)
$a = 10/3$	$b = 1$	$\mathbf{U}_6 \subset$ Aut(C)
$a = 2/9$	$b = -1/27$	$\mathbf{U}_6 \subset$ Aut(C)
$a = -6$	$b = 1$	$\mathbf{V}_8 \subset$ Aut(C)
$a^2 + 40\,a + 8 = 0$	$b = -5\,a - 1$	$\mathbf{V}_8 \subset$ Aut(C)

TABLE 4. $\mathbf{C}_4$-hyperelliptic polynomials that cancel all the determinants of Eq. (16)

The quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (15) is obtained by adjoining j_8 to $k[j_2, j_4, j_6]$ and j_8 satisfies an irreducible monic equation of degree 3. The invariant j_{10} is then rational in j_2, j_4, j_6 and j_8 (the invariants j_3, j_5, j_7 and j_9 are trivial).

Geometrically, singularities of the projective variety defined by Eq. (15) are on the curve $\mathbf{C}_2 \times \mathbf{C}_4$. Moreover, for non-zero J_2, we can set $J_2 = 1$, $J_4 = t$ for some parameter t and the intersection of the stratum with this hyperplane is a pencil a rational curves that can we parametrize with an other parameter u. So, we find for $t \neq 0$ the parametrization $(t, u) \mapsto (j_2(t,u) : 0 : j_4(t,u) : 0 : j_6(t,u) : 0 : j_8(t,u) : 0 : j_{10}(t,u))$, where

$$
\begin{aligned}
&j_2(t,u) = t\,,\quad j_4(t,u) = \frac{1}{96}\,(t+1)t^2\,,\\
&j_6(t,u) = -\frac{1}{124416000}\,(720\,u + 63\,t - 25)(-518400\,u^2 + (-90720\,t + 36000)u + 2700\,t^3 - 1269\,t^2 + 3150\,t - 625)\,,\\
&j_8(t,u) = \frac{8}{3}\,u^4 + (2/3\,t - \frac{10}{27})u^3 + (-\frac{1}{360}\,t^3 + \frac{139}{3600}\,t^2 - \frac{5}{72}\,t + \frac{25}{1296})u^2 + (-\frac{17}{14400}\,t^4 + \frac{241}{432000}\,t^3 - \frac{139}{51840}\,t^2 + \frac{25}{10368}\,t - \frac{125}{279936})u\\
&\qquad - \frac{1}{483840}\,t^6 - \frac{9707}{145152000}\,t^5 + \frac{82091}{967680000}\,t^4 - \frac{299}{18662400}\,t^3 + \frac{139}{2985984}\,t^2 - \frac{125}{4478976}\,t + \frac{625}{161243136}\,,\\
&j_{10}(t,u) = -\frac{5}{3}\,u^5 + (-\frac{31}{48}\,t + \frac{125}{432})u^4 + (-\frac{23}{3360}\,t^3 - \frac{3349}{40320}\,t^2 + \frac{155}{1728}\,t - \frac{625}{31104})u^3\\
&\qquad + (-\frac{1}{345600}\,t^4 - \frac{94723}{29030400}\,t^3 + \frac{3349}{387072}\,t^2 - \frac{775}{165888}\,t + \frac{3125}{4478976})u^2\\
&\qquad + (\frac{61}{3870720}\,t^6 + \frac{86893}{580608000}\,t^5 - \frac{735179}{13934592000}\,t^4 + \frac{105073}{418037760}\,t^3 - \frac{16745}{55738368}\,t^2 + \frac{3875}{35831808}\,t - \frac{15625}{1289945088})u + \frac{1}{44236800}\,t^7\\
&\qquad + \frac{734581}{139345920000}\,t^6 - \frac{23681183}{3344302080000}\,t^5 + \frac{736579}{401316249600}\,t^4 - \frac{108523}{24078974976}\,t^3 + \frac{83725}{24078974976}\,t^2 - \frac{19375}{20639121408}\,t + \frac{15625}{185752092672}\,.
\end{aligned}
\tag{17}
$$

The intersections of the stratum with the $J_4 = J_2^2/96$ and with the $J_2 = 0$, $J_6 = J_4$ hyperplanes are two curves which are not reachable by this parametrization. Furthermore, the rational curves $5040\,u - 579\,t - 175 = 0$ and $25401600\,u^2 + (-12186720\,t - 1764000)\,u + 185220\,t^3 - 2167119\,t^2 + 423150\,t + 30625 = 0$ are the loci of forms with multiple roots (discriminant equal to 0), the rational curve

$$
\begin{aligned}
&268738560000\,u^4 + 1492992000\,(78\,t - 25)\,u^3 - 3110400\,(90\,t^3 - 5409\,t^2 + 3900\,t - 625)\,u^2\\
&\qquad - 2880\,(130410\,t^4 - 323352\,t^3 + 405675\,t^2 - 146250\,t + 15625)\,u\\
&\qquad + 4665600\,t^6 - 55992060\,t^5 + 29516481\,t^4 - 31997700\,t^3 + 20283750\,t^2 - 4875000\,t + 390625 = 0
\end{aligned}
$$

is the locus of forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{C}_4$.

A parametrization for the intersection with $J_4 = J_2^2/96$ is of the form $t \mapsto (\,j_2(t) : 0 : j_4(t) : 0 : j_6(t) : 0 : j_8(t) : 0 : j_{10}(t)\,)$ where

$$
\begin{aligned}
&j_2(t) = t\,,\quad j_4(t) = \frac{1}{96}\,t^2\,,\quad j_6(t) = \frac{1}{7776000}\,(71\,t+2)(2183\,t^2 + 142\,t + 2)\,,\\
&j_{10}(t) = \frac{16012139}{8817984000}\,t^4 + \frac{51401}{196830000}\,t^3 + \frac{7301}{524880000}\,t^2 + \frac{31}{98415000}\,t + \frac{1}{393660000}\,,0,-\frac{905961157}{3174474240000}\,t^5\\
&\qquad - \frac{672811}{14171760000}\,t^4 - \frac{2449033}{793618560000}\,t^3 - \frac{77179}{793618560000}\,t^2 - \frac{337}{226748160000}\,t - \frac{1}{113374080000}
\end{aligned}
$$

with $t \neq 0$. The point $\left(1 : 0 : \frac{1}{96} : 0 : \frac{154993}{7776000} : 0 : \frac{16012139}{8817984000} : 0 : -\frac{905961157}{3174474240000}\right)$ is the only point on this line which is not reached by the parametrization. The points $13\,t - 14 = 0$ and $199904\,t^2 + 2296\,t - 49 = 0$, when defined, yield forms with multiple roots. The values $101\,t + 2 = 0$ and $28\,t + 1 = 0$, when defined too, yield forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{C}_4$.

Similarly, a parametrization for the intersection with $J_2 = 0$ and $J_6 = J_4$ is

$$j_2(t) = 0\,,\ \ j_4(t) = \frac{4}{3969}\,(t-9)(11\,t+6)\,,\ \ j_6(t) = \frac{4}{3969}\,(t+1)(11\,t+6)^2\,,$$
$$j_8(t) = \frac{32}{36756909}\,(373\,t^2+440\,t+127)(11\,t+6)^2\,,\ \ j_{10}(t) = -\frac{8}{992436543}\,(835\,t^2+646\,t+51)(11\,t+6)^3$$

with $11\,t+6 \neq 0$. The point $\left(0:0:\frac{4}{43659}:0:\frac{4}{43659}:0:\frac{11936}{4447585989}:0:-\frac{6680}{120084821703}\right)$, when defined, is the only point on this line which is not reached by the parametrization. Moreover, the point $t = 1/39$, when defined, yields a form with multiple roots. The roots of $667\,t^2+769\,t+302 = 0$, when defined too, yield forms with automorphism groups which contain $\mathbf{C}_2 \times \mathbf{C}_4$.

We can now easily check that the rational projective points defined on this variety are none other than the ones given by these parametrizations. Incidentally, this shows that over a finite field $\mathbb{F}_q$ of characteristic at least 11 there are $q^2-2\,q+2$ isomorphic classes of curves with automorphism group $\mathbf{C}_4$.

Remark 3.19. We are also able to deduce from these parametrizations explicit models over the field of moduli (see Section 4.5), but such expressions are far too large to be written down here.

3.2.4. *Stratum of dimension 3.* The stratum of dimension 3 corresponds to curves with automorphism group $\mathbf{D}_4$. Contrary to the other strata, we only give generators for the grevlex order $J_2 < J_3 < \ldots < J_{10}$ with weights 2, 3, ..., 10 (since generators for the lexical order are far too huge to be explicitly written here).

Lemma 3.20. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2, j_3, \ldots, j_{10})$ be its Shioda invariants. If the automorphism group of C contains $\mathbf{D}_4$, then j_2, j_3, ..., j_{10} satisfy the following 24 equations, from degree 16 up to degree 24 (cf. Appendix A for the complete set),*

$$\begin{aligned}
0 &= 49\,j_2^4\,j_4^2 - 1470\,j_2\,j_3^2\,j_4^2 - 2205\,j_2^2\,j_4^3 - 17496\,j_4^4 + 19845\,j_3\,j_4^2\,j_5 - 1092\,j_2^3\,j_4\,j_6 + 32760\,j_3^2\,j_4\,j_6 + 47880\,j_2\,j_4^2\,j_6\\
&\quad - 238140\,j_5^2\,j_6 + 133056\,j_4\,j_6^2 + 357210\,j_4\,j_5\,j_7 - 238140\,j_3\,j_6\,j_7 - 59535\,j_2\,j_7^2 - 884520\,j_4^2\,j_8 + 238140\,j_2\,j_6\,j_8 + 1786050\,j_8^2\\
&\quad + 119070\,j_3\,j_4\,j_9 + 59535\,j_2\,j_5\,j_9 - 714420\,j_7\,j_9 + 4410\,j_2^3\,j_{10} - 132300\,j_3^2\,j_{10} - 317520\,j_2\,j_4\,j_{10} + 2540160\,j_6\,j_{10},\\
0 &= -14\,j_2^3\,j_3\,j_4^2 + 420\,j_3^3\,j_4^2 + 630\,j_2\,j_3\,j_4^3 - 189\,j_2^2\,j_4^2\,j_5 + 360\,j_3\,j_4^2\,j_6 - 168\,j_2^3\,j_5\,j_6 + 5040\,j_3^2\,j_5\,j_6\\
&\quad + 3348\,j_2\,j_4\,j_5\,j_6 + 12096\,j_5\,j_6^2 + 252\,j_2^3\,j_4\,j_7 - 7560\,j_3^2\,j_4\,j_7 - 2916\,j_2\,j_4^2\,j_7 + 2268\,j_2^2\,j_6\,j_7 - 22032\,j_4\,j_6\,j_7\\
&\quad + 34020\,j_3\,j_7^2 - 68040\,j_3\,j_6\,j_8 - 34020\,j_2\,j_7\,j_8 - 42\,j_2^4\,j_9 + 1260\,j_2\,j_3^2\,j_9 + 756\,j_2^2\,j_4\,j_9 + 7776\,j_4^2\,j_9\\
&\quad - 34020\,j_3\,j_5\,j_9 - 24192\,j_2\,j_6\,j_9 + 408240\,j_8\,j_9 + 34020\,j_3\,j_4\,j_{10} + 17010\,j_2\,j_5\,j_{10} - 204120\,j_7\,j_{10},\\
0 &= 14\,j_2^7\,j_4 - 840\,j_2^4\,j_3^2\,j_4 + 12600\,j_2\,j_3^4\,j_4 - 1260\,j_2^5\,j_4^2 + 37800\,j_2^2\,j_3^2\,j_4^2 + 25191\,j_3^3\,j_4^3 + 94770\,j_2^3\,j_4^3 - 743580\,j_2\,j_4^4 + 11340\,j_2^3\,j_3\,j_4\,j_5\\
&\quad - 340200\,j_3^3\,j_4\,j_5 - 510300\,j_2\,j_3\,j_4^2\,j_5 + 76545\,j_2^2\,j_4\,j_5^2 + 1771470\,j_4^2\,j_5^2 + 546\,j_2^6\,j_6 - 32760\,j_2^3\,j_3^2\,j_6 + 491400\,j_3^4\,j_6\\
&\quad - 58446\,j_2^4\,j_4\,j_6 + 1753380\,j_2\,j_3^2\,j_4\,j_6 + 1524420\,j_2^2\,j_4^2\,j_6 - 2449440\,j_4^3\,j_6 - 3768930\,j_3\,j_4\,j_5\,j_6 - 2985255\,j_2\,j_5^2\,j_6 + 47250\,j_2^3\,j_6^2\\
&\quad - 1417500\,j_3^2\,j_6^2 + 3418200\,j_2\,j_4\,j_6^2 + 13608000\,j_6^3 + 8726130\,j_3\,j_4^2\,j_7 + 51030\,j_2^3\,j_5\,j_7 - 1530900\,j_2^2\,j_5\,j_7 + 688905\,j_2\,j_4\,j_5\,j_7\\
&\quad - 22963500\,j_5\,j_6\,j_7 + 221130\,j_2^3\,j_4\,j_8 - 6633900\,j_2^2\,j_4\,j_8 - 18676980\,j_2\,j_4^2\,j_8 + 20667150\,j_5^2\,j_8 + 61236000\,j_4\,j_6\,j_8\\
&\quad - 34020\,j_2^3\,j_3\,j_9 + 1020600\,j_3^3\,j_9 + 1530900\,j_2\,j_3\,j_4\,j_9 - 459270\,j_2^2\,j_5\,j_9 + 11022480\,j_4\,j_5\,j_9 - 30618000\,j_3\,j_6\,j_9\\
&\quad + 34020\,j_2^4\,j_{10} - 1020600\,j_2\,j_3^2\,j_{10} - 1530900\,j_2^2\,j_4\,j_{10} - 44089920\,j_4^2\,j_{10} + 13778100\,j_3\,j_5\,j_{10} + 30618000\,j_2\,j_6\,j_{10},\\
0 &= \cdots
\end{aligned} \tag{18}$$

Furthermore, a curve C with automorphism group $\mathbf{D}_4$ is K-isomorphic to the curve $y^2 = a_0\,x^8 + a_6\,x^6 + a_4\,x^4 + a_2\,x^2 + a_0$ where:

- *a_4 is the solution of the linear equation*

$$\begin{aligned}
0 = (&-11022480 j_5 j_4 j_3 + 2933280 j_4 {j_3}^2 j_2 + 784 {j_2}^6 + 55339200 {j_6}^2 + 4408992 {j_4}^3 + 705600 {j_3}^4 - 36741600 j_9 j_3 - 4286520 {j_5}^2 j_2\\
&+ 32315760 j_6 j_4 j_2 + 36741600 j_{10} j_2 - 64297800 j_7 j_5 + 10054800 j_6 {j_3}^2 - 335160 j_6 {j_2}^3 - 47040 {j_3}^2 {j_2}^3 + 2812320 {j_4}^2 {j_2}^2\\
&- 97776 j_4 {j_2}^4)X - 410791500 j_7 j_6 - 10716300 j_9 {j_2}^2 + 4365900 j_7 {j_2}^3 + 150793650 j_6 j_4 j_3 + 39590775 j_6 j_5 j_2 - 196465500 j_7 j_4 j_2\\
&+ 10716300 {j_4}^2 j_3 j_2 + 1205583750 j_8 j_5 - 1157360400 j_9 j_4 + 254435580 j_5 {j_4}^2 + 321489000 j_{10} j_3 - 75014100 {j_5}^2 j_3 - 130977000 j_7 {j_3}^2\\
&+ 7144200 j_4 {j_3}^3 + 5120010 j_5 j_4 {j_2}^2 - 238140 j_4 j_3 {j_2}^3 - 185220 j_5 {j_2}^4 + 5556600 j_5 {j_3}^2 j_2 \quad \text{if this equation is non trivial,}
\end{aligned} \tag{19}$$

$$\begin{aligned}
0 = (&-483840 j_4 {j_3}^2 j_2 + 2449440 j_5 j_4 j_3 - 112 {j_2}^6 - 9072000 {j_6}^2 - 629856 {j_4}^3 - 100800 {j_3}^4 + 612360 {j_5}^2 j_2 - 6713280 j_6 j_4 j_2 + 9185400 j_7 j_5\\
&- 2797200 j_6 {j_3}^2 + 93240 j_6 {j_2}^3 + 6720 {j_3}^2 {j_2}^3 - 498960 {j_4}^2 {j_2}^2 + 16128 j_4 {j_2}^4 + 18370800 j_8 j_4)X + 89302500 j_7 j_6 + 26460 {j_5}^2 j_2^4\\
&- 2381400 {j_4}^2 j_3 j_2 - 22027950 j_6 j_4 j_3 - 11013975 j_6 j_5 j_2 + 41079150 j_7 j_4 j_2 + 10716300 {j_5}^2 j_3 - 241116750 j_8 j_5 + 257191200 j_9 j_4\\
&- 56030940 j_5 {j_4}^2 + 23814000 j_7 {j_3}^2 - 1587600 j_4 {j_3}^3 - 793800 j_7 {j_2}^3 - 793800 j_5 {j_3}^2 j_2 - 476280 j_5 j_4 {j_2}^2 + 52920 j_4 j_3 {j_2}^3 \quad \text{otherwise;}
\end{aligned} \tag{20}$$

- a_0 *is any root of the quadratic equation*

$$\begin{aligned}0 = (&-529079040 j_5 j_4 j_3 + 1551156480 j_6 j_4 j_2 + 140797440 j_4 {j_3}^2 j_2 - 2257920 {j_3}^2 {j_2}^3 - 4693248 j_4 {j_2}^4 - 1763596800 j_9 j_3\\ &+ 1763596800 j_{10} j_2 + 2656281600 {j_6}^2 + 33868800 {j_3}^4 + 37632 {j_2}^6 + 211631616 {j_4}^3 + 134991360 {j_4}^2 {j_2}^2 - 16087680 j_6 {j_2}^3\\ &- 205752960 {j_5}^2 j_2 + 482630400 j_6 {j_3}^2 - 3086294400 j_7 j_5) X^2 - 14647500 j_4 {j_3}^2 {j_2}^2 - 307355310 j_6 j_4 {j_2}^2 - 140277690 j_6 {j_3}^2 j_2\\ &- 382571910 j_7 j_4 j_3 + 1130212440 j_6 j_5 j_3 - 105019740 j_8 j_4 j_2 + 555660 j_5 j_3 {j_2}^3 + 206671500 j_9 j_3 j_2 + 36996750 j_5 j_4 j_3 j_2\\ &+ 3472081200 {j_7}^2 - 3724 {j_2}^7 + 507952620 j_7 j_5 j_2 + 8024365440 j_{10} j_4 + 530456850 j_5 {j_2}^2 j_4 - 472252032 j_6 {j_4}^2 - 4089340080 j_9 j_5\\ &+ 39293100 {j_4}^2 {j_3}^2 - 16669800 j_5 {j_3}^3 - 400029840 {j_6}^2 j_2 - 114388848 {j_4}^3 j_2 - 3351600 {j_3}^4 j_2 - 206671500 j_{10} {j_2}^2 + 27862380 {j_5}^2 {j_2}^2\\ &+ 10835370 j_8 {j_2}^3 - 15739920 {j_4}^2 {j_2}^3 + 4675923 j_6 {j_2}^4 + 223440 {j_3}^2 {j_2}^4 + 488250 j_4 {j_2}^5 - 325061100 j_8 {j_3}^2 - 13974055200 j_8 j_6\end{aligned} \tag{21}$$

if this equation is non trivial,

$$\begin{aligned}0 = (&117573120 j_5 j_4 j_3 - 322237440 j_6 j_4 j_2 - 23224320 j_4 {j_3}^2 j_2 + 322560 {j_3}^2 {j_2}^3 + 774144 j_4 {j_2}^4 + 881798400 j_8 j_4 - 435456000 {j_6}^2\\ &- 4838400 {j_3}^4 - 5376 {j_2}^6 - 30233088 j_4 {j_3}^3 - 23950080 {j_4}^2 {j_2}^2 + 4475520 j_6 {j_2}^3 + 29393280 {j_5}^2 j_2 - 134265600 j_6 {j_3}^2 + 440899200 j_7 j_5) X^2\\ &+ 151099830 j_7 j_4 j_3 + 2457000 j_4 {j_3}^2 {j_2}^2 + 34838370 j_6 {j_3}^2 j_2 - 257905620 j_6 j_5 j_3 - 184779630 j_8 j_4 j_2 - 79380 j_5 j_3 {j_2}^3 - 72564660 j_7 j_5 j_2\\ &- 10206000 j_5 j_4 j_3 j_2 - 868020300 {j_7}^2 + 532 {j_2}^7 - 2469035520 j_{10} j_4 - 96446700 {j_5}^2 j_4 + 62215776 j_6 {j_4}^2 + 1080203040 j_9 j_5\\ &+ 66418380 j_6 j_4 {j_2}^2 + 2381400 j_5 {j_3}^3 + 20230560 {j_6}^2 j_2 + 15422724 {j_4}^3 j_2 + 478800 {j_3}^4 j_2 - 3980340 {j_5}^2 {j_2}^2 - 1547910 j_8 {j_2}^3\\ &+ 2999430 {j_4}^2 {j_2}^3 - 1161279 j_6 {j_2}^4 - 31920 {j_3}^2 {j_2}^4 - 81900 j_4 {j_2}^5 + 46437300 j_8 {j_3}^2 + 3815002800 j_8 j_6 - 11736900 {j_4}^2 {j_3}^2 \quad \text{otherwise;}\end{aligned} \tag{22}$$

- a_2 *is any root of*

$$\begin{aligned}0 &= 15750\, a_0\, X^4 + (105000\, a_0^2\, a_4 + 510\, a_4^3 - 23100\, a_4\, j_2 - 686000\, j_3)\, X^2 + 705600\, a_0^3\, a_4^2 - 10804500\, a_0^3\, j_2\\ &\quad + 3024\, a_0\, a_4^4 - 244755\, a_0\, a_4^2\, j_2 - 1440600\, a_0\, a_4\, j_3 + 15126300\, a_0\, j_4 + 2881200\, a_0\, j_2^2 \quad \text{if } a_0 \neq 0,\\ 0 &= X - 1 \quad \text{otherwise;}\end{aligned}$$

- *and* a_6 *is any root of*

$$\begin{aligned}0 &= 5\, a_2\, X + 140\, a_0^2 + a_4^2 - 70\, j_2 \quad \text{if } a_2 \neq 0,\\ 0 &= 1575\, a_0\, X^2 - 24\, a_4^3 + 2940\, a_4\, j_2 - 68600\, j_3 \quad \text{otherwise.}\end{aligned}$$

Similarly to the automorphism group $\mathbf{C}_2^3$, Lemma 3.14 yields most of the time a model over an extension of k (of degree at most 8, see Section 4 for the rationality issue).

Remark 3.21. In case the j_is correspond to polynomials with multiple roots, it may happen that some of equations in Lemma 3.20 for a_4, a_0, a_2 or a_6 are inconsistent (typically '$0 = 1$'). It means that none of the polynomials $a_0\, x^8 + a_6\, x^6 + a_4\, x^4 + a_2\, x^2 + a_0$ have such invariants. Instead, we construct polynomials of the form $f(x) = a_8\, x^8 + a_6\, x^6 + a_4\, x^4 + a_2\, x^2$ where a_4 is the root of

$$\begin{aligned}0 = (&11741206932\, j_7 + 7172848242\, j_4\, j_3 - 407219400\, j_3\, {j_2}^2 - 340313967\, j_5\, j_2) X - 770798216160\, j_8 + 48652305408\, j_6\, j_2\\ &- 21652069824\, {j_4}^2 - 15830753400\, j_5\, j_3 + 1196327160\, j_4\, {j_2}^2 - 77421296\, {j_2}^4 - 6442857120\, {j_3}^2 j_2 \quad \text{if this equation is non trivial,}\\ 0 = (&488750830848\, {j_4}^2 + 19912489511040\, j_8 + 119825959260\, {j_3}^2 j_2 + 272672590848\, j_4\, {j_2}^2 - 11523989242\, {j_2}^4) X\\ &+ 663046656007680\, j_9 + 29540232087660\, j_7\, j_2 - 147027580680960\, j_5\, j_4 - 238603106730240\, j_6\, j_3\\ &- 32453930490450\, j_4\, j_3\, j_2 + 5621022644895\, j_5\, {j_2}^2 + 590440221000\, j_3\, {j_2}^3 j_2 \quad \text{otherwise, if this equation is non trivial,}\\ 0 = (&7023805824\, j_6\, j_2 - 83616540\, {j_3}^2 j_2 + 1090412288\, j_4\, {j_2}^2 + 1936166400\, {j_4}^2 - 40257382\, {j_2}^4) X + 231759118080\, j_9\\ &- 96001584000\, j_5\, j_4 - 23998355150\, j_4\, j_3\, j_2 - 191099623680\, j_6\, j_3 - 32526010860\, j_7\, j_2 + 310611000\, j_3\, {j_2}^3 - 617600655\, j_5\, {j_2}^2\\ &\text{otherwise, if this equation is non trivial,}\\ 0 = X \quad &\text{otherwise;}\end{aligned}$$

and

$$a_2 = 1\,,\ a_6 = -({a_4}^2 - 70\, j_2)/5a_2\,,\ a_8 = -17\, {a_4}^3/525 + 22\, a_4\, j_2/15 + 392\, j_3/9\,.$$

Remark 3.22. It may happen that a curve C annihilates both Eq. (19) and Eq. (20) too, so that we can not determine a_4. But, when it happens, the automorphism group is larger than $\mathbf{D}_4$ (see Table 5) and so, lemmas for larger automorphism group apply.

Curves $C : y^2 = a_0\, x^8 + a_6\, x^6 + a_4\, x^4 + a_2\, x^2 + a_0$	Singular	Aut C
$a_0 = 0$	Yes	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$a_6 + a_2 = 0$	No	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$a_6 - a_2 = 0$	No	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$\begin{cases} 308 a_0 a_2 a_6 a_4 - 224 a_0 {a_4}^3 + 3 {a_2}^3 a_6 + 15 {a_2}^2 {a_4}^2 + 3 a_2 {a_6}^3 + 15 {a_6}^2 {a_4}^2 = 0 \\ 21 a_0 {a_2}^2 + 21 a_0 {a_6}^2 + 11 a_2 a_6 a_4 - 8 {a_4}^3 = 0 \\ 196 {a_0}^2 - a_2 a_6 - 5 {a_4}^2 = 0 \end{cases}$	No	$\mathbf{D}_{12} \subset \operatorname{Aut} C$

TABLE 5. Automorphism groups of curves C which cancel both Eq. (19) and Eq. (20)

Remark 3.23. It may happen that a curve C cancels both Eq. (21) and Eq. (22) too, so that we can not determine a_0. But again, when it happens, the automorphism group is larger than $\mathbf{D}_4$ (see Table 6) and so, lemmas for larger automorphism group apply.

Curves $C : y^2 = a_0\,x^8 + a_6\,x^6 + a_4\,x^4 + a_2\,x^2 + a_0$	Singular	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$a_0 = 0$	Yes	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$a_6 + a_2 = 0$	No	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$a_6 - a_2 = 0$	No	$\mathbf{C}_2^3 \subset \operatorname{Aut} C$
$\begin{cases} 308a_0a_2a_6a_4 - 224a_0{a_4}^3 + 3{a_2}^3a_6 + 15{a_2}^2{a_4}^2 + 3a_2{a_6}^3 + 15{a_6}^2{a_4}^2 = 0 \\ 21a_0{a_2}^2 + 21a_0{a_6}^2 + 11a_2a_6a_4 - 8{a_4}^3 = 0 \\ 196{a_0}^2 - a_2a_6 - 5{a_4}^2 = 0 \end{cases}$	No	$\mathbf{D}_{12} \subset \operatorname{Aut} C$
$\begin{cases} 14a_0 + 2a_6 - a_4 = 0 \\ 14a_0 + 2a_2 - a_4 = 0 \end{cases}$	No	$\mathbf{C}_2 \times \mathbf{D}_8 \subset \operatorname{Aut} C$
$\begin{cases} 14a_0 - 2a_6 - a_4 = 0 \\ 14a_0 - 2a_2 - a_4 = 0 \end{cases}$	No	$\mathbf{C}_2 \times \mathbf{D}_8 \subset \operatorname{Aut} C$

TABLE 6. Automorphism groups of curves C which cancel both Eq. (21) and Eq. (22)

Here, the quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (18) is obtained by adjoining j_6 to $k[j_2, j_3, j_4, j_5]$ and j_6 satisfies an irreducible monic equation of degree 10. The invariants j_7, j_8, j_9 and j_{10} are then rational in j_2, j_3, j_4 and j_5.

Geometrically and contrary to the other strata, we did not find any rational intersection with trivial hyperplane of the projective variety defined by Eq. (18). For instance, intersecting with $j_2 = j_3 = 0$ is geometrically birationally equivalent to the elliptic curve $y^2 = x^3 - x$, of j-invariant 1728 (the curve with complex multiplication by $\sqrt{-1}$). Nevertheless, it is straightforward to adapt the enumeration algorithm at the end of Section 1.5 to the case of a field of degree 10 over $k[j_2, j_3, j_4, j_5]$ (instead of a field of degree 5 over $k[j_2, j_3, j_4, j_5, j_6, j_7]$ for the full $\mathcal{I}_8$), in order to find non-trivial curves with automorphism group contained in $\mathbf{D}_4$.

At last, we emphasize that we have encountered unexpected computational difficulties to check with MAGMA that a reconstructed model for $\mathbf{D}_4$ has the same weighted projective class as the 9-uple $(j_2 : j_3 : \ldots : j_{10})$ provided in input. Our first implementations took about one week only to define the degree 8 extensions needed for the reconstructed model (due to some unexpected internal Gröbner basis computations) that we had to design a more optimized version. We perform by 'hand', *i.e.* directly over the quotient field of $\mathcal{I}_8$ modulo the ideal defined by Eq. (18), the calculation of the Shioda invariants of the reconstructed model. This requires explicit expressions for each coordinate of the Shioda invariants in the basis 1, a_0, a_2, $a_0\,a_2$, a_2^2,..., $a_0a_2^3$. Moreover, we only check the equality of j_2, ..., j_6 since j_7, j_8, j_9 and j_{10} are rational in j_2, j_3, j_4, j_5 and j_6. The corresponding routine took about half a day on a computer to check finally that everything was fine.

3.2.5. *Generic stratum.* Our main tool for reconstructing curves C with automorphism group $\mathbf{C}_2$ is the conic and quartic method developed in Proposition. 2.8. It requires that C does not cancel all the 364 fundamental conic determinants $R(q_1, q_2, q_3)$ (or equivalently Eq. (18)), otherwise Algorithm 3 fails. To check that this cannot happen, we have computed a Gröbner basis (for the grevlex order $J_2 < J_3 < \ldots < J_{10}$ with weights 2, 3, ..., 10) of a polynomial system which concatenates Relations (1) and the *fundamental determinants* $R(q_1, q_2, q_3)$ with $\{q_1, q_2, q_3\}$ any subset of size 3 of the set of 14 covariants of order 2 given in Tab. 2. From Lemma 3.2, these 364 fundamental determinants all vanish for curves with automorphism group contained in $\mathbf{D}_4$. Surprisingly, the Gröbner basis that we obtain is equal to the system of 24 polynomials (18) of Lemma 3.20.

After some further Gröbner computations, still for the 'grevlex' order $J_2 < J_3 < \ldots < J_{10}$ with weights 2, 3, ..., 10, we observe that we can reduce the set of 364 fundamental conic determinants $R(q_1, q_2, q_3)$ to a subset of only 19 elements, since the determinants of these 19 conics generate the same ideal as the one defined by Eq. (18).

Lemma 3.24. *Let C be a hyperelliptic curve of genus 3 over k, let $(j_2 : j_3 : \ldots : j_{10})$ be its Shioda invariants. If j_2, j_3, ... j_{10} do not cancel Eq. (18), nor Eq. (15) and Eq. (7), then its automorphism group is $\mathbf{C}_2$. Some*

of the 364 *fundamental determinants are non-zero among which at least one of the following* 19 *determinants,*

$$\begin{array}{lllll} R(C_{5,2},C_{6,2},C_{7,2}), & R(C_{5,2},C_{8,2},C_{9,2}), & R(C_{5,2},C_{7,2},C'_{8,2}), & R(C_{5,2},C_{6,2},C''_{9,2}), & R(C_{5,2},C'_{7,2},C'_{9,2}),\\ R(C_{5,2},C_{6,2},C'_{7,2}), & R(C_{5,2},C'_{8,2},C_{11,2}), & R(C_{5,2},C_{7,2},C_{11,2}), & R(C_{5,2},C_{7,2},C'_{11,2}), & R(C_{5,2},C'_{7,2},C'_{10,2}),\\ R(C_{6,2},C'_{7,2},C_{9,2}), & R(C_{5,2},C_{7,2},C_{10,2}), & R(C_{5,2},C_{7,2},C_{9,2}), & R(C_{5,2},C_{6,2},C'_{10,2}), & R(C_{5,2},C_{6,2},C_{10,2}),\\ R(C_{5,2},C'_{7,2},C'_{8,2}), & R(C_{5,2},C_{6,2},C'_{9,2}), & R(C_{5,2},C_{6,2},C'_{8,2}), & R(C_{5,2},C_{6,2},C_{8,2}). & \end{array}$$

With one of the corresponding non-singular conics, we can use Proposition. 2.8 to reconstruct the curve over k*.*

There are examples of curves with automorphism group $\mathbf{C}_2$ for which many determinants R are zero.

- The curve defined over $\mathbb{Q}$ by

$$y^2 = 2581381040\,x^8 + 7704021083264\,x^7 + 101567018399840\,x^6 - 96172789044745280\,x^5 - 1962596803409291000\,x^4 - 15122894980514300000\,x^3 + 266225999081701799000\,x^2 + 3116782046067938990000\,x + 191614097302577354466875$$

  has Shioda invariants in the class $(0:1:0:1:0:0:0:0:0:0)$ for which the only non-zero determinants in the list of Lemma 3.24 are $R(C_{5,2},C_{6,2},C'_{9,2})$ and $R(C_{5,2},C_{6,2},C''_{9,2})$. These two discriminants are zero modulo 19 and 113 too, but then the automorphism group is larger than $\mathbf{C}_2$.
- Similarly, the singular form $8\,x^5z^3 - 125z^8$ has Shioda invariants in the class $(0:0:0:1:0:0:0:0:0)$ (resp. $16\,x^7z - 35\,x^2z^6$, in the class $(0:0:0:1:0:0:0:0:3/32)$) for which 18 determinants in the list are zero in fields of characteristic 17 (resp. characteristic 13). Note that over $\mathbb{Q}$, these two forms already have 17 determinants in the list which are zero.

4. Field of definition

4.1. **Field of definition and field of moduli: general facts.** Let K be an algebraically closed field and let $k \subset K$ be a subfield. Let C be a curve defined over K of genus $g \geq 2$.

Definition 4.1. We say that k is a *field of definition* of C if there exists a curve $\mathcal{C}/k$ such that $\mathcal{C}$ is K-isomorphic to C. The curve $\mathcal{C}/k$ is a model of C over k.
The *field of moduli* of C, denoted $\mathbf{M}_C$, is the intersection of all fields of definition of C.

This is the definition given by [Koi72]. Usually, one defines the field of moduli of a curve C as the subfield K^H of K fixed by

$$H = \{\sigma \in \mathrm{Aut}(K),\ C \simeq {}^{\sigma}C\}.$$

As shown in [Koi72] (or [Hug05, Th.1.5.8]), K^H is a purely inseparable extension of $\mathbf{M}_C$. Hence, when $\mathbf{M}_C$ is a perfect field, both notions coincide.
Another common definition is field of moduli relative to a Galois extension K/k, as used for instance in [DE99]. In [HH03], it is proved that there is always a field of definition for C which is a finite extension of $\mathbf{M}_C$ and [Hug05, Cor.1.5.9] shows that this extension can be chosen separable. Hence, $\mathbf{M}_C$ is the field of moduli relative to the Galois extension $\mathbf{M}_C^{\mathrm{sep}}/\mathbf{M}_C$.

Remark 4.2. As a consequence of Torelli theorem (see for instance the formulation of [Mat58]), C can be defined over k if and only if the principally polarized abelian variety $\mathrm{Jac}\,C$ can be. Hence the field of moduli of C coincides with the field of moduli of $\mathrm{Jac}\,C$. Note however that, when C is non hyperelliptic, a model A of $\mathrm{Jac}\,C$ over k is not necessarily the Jacobian of a curve over k (see [LRZ10]).

A classical problem is to know whether $\mathbf{M}_C$ is a field of definition or if there is an obstruction. One finds several sufficient conditions in the literature. For instance, the moduli field $\mathbf{M}_C$ is a field of definition

- when C has no automorphism (or more generally when $\mathrm{Aut}(C)$ has no center and has a complement in the automorphism group of $\mathrm{Aut}(C)$ [DE99, Cor.4.3]);
- when K is the algebraic closure of a finite field [Hug07, Cor.2.11];
- when $K \subset \mathbb{C}$ and $\mathrm{Jac}(C)$ has complex multiplication [Mil72, Mil74].

The difference between a field of moduli and a field of definition boils down to Weil's cocycle relations.

Theorem 4.3 ([Wei56, proof of Th.1]). *Assume that K/k is a Galois extension. Then k is a field of definition of C if and only if there exists a finite extension k' of k such that for all $\sigma \in \mathrm{Gal}(K/k)$, there exists a k'-isomorphism $f_\sigma : C \to {}^\sigma C$ such that for all $\sigma, \tau \in \mathrm{Gal}(K/k)$,*

$$f_\tau^\sigma \circ f_\sigma = f_{\tau\sigma}.$$

Assume now that $K/\mathbf{M}_C$ is a Galois extension and let $\Gamma := \mathrm{Gal}(K/\mathbf{M}_C)$. By definition of the field of moduli, for all $\sigma \in \Gamma$, there exists, a K-isomorphism $F_\sigma : C \to {}^\sigma C$. But only when C has a trivial automorphism group, this forces Weil's cocycle relations. Following this remark, we consider the curve $B = C/\operatorname{Aut}(C)$. The isomorphism F_σ induces an isomorphism $f^\sigma : B \to {}^\sigma C/\operatorname{Aut}({}^\sigma C) = {}^\sigma B$ and the diagram

$$\begin{array}{ccc} C & \xrightarrow{F_\sigma} & {}^\sigma C \\ \downarrow & & \downarrow \\ B & \xrightarrow{f_\sigma} & {}^\sigma B \end{array}$$

is commutative. The curve B admits a model over $\mathbf{M}_C$. Indeed we have to check the cocycle relations

$$\forall \sigma, \tau \in G,\ f_\tau^\sigma \circ f_\sigma = f_{\tau\sigma}.$$

But

$$F_{\tau\sigma}^{-1} \circ (F_\tau^\sigma \circ F_\sigma) \in \operatorname{Aut}(C)$$

so $f_{\tau\sigma}^{-1} \circ (f_\tau^\sigma \circ f_\sigma) = \mathrm{id}$. Hence there exists a model $\mathcal{B}/\mathbf{M}_C$ and a K-isomorphism $\phi : B \to \mathcal{B}$ such that for all $\sigma \in \Gamma$, $f_\sigma = (\phi^{-1})^\sigma \circ \phi$. One of the main results of [DE99] is the following.

Theorem 4.4 ([DE99, Cor.4.3(c)]). *If $\#\operatorname{Aut}(C)$ is coprime to p and $\mathcal{B}(\mathbf{M}_C) \neq \emptyset$ then $\mathbf{M}_C$ is a field of definition of C.*

If $\mathcal{B}$ is a genus 0 curve, its canonical divisor is of degree -2. Hence to get a rational point on $\mathcal{B}$, it is enough that $\mathcal{B}$ has a rational divisor of odd degree [Hug05, Lem.4.0.4] (this is the case for instance if $\mathcal{B}(k') \neq \emptyset$ for $k'/\mathbf{M}_C$ an extension of odd degree [Hug05, Lem.4.0.5]).

Finally, the field of moduli is related to the moduli space as follows. Assume that the genus g of C is greater than or equal to 2 and let M_g be the coarse moduli space of curves of genus g viewed as a scheme over the prime field of K. The curve C gives a morphism $\operatorname{Spec} K \to \mathsf{M}_g$ whose image $[C]$ is a closed point of M_g. Let $K([C])$ be the residue field at $[C]$. If $K([C])$ is perfect then we have the equality $\mathbf{M}_C = K([C])$. Note that counterexamples are known if $K([C])$ is not perfect (in that case $\mathbf{M}_C$ is a purely inseparable extension of $K([C])$, see [Sek85]).

4.2. **The hyperelliptic case.** The problem to know if the field of moduli is a field of definition has been first addressed by Mestre in [Mes91] under the conditions that the the genus is even. Note that in this case, he showed [Mes91, p.322], that if a hyperelliptic curve C is defined over a field k, it has a hyperelliptic equation (in the sense of Section 1.3) over k. However, in general, this is not true and we will have to distinguish two problems: is the field of moduli a field of definition ? And if so, has the model over the field of moduli a hyperelliptic equation ? This motivates the following terminology.

Definition 4.5. An hyperelliptic curve C over a field k can be *hyperelliptically defined* over a subfield $k \subset K$ if there exists a model of C over k which has a hyperelliptic equation.

From now on, let K be an algebraically closed field of characteristic $p \neq 2$. Let $g \geq 2$ be an integer, $n = 2g + 2$ and let $f \in K[x]$ be a hyperelliptic polynomial of degree n. Let C be the hyperelliptic curve of genus g defined by the equation $y^2 = f(x)$. In the sequel we assume that K is a Galois extension of $\mathbf{M}_C$ and that $K([C])$ (and $\mathbf{M}_C$) are perfect fields (we are typically interested in the cases $p = 0$ or K is the Galois closure of a finite field). We denote $\Gamma := \mathrm{Gal}(K/\mathbf{M}_C)$.

From the form of the isomorphisms between hyperelliptic equations and Weil cocycle relations, we get the following result.

Lemma 4.6. *C has a model over k if and only if there exists a finite Galois extension k'/k such that for all $\sigma \in \mathrm{Gal}(k'/k)$, there exist $(M_\sigma, e_\sigma) \in (\mathrm{GL}_2(k'), k'^*)$ such that*

- *(M_σ, e_σ) is an isomorphism between C and ${}^\sigma C$;*
- *for all $\sigma, \tau \in \mathrm{Gal}(k'/k)$, there exists $\lambda \in k'^*$ such that*

$$M_{\tau\sigma} = \lambda M_\tau^\sigma M_\sigma, \quad e_{\tau\sigma} = \lambda^{g+1} e_\tau^\sigma e_\sigma.$$

If we wish to obtain a model with a hyperelliptic equation, we get the following version of the lemma.

Lemma 4.7. *C can be hyperelliptically defined over k if and only if there exists a finite Galois extension k'/k such that for all $\sigma \in \mathrm{Gal}(k'/k)$, there exist $M_\sigma \in \mathrm{GL}_2(k')$ such that*

- *there exists $e_\sigma \in k'^*$ such that (M_σ, e_σ) is an isomorphism between C and ${}^\sigma C$;*
- *for all $\sigma, \tau \in \mathrm{Gal}(k'/k)$,*

$$M_{\tau\sigma} = M_\tau^\sigma M_\sigma.$$

Proof. If C can be hyperelliptically defined over k, there exists a curve $\mathcal{C} : y^2 = \tilde{f}(x)$ with $\tilde{f} \in k[x]$, a finite extension k'/k and $(A, a) \in (\mathrm{GL}_2(k'), k'^*)$ such that (A, a) is a k'-isomorphism between C and $\mathcal{C}$. Defined then $M_\sigma = (A^\sigma)^{-1} A$ and $e_\sigma = e/e^\sigma$. One can check that the (M_σ, e_σ) satisfy the hypotheses.
Conversely, we can assume that there is a finite Galois extension k''/k containing k' and a splitting field of f. We denote r_i the roots of f. Following the proof of Hilbert 90 as in [Ser68, p.159,Prop.3], we can find a matrix $P \in M_2(k'')$ such that the matrix

$$A = \sum_{\tau \in \mathrm{Gal}(k''/k)} P^\tau M_\tau.$$

is invertible. Now for any lift $\bar{\sigma}$ of an element σ in $\mathrm{Gal}(k''/k)$ to Γ, we get

$$A^{\bar{\sigma}} = \sum_{\tau \in \mathrm{Gal}(k''/k)} P^{\tau\sigma} M_\tau^\sigma = \left(\sum_{\tau \in \mathrm{Gal}(k''/k)} P^{\tau\sigma} M_{\tau\sigma} \right) M_\sigma^{-1} = A M_\sigma^{-1}$$

since P and the M_τ are defined over k''. Let us denote $\tilde{r}_i = A.r_i$. For all $\sigma \in \mathrm{Gal}(k''/k)$, letting $\sigma(\infty) = \infty$, we get

$$\{\tilde{r}_i^\sigma\} = \{A^\sigma . r_i^\sigma\} = \{A M_\sigma^{-1} . r_i^\sigma\} = \{A.r_i\} = \{\tilde{r}_i\}.$$

Hence the polynomial $\tilde{f} = \prod (x - \tilde{r}_i) \in k[x]$ and the curve $\mathcal{C}/k : y^2 = \tilde{f}(x)$ is a model of C. □

Remark 4.8. The previous lemmas give an easy proof that for g even, C can be defined over k if and only if C can be hyperelliptically defined over k. Indeed, assuming the cocycle condition of Lemma 4.6 we can write

$$\det M_{\tau\sigma} = \lambda^2 (\det M_\tau)^\sigma (\det M_\sigma), \quad e_{\tau\sigma} = \lambda (\lambda^2)^{g/2} e_\tau^\sigma e_\sigma.$$

Hence we can obtain a good representative satisfying the cocycle condition of Lemma 4.7 letting

$$M'_\sigma = e_\sigma^{-1} \cdot (\det M_\sigma)^{g/2} \cdot M_\sigma.$$

Let now I_1 and I_2 be two homogeneous invariants of the same degree, of degree $2g+2$ binary forms, defined over the prime field of K and $I_2(f) \neq 0$. Their quotient $\iota = I_1/I_2$ satisfies $\iota(f) = \iota(M.f)$ for all $M \in \mathrm{GL}_2(K)$. The function ι is therefore a function on the moduli space of hyperelliptic curves of genus g. Hence $\iota(f) \in \mathbf{M}_C$. More specifically, we can prove the following result.

Proposition 4.9. *Let $\mathcal{J} = (I_1 : \ldots : I_m)$ be a m-uple of invariants of degree d_i, of degree n binary forms, and suppose each I_i is defined over the prime field of K. Then, there exists $\lambda \in K^*$ such that for $1 \leq i \leq m$, $I_i(f)/\lambda^{d_i} \in \mathbf{M}_C$.*

Proof. Let d be the gcd of the degree d_i of the invariants I_i which value at f is not zero. Then there exists $c_i \in \mathbb{Z}$ with $c_i = 0$ if $I_i(f) = 0$ such that $\sum c_i d_i = d$. We then define $I = \prod_i I_i^{c_i}$. From what we said above, the value $\mu_i = I_i(f)/I(f)^{d_i/d} \in \mathbf{M}_C$. If we let $\lambda = I(f)^{1/d}$ for any choice of a dth root of $I(f)$ we get the result. □

Note that the corresponding representative $(I_1(f)/\lambda^{d_1}, \ldots, I_m(f)/\lambda^{d_m})$ for the m-uple of invariants is the one constructed in Section 1.4 and used in our algorithms.

Corollary 4.10. *With the notation of Section 2.2, there exists $\lambda \in K^*$ such that the change of variables $x_i = \lambda^{e_i} \cdot x'_i$ for some integers e_1, e_2, e_3 makes the conic $\mathcal{Q}$ and the curve $\mathcal{H}$ defined over $\mathbf{M}_C$.*

Proof. Thanks to Proposition 4.9 we can find $\lambda \in K^*$ for the uple of all the A_{ij} and h_I. Let d_1, d_2, d_3 be the degree of q_1, q_2 and q_3. From the formula for the coefficients A_{ij} and h_I, one sees easily that $\deg(A_{ij}) + \deg(x_i^*) + \deg(x_j^*) = 2(d_1+d_2+d_3)$ is a constant c and $\deg(h_I) + \sum_{i\in I} \deg(x_i^*) = 1 + n/2 \cdot d_1 \cdot d_2 \cdot d_3$ is a constant c'. Hence, if we make the change of variables $x_i = \lambda^{e_i} \cdot x'_i$ where $e_i = \deg(x_i^*)$, we get that

$$\sum A_{ij} \lambda^{\deg(x_i^*)+\deg(x_j^*)} x'_i x'_j = \lambda^c \cdot \sum \left[A_{ij} / \lambda^{\deg(A_{ij})} \right] x_i x_j$$

and similarly

$$\sum_I h_I x_I = \lambda^{c'} \cdot \sum \left[h_I / \lambda^{\deg(h_I)} \right] x'_I.$$

This gives the result. □

We then find a refinement of Proposition 2.8.

Proposition 4.11. *Let (q_1, q_2, q_3) be three covariants of order 2 for degree n binary forms and assume they are defined over the prime field of K. If $R(q_1, q_2, q_3)$ evaluated at a form f in non-zero, there exists a non-singular conic $\mathcal{Q}$ and a plane curve $\mathcal{H}$ of degree $n/2$ defined over $\mathbf{M}_C$ such that there is a K-isomorphism $\mathcal{Q} \to \mathbb{P}^1$ mapping the intersections points of $\mathcal{Q} \cap \mathcal{H}$ on the roots of $f(X, Z)$. In particular the field of moduli is a field of definition if $\mathcal{Q}$ has a $\mathbf{M}_C$-rational point and in this case C can be hyperelliptically defined over $\mathbf{M}_C$.*

4.3. **The hyperelliptic case with no extra-automorphism.** Besides the hypotheses of Section 4.2, we assume that $\mathrm{Aut}(C) = \langle \iota \rangle$. In particular $C/\langle \iota \rangle = B$ is the genus 0 curve Q defined in Section 1.3.
Let D be the image of the ramification divisor W of C by the map $\phi \circ \rho : C \to B \to \mathcal{B}$. The divisor D is defined over $\mathbf{M}_C$. Indeed for any $\sigma \in \Gamma$

$${}^\sigma D = \phi^\sigma \circ \rho^\sigma({}^\sigma W) = \phi^\sigma \circ \rho^\sigma(F_\sigma(W)) = \phi^\sigma \circ f_\sigma \circ \rho(W) = \phi \circ \rho(W) = D.$$

Assume now that there exists $\Phi : C \to C'$ an isomorphism onto a model $C'/\mathbf{M}_C$ of C. Let Q' be the quotient of C' by the hyperelliptic involution and D' the image of the ramification divisor W' on C'. The induced K-isomorphism $\tilde{\Phi} : \mathcal{B} \to Q'$ maps the divisor D onto D'. Now for any $\sigma \in \Gamma$, we have $\tilde{\Phi}^{-1} \circ \tilde{\Phi}^\sigma(D) = D$. Since C has no extra-automorphism, this means that $\tilde{\Phi}^{-1} \circ \tilde{\Phi}^\sigma = \mathrm{id}$ hence $\tilde{\Phi}$ is defined over $\mathbf{M}_C$. Hence we get

Proposition 4.12. *The curve $\mathcal{B}$ has a rational point if and only if C can be hyperelliptically defined over $\mathbf{M}_C$.*

With the notation and hypotheses of Proposition 4.11, using $Q' = \mathcal{Q}$, we get a sufficient and necessary condition.

Corollary 4.13. *Let (q_1, q_2, q_3) be three covariants of order 2 for degree n binary forms and assume they are defined over the prime field of K. If $R(q_1, q_2, q_3)$ evaluated at a form f in non-zero, there exists a non-singular conic $\mathcal{Q}$ and a plane curve $\mathcal{H}$ of degree $n/2$ defined over $\mathbf{M}_C$ such that there is a K-isomorphism $\mathcal{Q} \to \mathbb{P}^1$ mapping the intersections points of $\mathcal{Q} \cap \mathcal{H}$ on the roots of $f(X, Z)$. In particular the field of moduli is a field of definition if and only if $\mathcal{Q}$ has a $\mathbf{M}_C$-rational point and in this case C can be hyperelliptically defined over $\mathbf{M}_C$.*

Note that the equivalence does not hold if C has extra-automorphism (see Remark 4.22).

If 'to be defined' and 'to be hyperelliptically defined' over $\mathbf{M}_C$ are the same problem when g is even, it is not the case for odd genus. Indeed,

Proposition 4.14. *Let g be odd. Then $\mathbf{M}_C$ is a field of definition for C.*

Proof. Consider the rational canonical divisor κ on $\mathcal{B}$ of degree -2. It is the negative of the intersection of a line with a place model of $\mathcal{B}$. If $\mathbf{M}_C$ is finite, we get the result directly from Section 4.1. Hence we assume that $\mathbf{M}_C$ is infinite. We can find a line such that $\mathrm{Supp}(-\kappa) \cap \mathrm{Supp}(D) = \emptyset$. Since $\deg D = 2g + 2$,

$D - 2\cdot\frac{g+1}{2}(-\kappa)$ is a divisor of degree 0, it is the divisor of a function $u \in \mathbf{M}_C(\tilde{Q})$. If we consider the degree 2 extension of the form $z^2 = u$, this defines a hyperelliptic curve over $\mathbf{M}_C$ with the same ramification as C, hence C admits a model over $\mathbf{M}_C$ (note that the crucial fact is that each point in the divisor $2\cdot\frac{g+1}{2}(-\kappa)$ has even multiplicity and hence does not contribute to the ramification). □

On the other hand, it is easy to exhibit examples for which $\mathbb{Q}$ is a field of moduli of a genus 3 curve C with no extra automorphism but C cannot be hyperelliptically defined over $\mathbb{Q}$. Indeed, consider the hyperelliptic genus 3 curve over $\mathbb{Q}$ which is a degree 2 cover of the conic $Q : x_1^2+x_2^2+x_3^2 = 0$ ramified over the intersection points of Q and

$$H : x_2^4 + x_1x_2^3 - (x_1^4 + x_1^3x_3 + x_1^2x_3^2 + x_1x_3^3 + 2x_3^4) = 0.$$

The curve C has a hyperelliptic equation

$$y^2 = -x^8 + (2i+2)x^7 - 8x^6 + (-2i-6)x^5 - 14x^4 + (-2i+6)x^3 - 8x^2 + (2i-2)x - 1$$

over $\mathbb{Q}(i)$ and we can check using our programs that C has no extra-automorphism. Since $Q \simeq \mathcal{B}$ and $Q(\mathbb{Q}) = \emptyset$, Proposition 4.12 shows that C cannot be hyperelliptically defined over $\mathbb{Q}$.

4.4. **The hyperelliptic case with extra-automorphisms.** When C has extra-automorphisms, the issues are less clear. In [Hug07] and [Hug05], the following results are proved.

Proposition 4.15 ([Hug07, Th.5.4], [Hug05, Prop.4.2.2])**.** *If $\overline{\mathrm{Aut}}(C)$ is not cyclic or is cyclic and of order divisible by p then C can be hyperelliptically defined over its field of moduli.*

Remark 4.16. In [FGD06] and [Fue10], one can find a family of hyperelliptic curves, among which for instance the genus 5 curve

$$C : y^2 = (x^4 - 2\,(1 - 2\,\frac{r_3-r_1}{r_3-r_2}\,\frac{q_4-r_2}{q_4-r_1})\,x^2 + 1)\cdot \\ (x^4 - 2\,(1 - 2\,\frac{r_3-r_1}{r_3-r_2}\,\frac{q_5-r_2}{q_5-r_1})\,x^2 + 1)\cdot(x^4 - 2\,(1 - 2\,\frac{r_3-r_1}{r_3-r_2}\,\frac{q_6-r_2}{q_6-r_1})\,x^2 + 1)\,,$$

with $q_4 = 3, q_5 = -1, q_6 = 7$ and r_1, r_2, r_3 the conjugate roots of $X^3 - 3X + 1 = 0$ (these values of q_i are only to fix ideas). This curve was claimed to be a curve with automorphism group $(\mathbb{Z}/2\mathbb{Z})^3$, field of moduli $\mathbb{Q}$ but with no hyperelliptic equation over $\mathbb{Q}$, contradicting Huggins' general result without the respective authors being aware of it. It took us a long month of discussions to realize that there was a subtle gap at page 406 point d) of [FGD06]: it is claimed there that a certain extension is Galois which is not always the case. It took us even longer to find an explicit hyperelliptic model of the curve over $\mathbb{Q}$ because, although the curve is defined over a cubic Galois extension of $\mathbb{Q}$, the $\bar{\mathbb{Q}}$-isomorphism we found between C and a model over $\mathbb{Q}$ is defined over a 12 degree extension of $\mathbb{Q}$. A faster and more systematic way to derive an hyperelliptic equation over the field of moduli is worked out in a work in progress by the authors of the present article.

We can add the following result which is implicitly contained in [Cou94].

Proposition 4.17. *If the signature of C has at least one odd exponent (with the convention of Table 3), then the field of moduli is a field of definition.*

Proof. Let s be one of the ramification index with odd signature exponent e. Consider the corresponding part Z of the ramification divisor W. Since Galois action respect the ramification index, the image of Z in the curve $\mathcal{B}$ (with the notation of Section 4.1) is a rational divisor of odd degree e. Hence by the remark following Theorem 4.4, we get the result. □

4.5. **The hyperelliptic case of genus** 3. Let now C be a genus 3 hyperelliptic curve. In this section, we want to address several issues which will be cut out according to the automorphism groups reviewed in Table 3.

Issue I: Can we compute the automorphism group from the invariants ?
Issue II: Is the field of moduli automatically a field of definition ?
Issue III: Can the curve be hyperelliptically defined over its field of moduli ?
Issue IV: Can we (hyperelliptically) reconstruct the curve from its invariants over K ?
Issue V: Can we reconstruct the curve over its field of moduli ?
Issue VI: Can we hyperelliptically reconstruct the curve over its field of moduli ?

Issue VII: What is the number of $\bar{F}_q$-isomorphism classes of hyperelliptic genus 3 curves with given automorphism group over a finite field $\mathbb{F}_q$?

The following table gather our state of knowledge (we **emphasize** what is proved in the present article). Note that the practical results are valid for $p = 0$ or $p \geq 11$.

#	Dim. 0	Dim. 1	$\mathbf{C}_2^3$	$\mathbf{C}_4$	$\mathbf{D}_4$	$\mathbf{C}_2$
I	yes	**yes**	**yes**	**yes**	**yes**	**yes**
II	yes	yes	yes	**yes**	no (counterex. exist)	**yes**
III	yes	yes	yes	**yes**	no	**computable** (possible theoretical obstruction)
IV	yes	**yes**	**yes**	**yes**	**yes**	**yes**
V	yes	**yes**	no **(over a cubic ext.)**	**yes**	no **(over a deg. 8 ext.)**	**yes**
VI	yes	**yes**	no **(over a cubic ext.)**	**yes**	no **(over a deg. 8 ext.)**	**yes** **(if no theoretical obstruction)**
VII	1	$\mathbf{q-3}$	$\mathbf{q^2-2q+2}$	$\mathbf{q^2-2q+2}$	q^3-2q^2+3 (unproven)	q^5-q^3+q-2 (unproven)

TABLE 7. Issues addressed in the present paper.

Remark 4.18. Similar issues for genus 2 are completely solved by [CQ05] (see also [CNP05, Th.5] for $p = 2$). In particular if the curve has extra-automorphism, then $\mathbf{M}_C$ is always a field of definition and moreover C can be hyperelliptically defined over $\mathbf{M}_C$. Actually, in [CQ05] for the case of automorphism group $\mathbf{D}_4$, the authors assumed that $p \neq 2, 3, 5$. A careful analysis and some improvements made during the implementation of their work in MAGMA by the two authors removed this restriction (see [LR08]).

STRATUM OF DIMENSION 0. This corresponds to cases 8,9,10 and 11 in Table 3. There is nothing to prove for them.

STRATUM OF DIMENSION 1. By Proposition 4.15, we have that cases 5,6 and 7 are hyperelliptically defined over their field of moduli. Moreover in Section 3.2.2 explicit equations for their stratum and an explicit hyperelliptic equation over the field of moduli are given.

STRATUM $\mathbf{C}_2^3$. By Proposition 4.15, we know that this case is hyperelliptically defined over its field of moduli. In Lemma 3.14, explicit equations for the stratum is given. Moreover an explicit hyperelliptic equation over an extension at most cubic of the field of moduli is given. To write a hyperelliptic equation over the field of moduli is a work in progress as mentioned in the introduction.

STRATUM $\mathbf{C}_4$. In Lemma 3.16 explicit equations for the stratum are given. Moreover since the signature is $(2^3, 4^2)$, the following result is a consequence of Proposition 4.17.

Proposition 4.19. *Let C/k be a hyperelliptic curve of genus 3 with $\mathrm{Aut}(C) \simeq \mathbf{C}_4$. Then C can be defined over its field of moduli.*

Remark 4.20. Proposition 4.19 has no analogue for even genus greater than 3, since there can be an obstruction as shown in [Hug05, Chap5].

Thanks to Lemma 3.16, the procedure is effective when $p \geq 11$ or $p = 0$. Indeed let $\{j_i\}$ be the Shioda invariants of such a curve C/k. In Lemma 3.16, it is proved that we can find three order 2 covariants q_i, such that $R(q_1, q_2, q_3)$ is non-zero. Now, we can use the first part of Proposition 4.11 to construct a conic $\mathcal{Q}$ and an quartic $\mathcal{H}$ over $\mathbf{M}_C$ such that there exists a K-isomorphism $\mathcal{Q} \to \mathbb{P}^1$ mapping the intersection divisor of $\mathcal{Q} \cap \mathcal{H}$ on the ramification divisor of C. Since $g = 3$ is odd, we can moreover proceed as in the proof of Proposition 4.14 and get a curve over $\mathbf{M}_C$, K-isomorphic to C. Hence C can be defined over its field of moduli. Moreover we can effectively construct a hyperelliptic equation over at most a quadratic extension of the field of moduli. Actually one can do better.

Proposition 4.21. *Let C/k be a hyperelliptic curve of genus 3 with $\mathrm{Aut}(C) \simeq \mathbf{C}_4$. Then C can be hyperelliptically defined over its field of moduli.*

Proof. By Proposition 4.19, we know that there is a model $\mathcal{C}$ of C over $\mathbf{M}_C$. Let $\mathcal{B} = \mathcal{C}/\langle\iota\rangle$ be seen as a plane non singular conic. The signature of C singles out on $\mathcal{B}$ a degree 2 effective rational divisor Z. If the points of Z are rational then we are done since then $\mathcal{B} \simeq \mathbb{P}^1$. Otherwise, after a quadratic extension F of $\mathbf{M}_C$, there is a map $\phi : \mathcal{B} \to \mathbb{P}^1$ sending one of this point to 0 and the other to ∞. Since these points were branches of $\mathcal{C}/\mathcal{B}$, there exists a K-isomorphism Φ lifting ϕ such that the curve $D = \Phi(\mathcal{C})$ is of the form $y^2 = xg(x)$ with $g \in F[x]$ of degree 6. Moreover since there is an involution which fixes 0 and ∞ and which is then $x \mapsto -x$, the polynomial g has only even power monomials. Let σ be the Galois involution of $F/\mathbf{M}_C$. Since D has a model over $\mathbf{M}_C$, there exists a K-isomorphism $(M_\sigma, e_\sigma) : D \to {}^\sigma D$ and M_σ is actually given by $\phi^\sigma \circ \phi^{-1}$ and hence is defined over F. Since the isomorphism preserves the ramification indexes, M_σ maps $\{0, \infty\}$ onto $\{0, \infty\}$. It is then easy to see that M_σ is either the map $x \mapsto ax$ or $x \mapsto a/x$ for some $a \in F$. By Lemma 4.6, since there exists $\lambda \in F^*$ such that $M_\sigma^\sigma M_\sigma = \lambda \cdot \mathrm{id}$, we get in the first case $a = \lambda = 1$ (and we are done) and $a = a^\sigma$ in the later case and so $a \in \mathbf{M}_C$.
If $b = \sqrt{a} \notin F$, let $F' = F(b)$ and σ' and τ be the two generators of the $\mathbf{C}_2 \times \mathbf{C}_2$ Galois extension $F'/\mathbf{M}_C$ defined by $\sigma'_{|F} = \sigma$, $b^{\sigma'} = b$ and $\tau_{|F} = \mathrm{id}$ and $b^\tau = -b$. We define

$$M'_{\sigma'} = \frac{1}{b} \cdot M_\sigma = \begin{bmatrix} 0 & b \\ \frac{1}{b} & 0 \end{bmatrix} \text{ and } M'_\tau = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

One can check that the conditions of 4.7 are satisfied.
If $b \in F$, then we define

$$M'_\sigma = \frac{1}{b} \cdot M_\sigma \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & b \\ -\frac{1}{b} & 0 \end{bmatrix}.$$

One can again check that the conditions of Lemma 4.7 are satisfied. □

The previous proof can be turned out into an effective method. Let assume that among the five determinants of Lemma 3.16, $R(C_{5,2}, C_{6,2}, C_{7,2})$ is not zero (the other cases are obtained by permutation). Then, due to the degree of the covariants and the action of $\mathbf{C}_4$, the conic and quartic have the following forms

$$\begin{aligned} \mathcal{Q} &: A_{11}x_1^2 + A_{13}x_1x_3 - A_{22}x_2^2 - A_{33}x_3^2 = 0 \\ \mathcal{H} &: x_2 \cdot (h_{11}x_1^3 + h_{13}x_1^2x_3 + h_{31}x_3^2x_1 + h_{33}x_3^3 + h_{12}x_1x_2^2 + h_{32}x_2^2x_3) = 0. \end{aligned}$$

By a linear change of variables in x_1 and x_3 we can assume that $A_{13} = 0$ and $A_{11} = 1$ and we write

$$\mathcal{Q} : x_1^2 - ax_2^2 + cx_3^2 = 0.$$

The conic and the quartic have the involution $(x_1 : x_2 : x_3) \mapsto (x_1 : -x_2 : x_3)$ which hence stabilizes the ramification divisor of C and is the involution of $\overline{\mathrm{Aut}}(C)$. The divisor $\{x_2 = 0\} \cap \mathcal{Q}$ is the fixed divisor Z of the proof. The points of Z are $(\pm\alpha : 0 : 1)$ where α is a root of $X^2 = c$. The isomorphism $\phi^{-1} : \mathbb{P}^1 \to \mathcal{Q}$ is given by

$$(t : u) \mapsto (\alpha(at^2 + u^2) : 2\alpha tu : u^2 - at^2).$$

From this, we see that

$$M_\sigma = \phi^\sigma \circ \phi^{-1} = \begin{bmatrix} 0 & a \\ 1 & 0 \end{bmatrix}.$$

We can then apply the formula in the proof of Lemma 4.7 to get the result. As alternative, we have noticed that there is another choice of parametrization which only requires a quadratic extension and that we implemented. We normalize

$$\mathcal{Q} : ax_1^2 - x_2^2 + cx_3^2$$

and parametrize the conic $\mathcal{Q}$ with respect to the point $(0 : \sqrt{c} : 1)$ by

$$(t : u) \mapsto (2\sqrt{c}tu : \sqrt{c}(at^2 + u^2) : at^2 - u^2).$$

Plugging these expressions in $\mathcal{H}$ leads to a polynomial $g \in \mathbf{M}_C[x]$ times $\sqrt{c}$, which is what we need.

Remark 4.22. Note that contrary to the case with no extra-automorphism, it might happen that none of the conics $\mathcal{Q}$ contains arithmetic information, *i.e.* that all of them have no points over $\mathbf{M}_C$. In this way, for instance, we noticed that none of the non-degenerate conics among the 19 conics of Lemma 3.24 have a rational point for $(j_2 : j_3 : \ldots : j_9)$ equal to $(1 : 0 : 6 : 0 : 6 : 0 : 2963/2835 : 0 : 2963/2835)$ over $\mathbb{Q}$.

STRATUM $\mathbf{D}_4$. In [Hug05, Chap.5], Huggins constructs examples of curves

$$C : y^2 = (x^2 - a_1)(x^2 + \frac{1}{\bar{a_1}})(x^2 - a_2)(x^2 + \frac{1}{\bar{a_2}})$$

with $a_1, a_2 \in \mathbb{Q}(i) \setminus (\mathbb{Q} \cup i\mathbb{Q})$, $|a_i| > 1$ and $|a_1/a_2| \neq 1$, which are defined over $\mathbb{Q}(i)$ with geometric group of automorphism $(\mathbb{Z}/2\mathbb{Z})^2$. By [Hug07, Prop.5.0.5], C has field of moduli $\mathbb{Q}(i) \cap \mathbb{R} = \mathbb{Q}$ but cannot be defined over $\mathbb{Q}$. Hence the field of moduli is not always a field of definition and there is not always a hyperelliptic equation over the field of moduli. However Lemma 3.20 gives explicit equations for the stratum and shows how to construct a hyperelliptic equation over an extension of the field of moduli of degree at most 8.

Remark 4.23. In [GSS05, Cor.2], it is asserted that as soon as $\#\operatorname{Aut}(C) > 2$, the field of moduli of C is a field of definition. However, the 'moduli problem' is distinct from the usual one, which explains the contradiction with Huggins' result.

STRATUM $\mathbf{C}_2$. This is the generic case for which we can apply the results of Section 4.3. Note that as we can find a non singular conic $\mathcal{Q}$ in this case, we can compute the obstruction using Corollary 4.13. When the obstruction is trivial, we apply Mestre's method to obtain a model with a hyperelliptic equation over the field of moduli.

4.6. **The case of curves over finite fields.**

Proposition 4.24. *Let C be a (hyperelliptic) curve of genus $g \geq 2$ defined over a finite field $k = \mathbb{F}_q$, such that all the $\bar{k}$-automorphisms of C are defined over k. Let k' be the extension of k of degree equal to the exponent e of $\operatorname{Aut}(C)$. Then there exists a k'-isomorphism from C to a curve defined over $\mathbf{M}_C$.*

Proof. Let $\mathcal{C}$ be a model of C over $\mathbf{M}_C$ and $\Phi : C \to \mathcal{C}$. We want to show that for the Frobenius automorphism $(x \mapsto x^{q^e}) = \tau \in \operatorname{Gal}(\bar{k'}/k')$ we have $\Phi^\tau = \Phi$. If $\gamma = x \mapsto x^q$, we have that $(\Phi^\gamma)^{-1} \circ \Phi = f \in \operatorname{Aut}(C)$. Since γ acts trivially on the automorphism group, we get

$$(\Phi^\tau)^{-1} \circ \Phi = ((\Phi^{\gamma^e})^{-1} \circ \Phi^{\gamma^{e-1}}) \cdots \circ ((\Phi^\gamma)^{-1} \circ \Phi^\gamma) = f \circ \cdots \circ f = 1.$$

□

Assume one knows how to compute a k'-isomorphism $F_\sigma : C \to {}^\sigma C$ for σ a generator of $\operatorname{Gal}(k'/\mathbf{M}_C)$ and that we know the automorphism of C explicitly. If C is given by a hyperelliptic equation, we can represent all these morphisms by couple (M_σ, e_σ) and we will actually only care about the matrix M_σ. Two different F_σ differ by an automorphism of C, hence we have a finite explicit set $\mathcal{F}$ of k'-isomorphisms represented by M_σ.

Now pick one of the M_σ in $\mathcal{F}$. Let $m = \deg(k'/\mathbf{M}_C)$ If

$$M_\sigma^{\sigma^{m-1}} \cdots M_\sigma^\sigma \cdot M_\sigma \neq \lambda \cdot \mathrm{id}$$

for $\lambda \in \mathbf{M}_C$ then pick another $M_\sigma \in \mathcal{F}$. Since there exists a k'-isomorphism from C to a model over $\mathbf{M}_C$, we know that there is at least one solution to the previous equation and the procedure produces a good M_σ and λ after a finite number of trials (less than $\#\overline{\operatorname{Aut}}(C)$). We let $M'_\sigma = \frac{1}{a} \cdot M_\sigma$ for a a solution of $\operatorname{Norm}_{k'/\mathbf{M}_C}(a) = \lambda$ (which always exists in a finite field). We can then apply the formula of Lemma 4.7

$$A = P + \sum_{i=1}^{m-1} P^{\sigma^i} \cdot M^{\sigma^i} \cdots M^\sigma \cdot M$$

for a random matrix $P \in M_2(k')$. We repeat this last part until we find A invertible. We can then apply A to the hyperelliptic polynomial of C to get a hyperelliptic equation defined over $\mathbf{M}_C$.

Practically, in the hyperelliptic case, M_σ can be computed thanks for instance to the function `IsGL2-Equivalent` of Magma. However, when C has extra-automorphisms, a better strategy is available. Since the automorphisms of C are defined over k', we can assume that $\overline{\mathrm{Aut}}(C)$ is given by one of the canonical expression of [Hug05, Lem.2.2.1]. Moreover, if $p \nmid \#\,\mathrm{Aut}(C)$ then by [Hug05, Lem.3.1.2], we have that M_σ belongs to the normalizer $\mathcal{N}$ of $\overline{\mathrm{Aut}}(C)$ in $\mathrm{PGL}_2(k')$. Moreover when $\overline{\mathrm{Aut}}(C)$ is not cyclic, then $\mathcal{N}$ is finite so we can easily find the isomorphism.

Remark 4.25. If C is non hyperelliptic, we can assume that it is canonically embedded. Then, isomorphisms and automorphisms are given by $(g+1)\times(g+1)$ matrices and the same strategy can be applied.

As an application, we use the algorithms developed in this article to exhibit a hyperelliptic equation over $\mathbb{F}_p$, $p = 11, 13, \ldots, 47$, for all the $\bar{\mathbb{F}}_p$-isomorphism classes of hyperelliptic curves of genus 3. As expected (*cf.* Table 7), we have obtained

- one model for each stratum of dimension 0,
- $p-3$ models for each stratum of dimension 1,
- $p^2-2\,p+2$ models for each stratum of dimension 2,
- $p^3-2\,p^2+3$ models for the stratum $\mathbf{D}_4$,
- p^5-p^3+p-2 models for the stratum $\mathbf{C}_2$,

that is a total number of p^5 non-$\bar{\mathbb{F}}_p$-isomorphic curves as predicted in [BG01].

5. Conclusion

Although a big part of the tasks we addressed in the introduction is now completed, we wish to pursue our researches in two directions. The first one is geometric and would be to give a conceptual proof of the existence of a non singular conic $\mathcal{Q}$ when the curve has no extra-automorphism. We could then hope to prove such a statement without restricting to the genus 3 case. The second one is arithmetic and is related to the explicit hyperelliptic descent over the field of moduli in the case $\mathbf{C}_2^3$. Here also, as we saw for instance with the genus 5 example of Fuertes and González-Diez, the problem is not limited to genus 3 curves, neither is it to this particular automorphism group nor to hyperelliptic curves. We will use covariants in a forthcoming article to get both theoretical and practical results on this issue in a general context.

References

[Bed07] Leonid Bedratyuk. On complete system of invariants for the binary form of degree 7. *Journal of Symbolic Computation*, 42:935, 2007.

[BG01] Bradley W. Brock and Andrew Granville. More points than expected on curves over finite field extensions. *Finite Fields Appl.*, 7(1):70–91, 2001. Dedicated to Professor Chao Ko on the occasion of his 90th birthday.

[BK86] F. A. Bogomolov and P. I. Katsylo. Rationality of some quotient varieties. *Mathematics of the USSR-Sbornik*, 54:571–576, 1986.

[Bol87] Oskar Bolza. On Binary Sextics with Linear Transformations into Themselves. *Amer. J. Math.*, 10(1):47–70, 1887.

[BP10a] A.E. Brouwer and M. Popoviciu. The invariants of the binary decimic, 2010. available on http://arxiv.org/abs/1002.1008.

[BP10b] A.E. Brouwer and M. Popoviciu. The invariants of the binary nonic, 2010. available on http://arxiv.org/abs/1002.0761.

[Bra88] Rolf Brandt. *Über die Automorphismengruppen von algebraischen Funktionenkörpern*. PhD thesis, Universität Essen, 1988.

[BS86] Rolf Brandt and Henning Stichtenoth. Die Automorphismengruppen hyperelliptischer Kurven. *Manuscripta Math.*, 55(1):83–92, 1986.

[BV05] H. Babu and P. Venkataraman. Group action on genus 3 curves and their Weierstrass points. In *Computational aspects of algebraic curves*, volume 13 of *Lecture Notes Ser. Comput.*, pages 264–272. World Sci. Publ., Hackensack, NJ, 2005.

[Car03] Gabriel Cardona. On the number of curves of genus 2 over a finite field. *Finite Fields Appl.*, 9(4):505–526, 2003.

[CFA+06] Henri Cohen, Gerhard Frey, Roberto Avanzi, Christophe Doche, Tanja Lange, Kim Nguyen, and Frederik Vercauteren, editors. *Handbook of elliptic and hyperelliptic curve cryptography*. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2006.

[Cle72] A Clebsch. *Theorie der binären algebraischen formen*. Verlag von B.G. Teubner, Leipzig, 1872.

[CNP05] G. Cardona, E. Nart, and J. Pujolàs. Curves of genus two over fields of even characteristic. *Math. Zeitschrift*, 250:177–201, 2005.

[Cou94] Jean-Marc Couveignes. Calcul et rationalité de fonctions de Belyĭ en genre 0. *Ann. Inst. Fourier (Grenoble)*, 44(1):1–38, 1994.

[CQ05] G. Cardona and J. Quer. Field of moduli and field of definition for curves of genus 2. In *Computational aspects of aglebraic curves*, volume 13 of *Lecture Notes Ser. Comput.*, pages 71–83, Hackensack, NJ„ 2005. World Sci. Publ.

[Cro02] H. Croeni. *Zur Berechnung von Kovarianten von Quantiken*. PhD thesis, Univ. des Saarlandes, Saarbrücken, 2002.

[DE99] Pierre Dèbes and Michel Emsalem. On fields of moduli of curves. *J. Algebra*, 211(1):42–56, 1999.

[Dix90] J. Dixmier. Quelques aspects de la théorie des invariants. *Gaz. Math., Soc. Math. Fr.*, 43:39–64, 1990.

[DL86] J. Dixmier and D. Lazard. Le nombre minimum d'invariants fondamentaux pour les formes binaires de degré 7. *Portugal. Math.*, 43(3):377–392, 1985/86.

[FGD06] Y. Fuertes and G. González-Diez. Fields of moduli and definition of hyperelliptic covers. *Arch. Math. (Basel)*, 86(5):398–408, 2006.

[Fue10] Yolanda Fuertes. Fields of moduli and definition of hyperelliptic curves of odd genus. *Arch. Math. (Basel)*, 95(1):15–81, 2010.

[Gor68] Paul Gordan. Beweis, dass jede Covariante und Invariante einer binären Form eine ganze Function mit numerischen Coefficienten einer endlichen Anzahl solcher Formen ist. *Journal für die reine und angewandte Mathematik*, (69):323–354, 1868.

[Gro71] A. Grothendieck. *Revêtements étales et géométrie algébrique (SGA 1)*, volume 224 of *Lecture Notes in Math.* Springer-Verlag, Heidelberg, 1971.

[GSS05] Jaime Gutierrez, D. Sevilla, and T. Shaska. Hyperelliptic curves of genus 3 with prescribed automorphism group. In *Computational aspects of algebraic curves*, volume 13 of *Lecture Notes Ser. Comput.*, pages 109–123. World Sci. Publ., Hackensack, NJ, 2005.

[GY03] J.H. Grace and A. Young. *The algebra of invariants*. Chelsea publishing company, New-York, 1903.

[Har77] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.

[HH03] H. Hammer and F. Herrlich. A remark on the moduli field of a curve. *Arch. Math. (Basel)*, 81(1):5–10, 2003.

[Hug05] B. Huggins. *Fields of moduli and fields of definition of curves*. PhD thesis, University of California, Berkeley, Berkeley, California, 2005. http://arxiv.org/abs/math.NT/0610247.

[Hug07] B. Huggins. Fields of moduli of hyperelliptic curves. *Math. Res. Lett.*, 14(2):249–262, 2007.

[Koi72] Shoji Koizumi. The fields of moduli for polarized abelian varieties and for curves. *Nagoya Math. J.*, 48:37–55, 1972.

[LR08] R. Lercier and C. Ritzenthaler. Invariants and reconstructions for genus 2 curves in any characteristic, 2008. Available in MAGMA 2.15 and later, http://magma.maths.usyd.edu.au/magma/handbook/text/1367.

[LRZ10] Gilles Lachaud, Christophe Ritzenthaler, and Alexey Zykin. Jacobians among abelian threefolds: a formula of Klein and a question of Serre. *Math. Res. Lett.*, 17(2), 2010.

[Mae90] Takashi Maeda. On the invariant field of binary octavics. *Hiroshima Mathematical Journal*, 20(3):619–632, 1990.

[Mat58] Teruhisa Matsusaka. On a theorem of Torelli. *Amer. J. Math.*, 80:784–800, 1958.

[Mes91] Jean-François Mestre. Construction de courbes de genre 2 à partir de leurs modules. In *Effective methods in algebraic geometry*, volume 94 of *Prog. Math.*, pages 313–334, Boston, 1991. Birkäuser.

[MF82] David Mumford and John Fogarty. *Geometric invariant theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*. Springer-Verlag, Berlin, second edition, 1982.

[Mil72] J. S. Milne. Abelian varieties defined over their fields of moduli. I. *Bull. London Math. Soc.*, 4:370–372, 1972.

[Mil74] J. S. Milne. Correction: "Abelian varieties defined over their fields of moduli. I" (Bull. London Math. Soc. **4** (1972), 370–372). *Bull. London Math. Soc.*, 6:145–146, 1974.

[MSSV02] K. Magaard, T. Shaska, S. Shpectorov, and H. Völklein. The locus of curves with prescribed automorphism group. *Sūrikaisekikenkyūsho Kōkyūroku*, (1267):112–141, 2002. Communications in arithmetic fundamental groups (Kyoto, 1999/2001).

[NS04] E. Nart and D. Sadornil. Hyperelliptic curves of genus three over finite fields of characteristic two. *Finite Fields and Their Applications*, 10:198–220, 2004.

[Pro07] Claudio Procesi. *Lie groups*. Universitext. Springer, New York, 2007. An approach through invariants and representations.

[Rie93] John F. X. Ries. Subvarieties of moduli space determined by finite groups acting on surfaces. *Trans. Amer. Math. Soc.*, 335(1):385–406, 1993.

[Roq70] P. Roquette. Abschätzung der Automorphismenanzahl von Funktionenkörpern. *Math. Z.*, 117:157–163, 1970.

[Sek85] Tsutomu Sekiguchi. Wild ramification of moduli spaces for curves or for abelian varieties. *Compositio Math.*, 54:33–372, 1985.

[Ser68] Jean-Pierre Serre. *Corps locaux*. Hermann, Paris, 1968. Deuxième édition, Publications de l'Université de Nancago, No. VIII.

[Shi67] T. Shioda. On the graded ring of invariants of binary octavics. *American J. of Math.*, 89(4):1022–1046, 1967.

[Sin72] David Singerman. Finitely maximal Fuchsian groups. *J. London Math. Soc. (2)*, 6:29–38, 1972.

[VG88] F. Von Gall. Das vollständige Formensystem der binären Form 7ter Ordnung. *Math. Ann.*, 31:318–336, 1888.

[Web99] Heinrich Weber. *Lehrbuch der Algebra, second ed., vol II*. Vieweg, Braunshweig, 1899.

[Wei56] A. Weil. The field of definition of a variety. *American Journal of Mathematics*, 78:509–524, 1956.

Appendix A. Stratum equations for the automorphism group $\mathbf{D}_4$

$$
\begin{aligned}
0 = {} & 49\,j_2^4\,j_4^2 - 1470\,j_2\,j_3^2\,j_4^2 - 2205\,j_2^2\,j_4^3 - 17496\,j_4^4 + 19845\,j_3\,j_4^2\,j_5 - 1092\,j_2^3\,j_4\,j_6 + 32760\,j_3^2\,j_4\,j_6 + 47880\,j_2\,j_4^2\,j_6 \\
& - 238140\,j_5^2\,j_6 + 133056\,j_4\,j_6^2 + 357210\,j_4\,j_5\,j_7 - 238140\,j_3\,j_6\,j_7 - 59535\,j_2\,j_7^2 - 884520\,j_4^2\,j_8 + 238140\,j_2\,j_6\,j_8 + 1786050\,j_8^2 \\
& + 119070\,j_3\,j_4\,j_9 + 59535\,j_2\,j_5\,j_9 - 714420\,j_7\,j_9 + 4410\,j_2^2\,j_{10} - 132300\,j_2^2\,j_{10} - 317520\,j_2\,j_4\,j_{10} + 2540160\,j_6\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -14\,j_2^3\,j_3\,j_4^2 + 420\,j_3^3\,j_4^2 + 630\,j_2\,j_3\,j_4^3 - 189\,j_2^2\,j_4^2\,j_5 + 360\,j_3\,j_4^2\,j_6 - 168\,j_2^3\,j_5\,j_6 + 5040\,j_3^2\,j_5\,j_6 + 3348\,j_2\,j_4\,j_5\,j_6 + 12096\,j_5\,j_6^2 \\
& + 252\,j_2^3\,j_4\,j_7 - 7560\,j_3^2\,j_4\,j_7 - 2916\,j_2\,j_4^2\,j_7 + 2268\,j_2^2\,j_6\,j_7 - 22032\,j_4\,j_6\,j_7 + 34020\,j_3\,j_7^2 - 68040\,j_3\,j_6\,j_8 - 34020\,j_2\,j_7\,j_8 - 42\,j_2^4\,j_9 \\
& + 1260\,j_2\,j_3^2\,j_9 + 756\,j_2^2\,j_4\,j_9 + 7776\,j_4^2\,j_9 - 34020\,j_3\,j_5\,j_9 - 24192\,j_2\,j_6\,j_9 + 408240\,j_8\,j_9 + 34020\,j_3\,j_4\,j_{10} + 17010\,j_2\,j_5\,j_{10} - 204120\,j_7\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 14\,j_2^7\,j_4 - 840\,j_2^4\,j_3^2\,j_4 + 12600\,j_2\,j_3^4\,j_4 - 1260\,j_2^5\,j_4^2 + 37800\,j_2^2\,j_3^2\,j_4^2 + 25191\,j_2^3\,j_3^3 + 94770\,j_3^2\,j_4^3 - 743580\,j_2\,j_4^4 + 11340\,j_2^3\,j_3\,j_4\,j_5 - 340200\,j_3^3\,j_4\,j_5 \\
& - 510300\,j_2\,j_3\,j_4^2\,j_5 + 76545\,j_2^2\,j_4\,j_5^2 + 1771470\,j_4^2\,j_5^2 + 546\,j_2^2\,j_6 - 32760\,j_2^3\,j_6 + 491400\,j_3^2\,j_6 - 58446\,j_2^2\,j_4\,j_6 + 1753380\,j_2\,j_3^2\,j_4\,j_6 \\
& + 1524420\,j_2^2\,j_4^2\,j_6 - 2449440\,j_3^2\,j_6 - 3768930\,j_3\,j_4\,j_5\,j_6 - 2985255\,j_2\,j_5^2\,j_6 + 47250\,j_2^3\,j_7^2 - 1417500\,j_2^2\,j_6^2 + 3418200\,j_2\,j_4\,j_6^2 + 13608000\,j_6^3 \\
& + 8726130\,j_3\,j_4^2\,j_7 + 51030\,j_2^3\,j_5\,j_7 - 1530900\,j_3^2\,j_5\,j_7 + 688905\,j_2\,j_4\,j_5\,j_7 - 22963500\,j_5\,j_6\,j_7 + 221130\,j_2^2\,j_4\,j_8 - 6633900\,j_3^2\,j_4\,j_8 \\
& - 18676980\,j_2\,j_4^2\,j_8 + 20667150\,j_5^2\,j_8 + 61236000\,j_4\,j_6\,j_8 - 34020\,j_2^2\,j_3\,j_9 + 1020600\,j_3^2\,j_9 + 1530900\,j_2\,j_3\,j_4\,j_9 - 459270\,j_2^2\,j_5\,j_9 + 11022480\,j_4\,j_5\,j_9 \\
& - 30618000\,j_3\,j_6\,j_9 + 34020\,j_2^4\,j_{10} - 1020600\,j_2\,j_3^2\,j_{10} - 1530900\,j_2^2\,j_4\,j_{10} - 44089920\,j_4^2\,j_{10} + 13778100\,j_3\,j_5\,j_{10} + 30618000\,j_2\,j_6\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -1078\,j_2^7\,j_4 + 64680\,j_2^4\,j_3^2\,j_4 - 970200\,j_2\,j_3^4\,j_4 + 83790\,j_2^5\,j_4^2 - 2513700\,j_2^2\,j_3^2\,j_4^2 - 417312\,j_2^3\,j_3^3 - 35108640\,j_2^2\,j_3^4 - 3061800\,j_2\,j_4^4 \\
& - 873180\,j_2^3\,j_3\,j_4\,j_5 + 26195400\,j_3^3\,j_4\,j_5 + 33934950\,j_2\,j_3\,j_4^2\,j_5 - 5893965\,j_2^2\,j_4\,j_5^2 + 8266860\,j_4^2\,j_5^2 - 18522\,j_2^2\,j_6 + 1111320\,j_2^2\,j_3^2\,j_6 \\
& - 16669800\,j_3^4\,j_6 + 1907262\,j_2^2\,j_4\,j_6 - 57217860\,j_2\,j_3^2\,j_4\,j_6 - 47979540\,j_2^2\,j_4^2\,j_6 - 30093120\,j_3^4\,j_6 - 22095990\,j_3\,j_4\,j_5\,j_6 \\
& + 165566835\,j_2\,j_5^2\,j_6 + 476280\,j_2^3\,j_6^2 - 14288400\,j_3^2\,j_6^2 - 324505440\,j_2\,j_4\,j_6^2 - 47304810\,j_3\,j_4^2\,j_7 - 3929310\,j_2^3\,j_5\,j_7 + 117879300\,j_3^2\,j_5\,j_7 \\
& + 43401015\,j_2\,j_4\,j_5\,j_7 + 64297800\,j_2\,j_3\,j_6\,j_7 + 32148900\,j_5\,j_6\,j_7 + 32148900\,j_2^2\,j_7^2 - 289340100\,j_4\,j_7^2 - 2738610\,j_2^2\,j_4\,j_8 + 82158300\,j_3^2\,j_4\,j_8 \\
& + 409362660\,j_2\,j_4^2\,j_8 - 1591370550\,j_5^2\,j_8 - 64297800\,j_2^2\,j_6\,j_8 + 3735396000\,j_4\,j_6\,j_8 - 964467000\,j_3\,j_7\,j_8 + 1428840\,j_2^3\,j_3\,j_9 \\
& - 42865200\,j_3^2\,j_9 - 96446700\,j_2\,j_3\,j_4\,j_9 + 3214890\,j_2^2\,j_5\,j_9 + 308629440\,j_4\,j_5\,j_9 + 514382400\,j_3\,j_6\,j_9 - 192893400\,j_2\,j_7\,j_9 - 2619540\,j_2^4\,j_{10} \\
& + 78586200\,j_2\,j_3^2\,j_{10} + 150028200\,j_2^2\,j_4\,j_{10} - 1014068160\,j_4^2\,j_{10} - 578680200\,j_3\,j_5\,j_{10} - 1200225600\,j_2\,j_6\,j_{10} + 11573604000\,j_8\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 14\,j_2^7\,j_4 - 840\,j_2^4\,j_3^2\,j_4 + 12600\,j_2\,j_3^4\,j_4 - 1050\,j_2^5\,j_4^2 + 31500\,j_2^2\,j_3^2\,j_4^2 - 5994\,j_2^3\,j_3^3 + 746820\,j_2^2\,j_4^3 - 126360\,j_2\,j_4^4 + 11340\,j_2^3\,j_3\,j_4\,j_5 \\
& - 340200\,j_3^3\,j_4\,j_5 - 425250\,j_2\,j_3\,j_4^2\,j_5 + 76545\,j_2^2\,j_4\,j_5^2 + 4067820\,j_4^2\,j_5^2 + 546\,j_2^2\,j_6 - 32760\,j_2^3\,j_6 + 491400\,j_3^2\,j_6 - 55926\,j_2^2\,j_4\,j_6 \\
& + 1677780\,j_2\,j_3^2\,j_4\,j_6 + 1405620\,j_2^2\,j_4^2\,j_6 + 77760\,j_4^3\,j_6 - 852930\,j_3\,j_4\,j_5\,j_6 - 4005855\,j_2\,j_5^2\,j_6 + 117180\,j_2^2\,j_6^2 - 3515400\,j_3^2\,j_6^2 + 2371680\,j_2\,j_4\,j_6^2 \\
& + 2894130\,j_3\,j_4^2\,j_7 + 51030\,j_2^3\,j_5\,j_7 - 1530900\,j_3^2\,j_5\,j_7 + 2219805\,j_2\,j_4\,j_5\,j_7 - 1020600\,j_2\,j_3\,j_6\,j_7 - 1530900\,j_5\,j_6\,j_7 - 510300\,j_2\,j_7^2 - 4592700\,j_4\,j_7^2 \\
& + 221130\,j_2^2\,j_4\,j_8 - 6633900\,j_3^2\,j_4\,j_8 - 16635780\,j_2\,j_4^2\,j_8 + 20667150\,j_5^2\,j_8 + 1020600\,j_2^2\,j_6\,j_8 + 12247200\,j_4\,j_6\,j_8 + 15309000\,j_3\,j_7\,j_8 \\
& - 15120\,j_2^3\,j_3\,j_9 + 453600\,j_3^3\,j_9 + 1190700\,j_2\,j_3\,j_4\,j_9 + 51030\,j_2^2\,j_5\,j_9 - 44089920\,j_4\,j_5\,j_9 - 13608000\,j_3\,j_6\,j_9 - 3061800\,j_2\,j_7\,j_9 \\
& + 146966400\,j_9^2 + 34020\,j_2^4\,j_{10} - 1020600\,j_2\,j_3^2\,j_{10} - 2041200\,j_2^2\,j_4\,j_{10} - 19595520\,j_4^2\,j_{10} + 6123600\,j_3\,j_5\,j_{10} + 24494400\,j_2\,j_6\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -14\,j_2^6\,j_3\,j_4 + 840\,j_2^3\,j_3^3\,j_4 - 12600\,j_3^5\,j_4 + 1260\,j_2^4\,j_3\,j_4^2 - 37800\,j_2\,j_3^3\,j_4^2 - 28350\,j_2^2\,j_3\,j_4^3 + 1180980\,j_3\,j_4^4 - 378\,j_2^5\,j_4\,j_5 + 11340\,j_2^2\,j_3^2\,j_4\,j_5 \\
& + 41229\,j_2^3\,j_4^2\,j_5 - 726570\,j_2^2\,j_3^2\,j_5 - 1061424\,j_2\,j_4^3\,j_5 - 76545\,j_2\,j_3\,j_4\,j_5^2 + 2066715\,j_4\,j_5^3 + 16866\,j_2^3\,j_3\,j_4\,j_6 - 505980\,j_3^3\,j_4\,j_6 \\
& - 758970\,j_2\,j_3\,j_4^2\,j_6 + 22302\,j_2^2\,j_5\,j_6 - 669060\,j_2\,j_3^2\,j_5\,j_6 - 775899\,j_2^2\,j_4\,j_5\,j_6 - 2758536\,j_4^2\,j_5\,j_6 + 9032310\,j_3\,j_5^2\,j_6 - 7510320\,j_3\,j_4\,j_6^2 \\
& + 1701000\,j_2\,j_5\,j_6^2 + 126\,j_2^6\,j_7 - 7560\,j_2^3\,j_4^2\,j_7 + 113400\,j_3^4\,j_7 - 22491\,j_2^2\,j_4\,j_7 + 674730\,j_2\,j_3^2\,j_4\,j_7 + 261954\,j_2^2\,j_4^2\,j_7 + 629856\,j_4^3\,j_7 \\
& - 10563210\,j_3\,j_4\,j_5\,j_7 - 192780\,j_2^3\,j_6\,j_7 + 5783400\,j_3^2\,j_6\,j_7 + 3363120\,j_2\,j_4\,j_6\,j_7 + 6889050\,j_5\,j_7^2 + 14849730\,j_3\,j_4^2\,j_8 + 51030\,j_2^3\,j_5\,j_8 \\
& - 1530900\,j_3^2\,j_5\,j_8 + 3750705\,j_2\,j_4\,j_5\,j_8 - 41334300\,j_5\,j_6\,j_8 - 18370800\,j_4\,j_7\,j_8 + 1134\,j_2^5\,j_9 - 34020\,j_2^2\,j_3^2\,j_9 - 81648\,j_2^3\,j_4\,j_9 \\
& + 918540\,j_2^2\,j_4\,j_9 + 4531464\,j_2\,j_4^2\,j_9 + 459270\,j_2\,j_3\,j_5\,j_9 - 19289340\,j_2^2\,j_5\,j_9 + 1632960\,j_2^2\,j_6\,j_9 + 14696640\,j_4\,j_6\,j_9 + 9185400\,j_3\,j_7\,j_9 \\
& - 34020\,j_2^3\,j_3\,j_{10} + 1020600\,j_3^2\,j_{10} + 1530900\,j_2\,j_3\,j_4\,j_{10} - 459270\,j_2^2\,j_5\,j_{10} + 29393280\,j_4\,j_5\,j_{10} - 48988800\,j_3\,j_6\,j_{10} - 9185400\,j_2\,j_7\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -28\,j_2^6\,j_3\,j_4 + 1680\,j_2^3\,j_3^3\,j_4 - 25200\,j_3^5\,j_4 + 2520\,j_2^4\,j_3\,j_4^2 - 75600\,j_2\,j_3^3\,j_4^2 - 56700\,j_2^2\,j_3\,j_4^3 + 1771470\,j_3\,j_4^4 - 756\,j_2^5\,j_4\,j_5 \\
& + 22680\,j_2^2\,j_3^2\,j_4\,j_5 + 76788\,j_2^3\,j_4^2\,j_5 - 1283040\,j_2^2\,j_4^3\,j_5 - 2009853\,j_2\,j_4^4\,j_5 - 153090\,j_2\,j_3\,j_4\,j_5^2 + 4133430\,j_4\,j_5^3 + 18612\,j_2^3\,j_3\,j_4\,j_6 \\
& - 558360\,j_3^3\,j_4\,j_6 - 837540\,j_2\,j_3\,j_4^2\,j_6 + 29484\,j_2^2\,j_5\,j_6 - 884520\,j_2\,j_3^2\,j_5\,j_6 - 1075518\,j_2^2\,j_4\,j_5\,j_6 - 6333552\,j_4^2\,j_5\,j_6 + 11941020\,j_3\,j_5^2\,j_6 \\
& - 11088900\,j_3\,j_4\,j_6^2 + 1275750\,j_2\,j_5\,j_6^2 + 252\,j_2^5\,j_7 - 15120\,j_2^2\,j_3^2\,j_7 + 226800\,j_3^4\,j_7 - 37422\,j_2^3\,j_4\,j_7 + 1122660\,j_2\,j_3^2\,j_4\,j_7 \\
& + 591948\,j_2^2\,j_4^2\,j_7 + 1653372\,j_4^3\,j_7 - 11941020\,j_3\,j_4\,j_5\,j_7 - 226800\,j_2^3\,j_6\,j_7 + 6804000\,j_3^2\,j_6\,j_7 + 6123600\,j_2\,j_4\,j_6\,j_7 + 5103000\,j_6^2\,j_7 \\
& + 17452260\,j_3\,j_4^2\,j_8 + 102060\,j_2^2\,j_5\,j_8 - 3061800\,j_3^2\,j_5\,j_8 + 1377810\,j_2\,j_4\,j_5\,j_8 - 45927000\,j_5\,j_6\,j_8 + 2268\,j_2^2\,j_9 - 68040\,j_2^2\,j_3^2\,j_9 \\
& - 197316\,j_3^2\,j_4\,j_9 + 2857680\,j_2^2\,j_4\,j_9 + 7225848\,j_2\,j_4^2\,j_9 + 918540\,j_2\,j_3\,j_5\,j_9 - 24800580\,j_5^2\,j_9 + 2041200\,j_2^2\,j_6\,j_9 + 24494400\,j_4\,j_6\,j_9 \\
& - 68040\,j_2^2\,j_3\,j_{10} + 2041200\,j_3^2\,j_{10} + 3061800\,j_2\,j_3\,j_4\,j_{10} - 918540\,j_2^2\,j_5\,j_{10} + 22044960\,j_4\,j_5\,j_{10} - 61236000\,j_3\,j_6\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -56\,j_2^6\,j_3\,j_4 + 3360\,j_2^3\,j_3^3\,j_4 - 50400\,j_3^5\,j_4 + 5670\,j_2^4\,j_3\,j_4^2 - 170100\,j_2\,j_3^3\,j_4^2 - 141750\,j_2^2\,j_3\,j_4^3 + 1574640\,j_3\,j_4^4 - 1512\,j_2^5\,j_4\,j_5 \\
& + 45360\,j_2^2\,j_3^2\,j_4\,j_5 + 128061\,j_2^3\,j_4^2\,j_5 - 1545480\,j_2^2\,j_4^3\,j_5 - 2204496\,j_2\,j_4^4\,j_5 - 306180\,j_2\,j_3\,j_4\,j_5^2 + 8266860\,j_4\,j_5^3 + 22104\,j_2^3\,j_3\,j_4\,j_6 \\
& - 663120\,j_3^3\,j_4\,j_6 - 1010880\,j_2\,j_3\,j_4^2\,j_6 + 36288\,j_2^2\,j_5\,j_6 - 1088640\,j_2\,j_3^2\,j_5\,j_6 - 1145016\,j_2^2\,j_4\,j_5\,j_6 - 10392624\,j_4^2\,j_5\,j_6 + 11634840\,j_3\,j_5^2\,j_6 \\
& - 9979200\,j_3\,j_4\,j_6^2 - 3470040\,j_2\,j_5\,j_6^2 + 504\,j_2^6\,j_7 - 30240\,j_2^3\,j_3^2\,j_7 + 453600\,j_3^4\,j_7 - 63504\,j_2^3\,j_4\,j_7 + 1905120\,j_2\,j_3^2\,j_4\,j_7 \\
& + 702756\,j_2^2\,j_4^2\,j_7 + 944784\,j_4^3\,j_7 - 5511240\,j_3\,j_4\,j_5\,j_7 - 260820\,j_2^3\,j_6\,j_7 + 4762800\,j_3^2\,j_6\,j_7 + 8485560\,j_2\,j_4\,j_6\,j_7 - 1530900\,j_2\,j_3\,j_7^2 \\
& + 22657320\,j_3\,j_4^2\,j_8 + 204120\,j_2^2\,j_5\,j_8 - 6123600\,j_3^2\,j_5\,j_8 - 6429780\,j_2\,j_4\,j_5\,j_8 + 3061800\,j_2\,j_3\,j_6\,j_8 + 9185400\,j_5\,j_6\,j_8 \\
& + 1530900\,j_2^2\,j_7\,j_8 - 27556200\,j_4\,j_7\,j_8 + 6426\,j_2^5\,j_9 - 192780\,j_2^2\,j_3^2\,j_9 - 428652\,j_2^3\,j_4\,j_9 + 5715360\,j_3^2\,j_4\,j_9 + 9552816\,j_2\,j_4^2\,j_9 \\
& + 3367980\,j_2\,j_3\,j_5\,j_9 - 49601160\,j_2^2\,j_9 + 3946320\,j_2^2\,j_6\,j_9 + 57386880\,j_4\,j_6\,j_9 - 36741600\,j_3\,j_7\,j_9 - 136080\,j_2^3\,j_3\,j_{10} + 4082400\,j_3^3\,j_{10} \\
& + 4592700\,j_2\,j_3\,j_4\,j_{10} - 2602530\,j_2^2\,j_5\,j_{10} - 66134880\,j_4\,j_5\,j_{10} - 85730400\,j_3\,j_6\,j_{10} + 27556200\,j_2\,j_7\,j_{10} + 440899200\,j_9\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -56\,j_2^{10} + 5040\,j_2^7\,j_3^2 - 151200\,j_2^4\,j_3^4 + 1512000\,j_2\,j_3^6 + 7560\,j_2^5\,j_4 - 453600\,j_2^2\,j_3^2\,j_4 + 6804000\,j_2^2\,j_3^4\,j_4 - 138780\,j_2^6\,j_4^2 - 1879200\,j_2^3\,j_3^2\,j_4^2 \\
& + 181278000\,j_3^4\,j_4^2 - 12009060\,j_2^2\,j_4^3 + 513361800\,j_2\,j_3^2\,j_4^3 + 362167200\,j_2^2\,j_4^4 + 476171136\,j_4^5 - 68040\,j_2^6\,j_3\,j_5 + 4082400\,j_2^3\,j_3^3\,j_5 \\
& - 61236000\,j_3^5\,j_5 + 6123600\,j_2^3\,j_3\,j_4\,j_5 - 183708000\,j_2\,j_3^3\,j_4\,j_5 - 137781000\,j_2^2\,j_3\,j_4^2\,j_5 + 411374700\,j_3\,j_4^3\,j_5 - 918540\,j_2^5\,j_5^2 + 27556200\,j_2^2\,j_3^2\,j_5^2 \\
& + 68890500\,j_2^3\,j_4\,j_5^2 - 826686000\,j_3^2\,j_4\,j_5^2 - 2341292850\,j_2\,j_4^2\,j_5^2 - 124002900\,j_2\,j_3\,j_5^3 + 3348078300\,j_4^2\,j_5^3 - 15750\,j_2^7\,j_6 + 945000\,j_2^4\,j_3^2\,j_6 \\
& - 14175000\,j_2^4\,j_3\,j_6 + 1417500\,j_2^5\,j_4\,j_6 - 42525000\,j_2^2\,j_3^2\,j_4\,j_6 - 67845600\,j_2^3\,j_4^2\,j_6 + 1078555500\,j_3^2\,j_4^2\,j_6 + 1261461600\,j_2\,j_4^3\,j_6 \\
& - 12757500\,j_2^2\,j_3\,j_5\,j_6 + 382725000\,j_3^3\,j_5\,j_6 + 574087500\,j_2\,j_3\,j_4\,j_5\,j_6 - 86113125\,j_2^2\,j_5^2\,j_6 - 4842018000\,j_4\,j_5^2\,j_6 - 14742000\,j_2^2\,j_6^2 \\
& + 442260000\,j_2\,j_3^2\,j_6^2 + 663390000\,j_2^2\,j_4\,j_6^2 + 1539648000\,j_4^2\,j_6^2 - 5970510000\,j_3\,j_5\,j_6^2 + 48478500\,j_2^3\,j_3\,j_4\,j_7 - 1454355000\,j_3^3\,j_4\,j_7 \\
& - 2181532500\,j_2\,j_3\,j_4^2\,j_7 + 22963500\,j_2^2\,j_5\,j_7 - 688905000\,j_2\,j_3^2\,j_5\,j_7 - 378897750\,j_2^2\,j_4\,j_5\,j_7 - 6200145000\,j_4^2\,j_5\,j_7 + 9300217500\,j_3\,j_5^2\,j_7 \\
& + 4363065000\,j_3\,j_4\,j_6\,j_7 + 1033357500\,j_2\,j_5\,j_6\,j_7 + 850500\,j_2^6\,j_8 - 51030000\,j_2^3\,j_3^2\,j_8 + 765450000\,j_3^4\,j_8 - 125023500\,j_2^4\,j_4\,j_8 \\
& + 3750705000\,j_2\,j_3^2\,j_4\,j_8 + 3903795000\,j_2^2\,j_4^2\,j_8 + 23147208000\,j_4^3\,j_8 - 19633792500\,j_3\,j_4\,j_5\,j_8 - 4650108750\,j_2\,j_5^2\,j_8 + 76545000\,j_2^3\,j_6\,j_8 \\
& - 2296350000\,j_2^3\,j_6\,j_8 - 7807590000\,j_2\,j_4\,j_6\,j_8 - 11573604000\,j_3\,j_4^2\,j_9 - 91854000\,j_2^2\,j_5\,j_9 + 2755620000\,j_3^2\,j_5\,j_9 + 826686000\,j_2\,j_4\,j_5\,j_9 \\
& + 16533720000\,j_5\,j_6\,j_9 - 244944000\,j_2^2\,j_4\,j_{10} + 7348320000\,j_3^2\,j_4\,j_{10} + 22596084000\,j_2\,j_4^2\,j_{10} - 37200870000\,j_5^2\,j_{10} - 22044960000\,j_4\,j_6\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 616\,j_2^{10} - 55440\,j_2^7\,j_3^2 + 1663200\,j_2^4\,j_3^4 - 16632000\,j_2\,j_3^6 - 76860\,j_2^8\,j_4 + 4611600\,j_2^5\,j_3^2\,j_4 - 69174000\,j_2^2\,j_3^4\,j_4 + 1205280\,j_2^6\,j_4^2 \\
& + 22939200\,j_2^3\,j_3^2\,j_4^2 - 1772928000\,j_3^4\,j_4^2 + 126996660\,j_2^4\,j_4^3 - 5111164800\,j_2\,j_3^2\,j_4^3 - 4209100200\,j_2^2\,j_4^4 - 5237882496\,j_4^5 + 748440\,j_2^6\,j_3\,j_5 \\
& - 44906400\,j_2^3\,j_3^3\,j_5 + 673596000\,j_3^5\,j_5 - 62256600\,j_2^4\,j_3\,j_4\,j_5 + 1867698000\,j_2\,j_3^3\,j_4\,j_5 + 1285956000\,j_2^2\,j_3\,j_4^2\,j_5 - 3186677700\,j_3\,j_4^3\,j_5 \\
& + 10103940\,j_2^5\,j_5^2 - 303118200\,j_2^2\,j_3^2\,j_5^2 - 723350250\,j_2^3\,j_4\,j_5^2 + 9093546000\,j_3^2\,j_4\,j_5^2 + 25724696850\,j_2\,j_4^2\,j_5^2 + 1364031900\,j_2\,j_3\,j_5^3 \\
& - 36828861300\,j_4\,j_5^3 + 570150\,j_2^7\,j_6 - 34209000\,j_2^4\,j_3^2\,j_6 + 513135000\,j_2\,j_3^4\,j_6 - 58903200\,j_2^5\,j_4\,j_6 + 1767096000\,j_2^2\,j_3^2\,j_4\,j_6 + 1798151400\,j_3^4\,j_4^2\,j_6 \\
& - 9061834500\,j_3^2\,j_4\,j_6^2 - 9782013600\,j_2\,j_4^3\,j_6 + 160744500\,j_2^3\,j_3\,j_5\,j_6 - 4822335000\,j_3^3\,j_5\,j_6 - 10307331000\,j_2\,j_3\,j_4\,j_5\,j_6 - 947244375\,j_2^2\,j_5^2\,j_6 \\
& + 18659484000\,j_4\,j_5^2\,j_6 + 151956000\,j_2^2\,j_6^2 - 4558680000\,j_2\,j_3^2\,j_6^2 - 3458376000\,j_2^2\,j_4\,j_6^2 - 14696640000\,j_4^2\,j_6^2 + 38578680000\,j_3\,j_5\,j_6^2 \\
& - 543469500\,j_2^3\,j_3\,j_4\,j_7 + 16304085000\,j_3^3\,j_4\,j_7 + 31138506000\,j_2\,j_3\,j_4^2\,j_7 - 229635000\,j_2^2\,j_5\,j_7 + 6889050000\,j_2\,j_3^2\,j_5\,j_7 \\
& + 5717911500\,j_2^2\,j_4\,j_5\,j_7 + 103985289000\,j_4^2\,j_5\,j_7 - 102302392500\,j_3\,j_5^2\,j_7 - 1082565000\,j_3\,j_4\,j_6\,j_7 - 27900652500\,j_2\,j_5\,j_6\,j_7 \\
& + 688905000\,j_2^3\,j_7^2 - 20667150000\,j_3^2\,j_7^2 - 22733865000\,j_2\,j_4\,j_7^2 - 9355500\,j_2^6\,j_8 + 561330000\,j_2^3\,j_3^2\,j_8 - 8419950000\,j_3^4\,j_8 + 1536003000\,j_2^2\,j_4\,j_8 \\
& - 46080090000\,j_2\,j_3^2\,j_4\,j_8 - 56857626000\,j_2^2\,j_4^2\,j_8 - 300913704000\,j_4^3\,j_8 + 199437997500\,j_3\,j_4\,j_5\,j_8 + 60451413750\,j_2\,j_5^2\,j_8 - 841995000\,j_2^3\,j_6\,j_8 \\
& + 25259850000\,j_3^2\,j_6\,j_8 + 110684070000\,j_2\,j_4\,j_6\,j_8 + 186004350000\,j_5\,j_7\,j_8 - 15309000\,j_2^2\,j_3\,j_9 + 459270000\,j_2\,j_3^2\,j_9 + 688905000\,j_2^2\,j_3\,j_4\,j_9 \\
& + 99202320000\,j_3\,j_4^2\,j_9 + 482233500\,j_2^3\,j_5\,j_9 - 20667150000\,j_2^2\,j_5\,j_9 + 13226976000\,j_2\,j_4\,j_5\,j_9 - 22044960000\,j_2\,j_3\,j_6\,j_9 - 4133430000\,j_2^2\,j_7\,j_9 \\
& - 198404640000\,j_4\,j_7\,j_9 + 15309000\,j_2^3\,j_{10} - 459270000\,j_2^2\,j_3^2\,j_{10} + 2602530000\,j_2^3\,j_4\,j_{10} - 98743050000\,j_3^2\,j_4\,j_{10} - 289891224000\,j_2\,j_4^2\,j_{10} \\
& + 6200145000\,j_2\,j_3\,j_5\,j_{10} + 372008700000\,j_5^2\,j_{10} + 22044960000\,j_2^2\,j_6\,j_{10} + 308629440000\,j_4\,j_6\,j_{10} + 124002900000\,j_3\,j_7\,j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 56\,j_2^{10} - 5040\,j_2^7 j_3^2 + 151200\,j_2^4 j_3^4 - 1512000\,j_2 j_3^6 - 4830\,j_2^8 j_4 + 289800\,j_2^5 j_3^2 j_4 - 4347000\,j_2^2 j_3^4 j_4 - 91800\,j_2^6 j_4^2 + 8343000\,j_2^3 j_3^2 j_4^2 \\
& - 167670000\,j_3^4 j_4^2 + 14360085\,j_2^7 j_4^3 - 418045050\,j_2 j_3^2 j_4^3 - 422528400\,j_2^4 j_4^4 - 51018336\,j_4^5 + 68040\,j_2^6 j_3 j_5 - 4082400\,j_2^3 j_3^3 j_5 + 61236000\,j_3^5 j_5 \\
& - 3912300\,j_2^4 j_3 j_4 j_5 + 117369000\,j_2 j_3^3 j_4 j_5 + 38272500\,j_2^2 j_3 j_4^2 j_5 - 897544800\,j_3^3 j_4^3 j_5 + 918540\,j_2^5 j_5^2 - 27556200\,j_2^2 j_3^2 j_5^2 - 53964225\,j_3^4 j_4 j_5^2 \\
& + 826686000\,j_3^2 j_4^2 j_5^2 + 2604060900\,j_2 j_4^3 j_5^2 + 124002900\,j_2 j_3 j_5^3 - 3348078300\,j_4^4 j_5^3 + 112140\,j_2^7 j_6 - 6728400\,j_2^4 j_3^2 j_6 + 100926000\,j_2 j_3^4 j_6 \\
& - 11907270\,j_2^5 j_4 j_6 + 357218100\,j_2^2 j_3^2 j_4 j_6 + 370978380\,j_3^4 j_4^2 j_6 - 1867041900\,j_3^2 j_4^2 j_6 - 2435443200\,j_2 j_4^3 j_6 + 4592700\,j_2^3 j_3 j_5 j_6 \\
& - 137781000\,j_3^3 j_5 j_6 - 941612850\,j_2 j_3 j_4 j_5 j_6 - 551124000\,j_2^2 j_5^2 j_6 + 7581891600\,j_4 j_5^2 j_6 + 21120750\,j_2^4 j_6^2 - 633622500\,j_2^2 j_3^2 j_6^2 \\
& + 130734000\,j_2^2 j_4 j_6^2 - 4776408000\,j_4^2 j_6^2 + 4822335000\,j_3 j_5 j_6^2 - 44396100\,j_2^3 j_3 j_4 j_7 + 1331883000\,j_3^3 j_4 j_7 + 3699419850\,j_2 j_3 j_4^2 j_7 \\
& - 13012650\,j_2^4 j_5 j_7 + 390379500\,j_2 j_3^2 j_5 j_7 + 568346625\,j_2^2 j_4 j_5 j_7 + 744017400\,j_4^2 j_5 j_7 - 9300217500\,j_3 j_5^2 j_7 + 3444525000\,j_3 j_4 j_6 j_7 \\
& - 3444525000\,j_2 j_5 j_6 j_7 - 850500\,j_2^6 j_8 + 51030000\,j_2^3 j_3^2 j_8 - 765450000\,j_3^4 j_8 + 164061450\,j_2^4 j_4 j_8 - 4921843500\,j_2 j_3^2 j_4 j_8 - 7362098100\,j_2^2 j_4^2 j_8 \\
& - 5456127600\,j_4^3 j_8 + 17980420500\,j_3 j_4 j_5 j_8 + 8680203000\,j_2 j_5^2 j_8 + 76545000\,j_2^3 j_6 j_8 - 2296350000\,j_2^2 j_4 j_6 j_8 + 5051970000\,j_2 j_4 j_6 j_8 \\
& + 13778100000\,j_6^2 j_8 - 6633900\,j_2^4 j_3 j_9 + 199017000\,j_2 j_3^3 j_9 + 298525500\,j_2^2 j_3 j_4 j_9 + 7936185600\,j_3 j_4^2 j_9 - 6889050\,j_2^3 j_5 j_9 \\
& - 2480058000\,j_3^2 j_5 j_9 + 165337200\,j_2 j_4 j_5 j_9 - 5970510000\,j_2 j_3 j_6 j_9 + 6633900\,j_2^2 j_{10} - 199017000\,j_2^2 j_3^2 j_{10} - 169929900\,j_2^2 j_4 j_{10} \\
& - 3857868000\,j_3^2 j_4 j_{10} - 22320522000\,j_2 j_4^2 j_{10} + 2686729500\,j_2 j_3 j_5 j_{10} + 33480783000\,j_5^2 j_{10} + 5970510000\,j_2^2 j_6 j_{10} - 33067440000\,j_4 j_6 j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 616\,j_2^{10} - 55440\,j_2^7 j_3^2 + 1663200\,j_2^4 j_3^4 - 16632000\,j_2 j_3^6 - 49140\,j_2^8 j_4 + 2948400\,j_2^5 j_3^2 j_4 - 44226000\,j_2^2 j_3^4 j_4 - 1546560\,j_2^6 j_4^2 \\
& + 113205600\,j_2^3 j_3^2 j_4^2 - 2004264000\,j_3^4 j_4^2 + 179105580\,j_2^4 j_4^3 - 4990442400\,j_2 j_3^2 j_4^3 - 4995545400\,j_2^2 j_4^4 - 1836660096\,j_4^5 + 748440\,j_2^6 j_3 j_5 \\
& - 44906400\,j_2^3 j_3^3 j_5 + 673596000\,j_2^5 j_5 - 39803400\,j_2^4 j_3 j_4 j_5 + 1194102000\,j_2 j_3^3 j_4 j_5 + 275562000\,j_2^2 j_3 j_4^2 j_5 - 9733243500\,j_3 j_4^3 j_5 \\
& + 10103940\,j_2^5 j_4 j_5 - 303118200\,j_2^2 j_3^2 j_5^2 - 571791150\,j_2^2 j_4 j_5^2 + 9093546000\,j_3^2 j_4 j_5^2 + 29604216150\,j_2 j_4^2 j_5^2 + 1364031900\,j_2 j_3 j_5^3 \\
& - 36828861300\,j_5^4 + 1318590\,j_2^7 j_6 - 79115400\,j_2^4 j_3^2 j_6 + 1186731000\,j_2 j_3^4 j_6 - 137884680\,j_2^5 j_4 j_6 + 4136540400\,j_2^2 j_3^2 j_4 j_6 \\
& + 4083926040\,j_3^2 j_4^2 j_6 - 16477805700\,j_2^3 j_4^2 j_6 - 20814991200\,j_2 j_4^3 j_6 + 95426100\,j_2^3 j_3 j_5 j_6 - 2862783000\,j_3^3 j_5 j_6 - 12074864400\,j_2 j_3 j_4 j_5 j_6 \\
& - 5921138475\,j_2^2 j_5^2 j_6 + 57112192800\,j_4 j_5^2 j_6 + 274428000\,j_2^4 j_6^2 - 8232840000\,j_2 j_3^2 j_6^2 + 239112000\,j_2^2 j_4 j_6^2 - 41290560000\,j_4^2 j_6^2 \\
& + 78994440000\,j_3 j_5 j_6^2 - 510810300\,j_2^3 j_3 j_4 j_7 + 15324309000\,j_3^3 j_4 j_7 + 41435339400\,j_2 j_3 j_4^2 j_7 - 128595600\,j_2^4 j_5 j_7 + 3857868000\,j_2 j_3^2 j_5 j_7 \\
& + 4767222600\,j_2^2 j_4 j_5 j_7 + 44393038200\,j_4^2 j_5 j_7 - 102302392500\,j_3^2 j_5^2 j_7 - 21880935000\,j_3 j_4 j_6 j_7 - 42367657500\,j_2 j_5 j_6 j_7 \\
& - 229635000\,j_2^2 j_7^2 + 6889050000\,j_3^2 j_7^2 + 2066715000\,j_2 j_4 j_7^2 + 41334300000\,j_6 j_7^2 - 9355500\,j_2^7 j_8 + 561330000\,j_2^3 j_3^2 j_8 - 8419950000\,j_3^4 j_8 \\
& + 1839121200\,j_2^4 j_4 j_8 - 55173636000\,j_2 j_3^2 j_4 j_8 - 82264442400\,j_2^2 j_4^2 j_8 - 159385060800\,j_4^3 j_8 + 227545321500\,j_3 j_4 j_5 j_8 + 101372370750\,j_2 j_5^2 j_8 \\
& + 535815000\,j_2^3 j_6 j_8 - 16074450000\,j_3^2 j_6 j_8 + 102417210000\,j_2 j_4 j_6 j_8 - 82668600\,j_2^4 j_3 j_9 + 2480058000\,j_2 j_3^3 j_9 + 3720087000\,j_2^2 j_3 j_4 j_9 \\
& + 119704132800\,j_3 j_4^2 j_9 - 41334300\,j_2^3 j_5 j_9 - 32240754000\,j_3^2 j_5 j_9 + 992023200\,j_2 j_4 j_5 j_9 - 66134880000\,j_2 j_3 j_6 j_9 + 4133430000\,j_2^2 j_7 j_9 \\
& - 99202320000\,j_4 j_7 j_9 + 82668600\,j_2^5 j_{10} - 2480058000\,j_2^2 j_3^2 j_{10} - 2431069200\,j_2^3 j_4 j_{10} - 38670534000\,j_3^2 j_4 j_{10} - 262114574400\,j_2 j_4^2 j_{10} \\
& + 33480783000\,j_2 j_3 j_5 j_{10} + 342248004000\,j_5^2 j_{10} + 66134880000\,j_2^2 j_6 j_{10} + 44089920000\,j_4 j_6 j_{10} - 124002900000\,j_3 j_7 j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -4312\,j_2^{10} + 388080\,j_2^7 j_3^2 - 11642400\,j_2^4 j_3^4 + 116424000\,j_2 j_3^6 + 861420\,j_2^8 j_4 - 51685200\,j_2^5 j_3^2 j_4 + 775278000\,j_2^2 j_3^4 j_4 - 39329010\,j_2^6 j_4^2 \\
& + 849536100\,j_2^3 j_3^2 j_4^2 + 9910026000\,j_3^4 j_4^2 - 199178595\,j_2^4 j_4^3 + 33391225350\,j_2 j_3^2 j_4^3 + 13993956900\,j_2^2 j_4^4 - 22856214528\,j_4^5 \\
& - 5239080\,j_2^6 j_3 j_5 + 314344800\,j_2^3 j_3^3 j_5 - 4715172000\,j_3^5 j_5 + 697750200\,j_2^4 j_3 j_4 j_5 - 20932506000\,j_2 j_3^3 j_4 j_5 - 20387760750\,j_2^2 j_3 j_4^2 j_5 \\
& + 57992022900\,j_3 j_4^3 j_5 - 70727580\,j_2^5 j_5^2 + 2121827400\,j_2^2 j_3^2 j_5^2 + 6831641250\,j_2^2 j_4 j_5^2 - 63654822000\,j_3^2 j_4 j_5^2 - 131195068200\,j_2 j_4^2 j_5^2 \\
& - 9548223300\,j_2 j_3 j_5^3 + 257802029100\,j_5^4 + 4740750\,j_2^7 j_6 - 284445000\,j_2^4 j_3^2 j_6 + 4266675000\,j_2 j_3^4 j_6 - 457531200\,j_2^5 j_4 j_6 \\
& + 13725936000\,j_2^2 j_3^2 j_4 j_6 + 5540157000\,j_2^3 j_4^2 j_6 + 162696397500\,j_3^2 j_4^2 j_6 + 217767463200\,j_2 j_4^3 j_6 - 696559500\,j_2^3 j_3 j_5 j_6 + 20896785000\,j_3^3 j_5 j_6 \\
& + 27801144000\,j_2 j_3 j_4 j_5 j_6 - 40145938875\,j_2^2 j_5^2 j_6 - 939941982000\,j_4 j_5^2 j_6 - 358202250\,j_2^4 j_6^2 + 10746067500\,j_2^2 j_3^2 j_6^2 + 99499995000\,j_2^2 j_4 j_6^2 \\
& + 546924960000\,j_4^2 j_6^2 - 57868020000\,j_3 j_5 j_6^2 + 3589960500\,j_2^3 j_3 j_4 j_7 - 107698815000\,j_3^3 j_4 j_7 - 130754169000\,j_2 j_3 j_4^2 j_7 + 2786238000\,j_2^2 j_5 j_7 \\
& - 83587140000\,j_2 j_3^2 j_5 j_7 - 67994923500\,j_2^2 j_4 j_5 j_7 + 494771571000\,j_4^2 j_5 j_7 + 716116747500\,j_3 j_5^2 j_7 - 4822335000\,j_2^2 j_3 j_6 j_7 \\
& - 1107070335000\,j_3 j_4 j_6 j_7 - 16878172500\,j_2 j_5 j_6 j_7 - 12055837500\,j_3^2 j_7^2 + 289340100000\,j_2^2 j_3^2 j_7 + 65101522500\,j_2 j_4 j_7^2 + 65488500\,j_2^6 j_8 \\
& - 3929310000\,j_2^3 j_3^2 j_8 + 58939650000\,j_3^4 j_8 - 7215642000\,j_2^2 j_4 j_8 + 216469260000\,j_2 j_3^2 j_4 j_8 + 143384094000\,j_2^2 j_4^2 j_8 - 1180507608000\,j_4^3 j_8 \\
& - 585913702500\,j_3 j_4 j_5 j_8 + 54251268750\,j_2 j_5^2 j_8 + 10716300000\,j_2^3 j_6 j_8 - 176818950000\,j_2^2 j_6 j_8 - 22504230000\,j_2 j_4 j_6 j_8 \\
& + 72335025000\,j_2 j_3 j_7 j_8 - 589396500\,j_2^4 j_3 j_9 + 17681895000\,j_2 j_3^3 j_9 + 28934010000\,j_2^2 j_3 j_4 j_9 - 11573604000\,j_3 j_4^2 j_9 - 8680203000\,j_3^2 j_5 j_9 \\
& + 57868020000\,j_3 j_5 j_9 - 23147208000\,j_2 j_4 j_5 j_9 - 183248730000\,j_2 j_3 j_6 j_9 + 159137055000\,j_2^2 j_7 j_9 - 1041624360000\,j_4 j_7 j_9 + 678699000\,j_2^5 j_{10} \\
& - 20360970000\,j_2^2 j_3^2 j_{10} - 54170896500\,j_2^2 j_4 j_{10} + 636548220000\,j_3^2 j_4 j_{10} + 1157360400000\,j_2 j_4^2 j_{10} + 238705582500\,j_2 j_3 j_5 j_{10} \\
& - 4687309620000\,j_5^2 j_{10} + 234686970000\,j_2^2 j_6 j_{10} + 9126613440000\,j_4 j_6 j_{10} - 5208121800000\,j_3 j_7 j_{10} + 20832487200000\,j_{10}^2,
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 14\,j_2^7 j_3 j_4 - 840\,j_2^4 j_3^3 j_4 + 12600\,j_2 j_3^5 j_4 - 1260\,j_2^5 j_3 j_4^2 + 37800\,j_2^2 j_3^3 j_4^2 + 22050\,j_2^3 j_3 j_4^3 + 189000\,j_3^3 j_4^3 - 897480\,j_2 j_3 j_4^4 + 252\,j_2^6 j_4 j_5 \\
& - 3780\,j_2^3 j_3^2 j_4 j_5 - 113400\,j_3^4 j_4 j_5 - 37125\,j_2^4 j_4^2 j_5 + 603450\,j_2 j_3^2 j_4^2 j_5 + 1046844\,j_2^2 j_4^3 j_5 + 2834352\,j_4^4 j_5 + 76545\,j_2^5 j_3 j_4 j_5 \\
& - 2930580\,j_3^2 j_4^2 j_5^2 - 1377810\,j_2 j_4^3 j_5^2 + 294\,j_2^7 j_3 j_6 - 17640\,j_2^4 j_3^3 j_6 + 264600\,j_2 j_3^5 j_6 - 43326\,j_2^5 j_3 j_4 j_6 + 1299780\,j_2^2 j_3^3 j_4 j_6 \\
& + 1354320\,j_2^3 j_3 j_4^2 j_6 + 162000\,j_3^3 j_4^2 j_6 - 14364\,j_2^5 j_5 j_6 + 430920\,j_2^2 j_3^2 j_5 j_6 + 527094\,j_2^3 j_4 j_5 j_6 - 3252150\,j_3^2 j_4 j_5 j_6 - 1411344\,j_2 j_4^2 j_5 j_6 \\
& - 7424865\,j_2 j_3 j_5^2 j_6 + 20207880\,j_5^3 j_6 - 7560\,j_2^6 j_3 j_6^2 + 226800\,j_2^3 j_3^3 j_6^2 + 7850520\,j_2 j_3 j_4 j_6^2 - 1803060\,j_2^2 j_5 j_6^2 - 3110400\,j_4 j_5 j_6^2 \\
& - 84\,j_2^7 j_7 + 5040\,j_2^4 j_3^2 j_7 - 75600\,j_2 j_3^4 j_7 + 14742\,j_2^2 j_4 j_7 - 442260\,j_2^2 j_3^2 j_4 j_7 + 648\,j_3^2 j_4^2 j_7 + 31590\,j_3^2 j_4 j_7 - 3254256\,j_2 j_4^2 j_7 \\
& + 17010\,j_2^2 j_3 j_5 j_7 - 510300\,j_3^2 j_5 j_7 + 8190315\,j_2 j_3 j_4 j_5 j_7 - 38578680\,j_4 j_5^2 j_7 + 128520\,j_2^2 j_6 j_7 - 3855600\,j_2 j_3^2 j_6 j_7 + 447120\,j_2^2 j_4 j_6 j_7 \\
& - 15863040\,j_4^2 j_6 j_7 + 10716300\,j_3 j_5 j_6 j_7 + 29087100\,j_3 j_4 j_7^2 + 119070\,j_2^3 j_3 j_4 j_8 - 3572100\,j_3^3 j_4 j_8 - 20207880\,j_2 j_3 j_4^2 j_8 - 34020\,j_2^4 j_5 j_8 \\
& + 1020600\,j_2 j_3^2 j_5 j_8 - 2908710\,j_2^2 j_4 j_5 j_8 + 123084360\,j_4^2 j_5 j_8 + 6889050\,j_3 j_5^2 j_8 - 27556200\,j_3 j_4 j_6 j_8 + 4592700\,j_2 j_5 j_6 j_8 + 510300\,j_2^2 j_7 j_8 \\
& - 15309000\,j_3^2 j_7 j_8 - 33679800\,j_2 j_4 j_7 j_8 - 378\,j_2^6 j_9 - 11340\,j_2^3 j_3^2 j_9 + 680400\,j_3^4 j_9 + 6804\,j_2^2 j_4 j_9 + 1326780\,j_2 j_3^2 j_4 j_9 - 3102624\,j_2^2 j_4^2 j_9 \\
& - 1959552\,j_4^3 j_9 - 459270\,j_2^2 j_3 j_5 j_9 - 22657320\,j_3 j_4 j_5 j_9 + 8266860\,j_2 j_5^2 j_9 - 1360800\,j_2^3 j_6 j_9 - 8164800\,j_3^2 j_6 j_9 - 12247200\,j_2 j_4 j_6 j_9 \\
& - 9185400\,j_2 j_3 j_7 j_9 + 110224800\,j_5 j_7 j_9 + 34020\,j_2^3 j_3 j_{10} - 1020600\,j_2 j_3^2 j_{10} - 1530900\,j_2^2 j_3 j_4 j_{10} + 12247200\,j_3 j_4^2 j_{10} - 51030\,j_2^2 j_5 j_{10} \\
& + 15309000\,j_3^2 j_5 j_{10} + 13471920\,j_2 j_4 j_5 j_{10} + 48988800\,j_2 j_3 j_6 j_{10} - 220449600\,j_5 j_6 j_{10} + 9185400\,j_2^2 j_7 j_{10} - 146966400\,j_4 j_7 j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 350\,j_2^7 j_3 j_4 - 21000\,j_2^4 j_3^3 j_4 + 315000\,j_2 j_3^5 j_4 - 31500\,j_2^5 j_3 j_4^2 + 945000\,j_2^2 j_3^3 j_4^2 + 587655\,j_2^3 j_3 j_4^3 + 3632850\,j_3^3 j_4^3 - 16694100\,j_2 j_3 j_4^4 \\
& + 4536\,j_2^6 j_4 j_5 + 11340\,j_2^3 j_3^2 j_4 j_5 - 4422600\,j_3^4 j_4 j_5 - 487944\,j_2^4 j_4^2 j_5 + 1880820\,j_2^2 j_3^2 j_4^2 j_5 + 12203460\,j_2^2 j_4^3 j_5 + 11337408\,j_4^4 j_5 \\
& + 1913625\,j_2^2 j_3 j_4 j_5^2 + 12006630\,j_3 j_4^2 j_5^2 - 24800580\,j_2 j_4 j_5^3 + 6426\,j_2^7 j_3 j_6 - 385560\,j_2^4 j_3^3 j_6 + 5783400\,j_2^2 j_3^5 j_6 - 810990\,j_2^3 j_3 j_4 j_6 \\
& + 24329700\,j_2 j_3^3 j_4 j_6 + 23481900\,j_2^2 j_3 j_4^2 j_6 + 583200\,j_3 j_4^3 j_6 - 195048\,j_2^5 j_5 j_6 + 5851440\,j_2^2 j_3^2 j_5 j_6 + 7635060\,j_2^3 j_4 j_5 j_6 - 59960250\,j_2^2 j_3 j_4 j_5 j_6 \\
& - 2099520\,j_2 j_4^2 j_5 j_6 - 114128595\,j_2 j_3 j_5^2 j_6 + 99202320\,j_5^3 j_6 + 425250\,j_2^2 j_3 j_6^2 - 12757500\,j_3^2 j_4 j_6^2 + 119475000\,j_2 j_3 j_4 j_6^2 - 10206000\,j_2^2 j_5 j_6^2 \\
& - 143467200\,j_4 j_5 j_6^2 - 1512\,j_2^7 j_7 + 90720\,j_2^4 j_3^2 j_7 - 1360800\,j_2 j_3^4 j_7 + 233604\,j_2^5 j_4 j_7 - 7008120\,j_2^2 j_3^2 j_4 j_7 - 2980152\,j_2^3 j_4^2 j_7 \\
& + 84046410\,j_3^3 j_4^2 j_7 - 14696640\,j_2 j_4^3 j_7 + 663390\,j_2^3 j_3 j_5 j_7 - 19901700\,j_3^3 j_5 j_7 + 84276045\,j_2 j_3 j_4 j_5 j_7 - 99202320\,j_4 j_5^2 j_7 \\
& + 1701000\,j_2^2 j_6 j_7 - 51030000\,j_2 j_3^2 j_6 j_7 - 34700400\,j_2^2 j_4 j_6 j_7 - 22963500\,j_3 j_5 j_6 j_7 + 2602530\,j_2^2 j_3 j_4 j_8 - 78075900\,j_3^2 j_4 j_8 \\
& - 335267100\,j_2 j_3 j_4^2 j_8 - 612360\,j_2^4 j_5 j_8 + 18370800\,j_2 j_3^2 j_5 j_8 - 11941020\,j_2^2 j_4 j_5 j_8 + 363741840\,j_4^2 j_5 j_8 + 268672950\,j_3 j_5^2 j_8 \\
& + 275562000\,j_3 j_4 j_6 j_8 + 137781000\,j_2 j_5 j_6 j_8 - 12096\,j_2^7 j_9 - 124740\,j_2^3 j_3^2 j_9 + 14628600\,j_3^4 j_9 + 789264\,j_2^2 j_4 j_9 + 14594580\,j_2 j_3^2 j_4 j_9 \\
& - 36741600\,j_2^2 j_4^2 j_9 - 17635968\,j_4^3 j_9 - 11481750\,j_2^2 j_3 j_5 j_9 - 66134880\,j_3 j_4 j_5 j_9 + 148803480\,j_2^2 j_5^2 j_9 - 17690400\,j_2^2 j_6 j_9 \\
& - 234738000\,j_3^2 j_6 j_9 - 24494400\,j_2 j_4 j_6 j_9 + 489888000\,j_6^2 j_9 + 850500\,j_2^4 j_3 j_{10} - 25515000\,j_2 j_3^3 j_{10} - 38272500\,j_2^2 j_3 j_4 j_{10} \\
& - 330674400\,j_3 j_4^2 j_{10} + 6123600\,j_2^3 j_5 j_{10} + 160744500\,j_3^2 j_5 j_{10} - 55112400\,j_2 j_4 j_5 j_{10} + 765450000\,j_2 j_3 j_6 j_{10} - 1102248000\,j_5 j_6 j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -2702\,j_2^7 j_3 j_4 + 162120\,j_2^4 j_3^3 j_4 - 2431800\,j_2 j_3^5 j_4 + 243180\,j_2^5 j_3 j_4^2 - 7295400\,j_2^2 j_3^3 j_4^2 - 5174442\,j_2^3 j_3 j_4^3 - 8913240\,j_3^3 j_4^3 \\
& + 153148320\,j_2 j_3 j_4^4 - 61236\,j_2^6 j_4 j_5 + 1485540\,j_2^3 j_3^2 j_4 j_5 + 10546200\,j_3^4 j_4 j_5 + 6241941\,j_2^4 j_4^2 j_5 - 88770330\,j_2^2 j_3^2 j_4^2 j_5 - 162170424\,j_2^2 j_4^3 j_5 \\
& + 8503056\,j_4^4 j_5 - 14773185\,j_2^2 j_3 j_4 j_5^2 - 28737180\,j_3 j_4^2 j_5^2 + 334807830\,j_2 j_4 j_5^3 - 13230\,j_2^7 j_3 j_6 + 793800\,j_2^3 j_3^3 j_6 - 11907000\,j_3^5 j_6 \\
& + 2873358\,j_2^5 j_3 j_4 j_6 - 86200740\,j_2 j_3^3 j_4 j_6 - 102510360\,j_2^2 j_3 j_4^2 j_6 - 7639920\,j_3 j_4^3 j_6 + 2374596\,j_2^4 j_5 j_6 - 71237880\,j_2^2 j_3^2 j_5 j_6 \\
& - 89948394\,j_2^3 j_4 j_5 j_6 + 174223710\,j_3^2 j_4 j_5 j_6 - 299461536\,j_2 j_4^2 j_5 j_6 + 1034046405\,j_2 j_3 j_5^2 j_6 - 545612760\,j_5^3 j_6 + 340200\,j_2^3 j_3^2 j_6^2 \\
& - 10206000\,j_3^3 j_6^2 - 1055899800\,j_2 j_3 j_4 j_6^2 + 111755700\,j_2^2 j_5 j_6^2 + 545875200\,j_4 j_5 j_6^2 + 20412\,j_2^7 j_7 - 1224720\,j_2^4 j_3^2 j_7 + 18370800\,j_2 j_3^4 j_7 \\
& - 3024378\,j_2^5 j_4 j_7 + 90731340\,j_2^2 j_3^2 j_4 j_7 + 48522240\,j_3^2 j_4^2 j_7 - 205074990\,j_2^3 j_4 j_7 + 60466176\,j_2 j_4^2 j_7 - 1581930\,j_2^3 j_3 j_5 j_7 \\
& + 47457900\,j_3^3 j_5 j_7 - 939896055\,j_2 j_3 j_4 j_5 j_7 + 545612760\,j_4 j_5^2 j_7 - 18370800\,j_2^2 j_6 j_7 + 551124000\,j_2 j_3^2 j_6 j_7 + 468018000\,j_2^2 j_4 j_6 j_7 \\
& - 451396800\,j_4^2 j_6 j_7 - 390379500\,j_3 j_5 j_6 j_7 + 344452500\,j_3 j_4 j_7^2 - 5358150\,j_2^3 j_3 j_4 j_8 + 160744500\,j_3^2 j_4 j_8 + 1833405840\,j_2 j_3 j_4^2 j_8 \\
& + 8266860\,j_2^4 j_5 j_8 - 248005800\,j_2 j_3^2 j_5 j_8 + 85883490\,j_2^2 j_4 j_5 j_8 - 1173894120\,j_4^2 j_5 j_8 - 640681650\,j_3 j_5^2 j_8 + 413343000\,j_3 j_4 j_6 j_8 \\
& - 2824510500\,j_2 j_5 j_6 j_8 + 7654500\,j_2^3 j_7 j_8 - 229635000\,j_3^2 j_7 j_8 - 413343000\,j_2 j_4 j_7 j_8 + 2755620000\,j_6 j_7 j_8 + 184842\,j_2^6 j_9 \\
& - 4524660\,j_2^3 j_3^2 j_9 - 30618000\,j_3^4 j_9 + 15683220\,j_2^2 j_4 j_9 + 175032900\,j_2 j_3^2 j_4 j_9 + 579537504\,j_2^2 j_4^2 j_9 - 123451776\,j_4^3 j_9 + 88639110\,j_2^2 j_3 j_5 j_9 \\
& + 259028280\,j_3 j_4 j_5 j_9 - 2008846980\,j_2 j_5^2 j_9 + 175543200\,j_2^3 j_6 j_9 + 367416000\,j_3^2 j_6 j_9 + 1775844000\,j_2 j_4 j_6 j_9 - 137781000\,j_2 j_3 j_7 j_9 \\
& - 6565860\,j_2^4 j_3 j_{10} + 196975800\,j_2 j_3^3 j_{10} + 295463700\,j_2^2 j_3 j_4 j_{10} + 308629440\,j_3 j_4^2 j_{10} - 81596970\,j_2^3 j_5 j_{10} - 211264200\,j_3^2 j_5 j_{10} \\
& + 1499057280\,j_2 j_4 j_5 j_{10} - 5633712000\,j_2 j_3 j_6 j_{10} + 5511240000\,j_5 j_6 j_{10} + 137781000\,j_2^2 j_7 j_{10} - 3306744000\,j_4 j_7 j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = &-3374\, j_2^7\, j_3\, j_4 + 202440\, j_2^4\, j_3^3\, j_4 - 3036600\, j_2\, j_3^5\, j_4 + 303660\, j_2^5\, j_3\, j_4^2 - 9109800\, j_2^2\, j_3^3\, j_4^2 - 6342498\, j_2^3\, j_3\, j_4^3 - 14695560\, j_3^3\, j_4^3 \\
&+ 186990120\, j_2\, j_3\, j_4^4 - 68796\, j_2^6\, j_4\, j_5 + 1394820\, j_2^3\, j_3^2\, j_4\, j_5 + 20071800\, j_3^4\, j_4\, j_5 + 7003395\, j_2^4\, j_4^2\, j_5 - 87119550\, j_2\, j_3^2\, j_4^2\, j_5 \\
&- 180571842\, j_2^2\, j_4^3\, j_5 + 30862944\, j_4^4\, j_5 - 18447345\, j_2^2\, j_3\, j_4\, j_5^2 - 80940870\, j_3\, j_4^2\, j_5^2 + 376142130\, j_2\, j_4\, j_5^3 - 27510\, j_2^6\, j_3\, j_6 + 1650600\, j_2^3\, j_3^3\, j_6 \\
&- 24759000\, j_2^5\, j_6 + 4605246\, j_2^4\, j_3\, j_4\, j_6 - 138157380\, j_2\, j_3^3\, j_4\, j_6 - 151528320\, j_2^2\, j_3\, j_4^2\, j_6 - 6972480\, j_3\, j_4^3\, j_6 + 2696652\, j_2^5\, j_5\, j_6 \\
&- 80899560\, j_2^2\, j_3^2\, j_5\, j_6 - 102476502\, j_2^2\, j_4\, j_5\, j_6 + 296199990\, j_3^2\, j_4\, j_5\, j_6 - 301430808\, j_2\, j_4^2\, j_5\, j_6 + 1242554985\, j_2\, j_3\, j_5^2\, j_6 - 464781240\, j_5^3\, j_6 \\
&- 604800\, j_2^3\, j_3\, j_6^2 + 18140000\, j_3^3\, j_6^2 - 1279508400\, j_2\, j_3\, j_4\, j_6^2 + 129616200\, j_2^2\, j_5\, j_6^2 + 524880000\, j_4\, j_5\, j_6^2 + 22932\, j_2^7\, j_7 - 1375920\, j_2^4\, j_3^2\, j_7 \\
&+ 20638800\, j_2\, j_3^3\, j_7 - 3412206\, j_2^5\, j_4\, j_7 + 102366180\, j_2^2\, j_3^2\, j_4\, j_7 + 53584416\, j_2^3\, j_4^2\, j_7 - 405316710\, j_2^3\, j_3\, j_4\, j_7 + 83800008\, j_2\, j_4^3\, j_7 \\
&- 3010770\, j_2^3\, j_3\, j_5\, j_7 + 90323100\, j_3^3\, j_5\, j_7 - 1084106835\, j_2\, j_3\, j_4\, j_5\, j_7 + 349963740\, j_4\, j_5^2\, j_7 - 20865600\, j_2^4\, j_6\, j_7 + 625968000\, j_2\, j_3^2\, j_6\, j_7 \\
&+ 513434700\, j_2^2\, j_4\, j_6\, j_7 - 451396800\, j_4^2\, j_6\, j_7 + 68890500\, j_3\, j_5\, j_6\, j_7 + 229635000\, j_3\, j_4\, j_7^2 + 688905000\, j_7^3 - 11141550\, j_2^3\, j_3\, j_4\, j_8 \\
&+ 334246500\, j_3^3\, j_4\, j_8 + 2512513080\, j_2\, j_3\, j_4^2\, j_8 + 9287460\, j_2^4\, j_5\, j_8 - 278623800\, j_2\, j_3^2\, j_5\, j_8 + 105172830\, j_2^2\, j_4\, j_5\, j_8 - 1260236880\, j_4^2\, j_5\, j_8 \\
&- 1219361850\, j_3\, j_5^2\, j_8 - 520506000\, j_3\, j_4\, j_6\, j_8 - 3191926500\, j_2\, j_5\, j_6\, j_8 + 7654500\, j_2^3\, j_7\, j_8 - 229635000\, j_2^3\, j_7\, j_8 - 298525500\, j_2\, j_4\, j_7\, j_8 \\
&+ 205254\, j_2^6\, j_9 - 4116420\, j_2^3\, j_3^2\, j_9 - 61236000\, j_3^4\, j_9 - 17064432\, j_2^4\, j_4\, j_9 + 142986060\, j_2\, j_3^2\, j_4\, j_9 + 641875752\, j_2^2\, j_4^2\, j_9 - 219469824\, j_4^3\, j_9 \\
&+ 110684070\, j_2^2\, j_3\, j_5\, j_9 + 264539520\, j_3\, j_4\, j_5\, j_9 - 2256852780\, j_2\, j_5^2\, j_9 + 204120000\, j_2^3\, j_6\, j_9 + 979776000\, j_3^2\, j_6\, j_9 + 1837080000\, j_2\, j_4\, j_6\, j_9 \\
&- 137781000\, j_2\, j_3\, j_7\, j_9 - 8198820\, j_2^4\, j_3\, j_{10} + 245964600\, j_2\, j_3^3\, j_{10} + 368946900\, j_2^2\, j_3\, j_4\, j_{10} + 1179405360\, j_3\, j_4^2\, j_{10} - 91700910\, j_2^2\, j_5\, j_{10} \\
&- 569494800\, j_3^2\, j_5\, j_{10} + 1603770840\, j_2\, j_4\, j_5\, j_{10} - 7103376000\, j_2\, j_3\, j_6\, j_{10} + 4408992000\, j_5\, j_6\, j_{10} + 137781000\, j_2^2\, j_7\, j_{10} - 551124000\, j_4\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = &-392\, j_2^{11} + 35280\, j_2^8\, j_3^2 - 1058400\, j_2^5\, j_3^4 + 10584000\, j_2^2\, j_3^6 + 52220\, j_2^9\, j_4 - 3427200\, j_2^6\, j_3^2\, j_4 + 64638000\, j_2^3\, j_3^4\, j_4 - 264600000\, j_3^6\, j_4 \\
&- 1466060\, j_2^7\, j_4^2 + 44871600\, j_2^4\, j_3^2\, j_4^2 - 26694000\, j_2\, j_3^4\, j_4^2 - 34824420\, j_2^5\, j_4^3 + 1478487600\, j_2^2\, j_3^2\, j_4^3 + 1906270200\, j_3^2\, j_4^4 + 10926981000\, j_2^3\, j_4^4 \\
&- 6936954048\, j_2\, j_4^5 - 476280\, j_2^7\, j_3\, j_5 + 28576800\, j_2^4\, j_3^3\, j_5 - 428652000\, j_2\, j_3^5\, j_5 + 34360200\, j_2^5\, j_3\, j_4\, j_5 - 1030806000\, j_2^2\, j_3^3\, j_4\, j_5 \\
&- 414963000\, j_2^3\, j_3\, j_4^2\, j_5 - 5003370000\, j_3^3\, j_4^2\, j_5 - 3794372100\, j_2\, j_3\, j_4^3\, j_5 - 6429780\, j_2^6\, j_5^2 + 192893400\, j_2^3\, j_3^2\, j_5^2 + 417170250\, j_2^4\, j_4\, j_5^2 \\
&- 5557167000\, j_2\, j_3^2\, j_4\, j_5^2 - 15345304200\, j_2^2\, j_4^2\, j_5^2 + 20076660000\, j_4^3\, j_5^2 - 868020300\, j_2^2\, j_3\, j_5^3 + 18600435000\, j_3\, j_4\, j_5^3 + 23436548100\, j_2\, j_5^4 \\
&- 330750\, j_2^8\, j_6 + 19845000\, j_2^5\, j_3^2\, j_6 - 297675000\, j_2^2\, j_3^4\, j_6 + 35514000\, j_2^6\, j_4\, j_6 - 858330000\, j_2^3\, j_3^2\, j_4\, j_6 - 6212700000\, j_3^4\, j_4\, j_6 - 1287423000\, j_2^4\, j_4^2\, j_6 \\
&+ 1452802500\, j_2\, j_3^2\, j_4^2\, j_6 + 13651027200\, j_2^2\, j_4^3\, j_6 - 29029363200\, j_4^4\, j_6 + 233887500\, j_2^4\, j_3\, j_5\, j_6 - 7016625000\, j_2\, j_3^3\, j_5\, j_6 - 5401890000\, j_2^2\, j_3\, j_4\, j_5\, j_6 \\
&- 53512974000\, j_3\, j_4^2\, j_5\, j_6 + 960001875\, j_2^3\, j_5^2\, j_6 + 120175650000\, j_3^2\, j_5^2\, j_6 - 46983210000\, j_2\, j_4\, j_5^2\, j_6 - 97524000\, j_5^3\, j_6^2 + 2925720000\, j_2^2\, j_3^2\, j_6^2 \\
&+ 3212136000\, j_2^3\, j_4\, j_6^2 - 133688880000\, j_3^2\, j_4\, j_6^2 - 5787936000\, j_2\, j_4^2\, j_6^2 - 1224720000\, j_2\, j_3\, j_5\, j_6^2 + 40415760000\, j_5^2\, j_6^2 + 2646000\, j_2^6\, j_3\, j_7 \\
&- 158760000\, j_2^3\, j_3^3\, j_7 + 2381400000\, j_3^5\, j_7 - 60385500\, j_2^4\, j_3\, j_4\, j_7 + 1811565000\, j_2\, j_3^3\, j_4\, j_7 - 13778100000\, j_2^2\, j_3\, j_4^2\, j_7 + 58484754000\, j_3\, j_4^3\, j_7 \\
&+ 168399000\, j_2^2\, j_5\, j_7 - 5051970000\, j_2^2\, j_3^2\, j_5\, j_7 - 5370907500\, j_2^3\, j_4\, j_5\, j_7 - 130279590000\, j_3^2\, j_4\, j_5\, j_7 + 46845540000\, j_2\, j_4^2\, j_5\, j_7 \\
&+ 69234952500\, j_2\, j_3\, j_5^2\, j_7 - 111602610000\, j_3^2\, j_5^2\, j_7 - 2517480000\, j_2^2\, j_3\, j_6\, j_7 + 75524400000\, j_3^3\, j_6\, j_7 + 24308505000\, j_2\, j_3\, j_4\, j_6\, j_7 \\
&+ 14581822500\, j_2^2\, j_5\, j_6\, j_7 + 48945060000\, j_4\, j_5\, j_6\, j_7 - 382725000\, j_2^4\, j_7^2 + 11481750000\, j_2^2\, j_3^2\, j_7^2 + 5740875000\, j_2^2\, j_4\, j_7^2 - 33067440000\, j_4^2\, j_7^2 \\
&+ 5197500\, j_2^7\, j_8 - 311850000\, j_2^4\, j_3^2\, j_8 + 4677750000\, j_2\, j_3^4\, j_8 - 896427000\, j_2^5\, j_4\, j_8 + 26892810000\, j_2^2\, j_3^2\, j_4\, j_8 + 38918880000\, j_3^2\, j_4^2\, j_8 \\
&+ 60980850000\, j_3^2\, j_4^2\, j_8 - 155110788000\, j_2\, j_4^3\, j_8 + 153090000\, j_2^3\, j_3\, j_5\, j_8 - 4592700000\, j_3^3\, j_5\, j_8 - 44434372500\, j_2\, j_3\, j_4\, j_5\, j_8 \\
&- 41850978750\, j_2^2\, j_5^2\, j_8 + 301740390000\, j_4\, j_5^2\, j_8 + 161595000\, j_2^2\, j_6\, j_8 - 4847850000\, j_2\, j_3^2\, j_6\, j_8 - 37813230000\, j_2^2\, j_4\, j_6\, j_8 + 173910240000\, j_4^2\, j_6\, j_8 \\
&- 564902100000\, j_3\, j_5\, j_6\, j_8 - 68890500000\, j_3\, j_4\, j_7\, j_8 + 25515000\, j_2^5\, j_3\, j_9 - 765450000\, j_2^2\, j_3^3\, j_9 - 2393307000\, j_2^3\, j_3\, j_4\, j_9 + 37353960000\, j_3^3\, j_4\, j_9 \\
&+ 44824752000\, j_2\, j_3\, j_4^2\, j_9 - 803722500\, j_2^4\, j_5\, j_9 + 34445250000\, j_2\, j_3^2\, j_5\, j_9 + 10103940000\, j_2^2\, j_4\, j_5\, j_9 + 88179840000\, j_4^2\, j_5\, j_9 \\
&- 421609860000\, j_3\, j_5^2\, j_9 + 36741600000\, j_2^2\, j_3\, j_6\, j_9 + 367416000000\, j_3\, j_4\, j_6\, j_9 + 5051970000\, j_2^3\, j_7\, j_9 + 55112400000\, j_3^2\, j_7\, j_9 \\
&- 55112400000\, j_2\, j_4\, j_7\, j_9 - 30051000\, j_2^6\, j_{10} + 1037610000\, j_2^3\, j_3^2\, j_{10} - 4082400000\, j_3^4\, j_{10} + 1316574000\, j_2^4\, j_4\, j_{10} - 5051970000\, j_2\, j_3^2\, j_4\, j_{10} \\
&+ 83770848000\, j_2^2\, j_4^2\, j_{10} - 387991296000\, j_4^3\, j_{10} - 10333575000\, j_2^2\, j_3\, j_5\, j_{10} + 655837560000\, j_3\, j_4\, j_5\, j_{10} - 173604060000\, j_2\, j_5^2\, j_{10} \\
&- 18779040000\, j_2^2\, j_6\, j_{10} - 538876800000\, j_3^2\, j_6\, j_{10} + 102876480000\, j_2\, j_4\, j_6\, j_{10} - 206671500000\, j_2\, j_3\, j_7\, j_{10} + 496011600000\, j_5\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = &-8624\, j_2^{11} + 776160\, j_2^8\, j_3^2 - 23284800\, j_2^5\, j_3^4 + 232848000\, j_2^2\, j_3^6 + 836430\, j_2^9\, j_4 - 61299000\, j_2^6\, j_3^2\, j_4 + 1419579000\, j_2^3\, j_3^4\, j_4 - 10001880000\, j_3^6\, j_4 \\
&+ 2396520\, j_2^7\, j_4^2 + 169759800\, j_2^4\, j_3^2\, j_4^2 - 7249662000\, j_2\, j_3^4\, j_4^2 - 1755542565\, j_2^5\, j_4^3 + 33823449450\, j_2^2\, j_3^2\, j_4^3 + 57104013420\, j_2^3\, j_4^4 \\
&+ 610877651400\, j_2^2\, j_3^3\, j_4^4 - 63190933056\, j_2\, j_4^5 - 10478160\, j_2^7\, j_3\, j_5 + 628689600\, j_2^4\, j_3^3\, j_5 - 9430344000\, j_2\, j_3^5\, j_5 + 377451900\, j_2^5\, j_3\, j_4\, j_5 \\
&- 11323557000\, j_2^2\, j_3^3\, j_4\, j_5 + 17861010300\, j_2^3\, j_3\, j_4^2\, j_5 - 408842154000\, j_3^3\, j_4^2\, j_5 - 515879620200\, j_2\, j_3\, j_4^3\, j_5 - 141455160\, j_2^6\, j_5^2 \\
&+ 4243654800\, j_2^3\, j_3^2\, j_5^2 + 8045262225\, j_2^4\, j_4\, j_5^2 - 164923857000\, j_2\, j_3^2\, j_4\, j_5^2 - 381143580300\, j_2^2\, j_4^2\, j_5^2 + 244888012800\, j_4^3\, j_5^2 - 19096446600\, j_2^2\, j_3\, j_5^3 \\
&+ 1328071059000\, j_3\, j_4\, j_5^3 + 515604058200\, j_2\, j_5^4 - 15245370\, j_2^8\, j_6 + 914722200\, j_2^5\, j_3^2\, j_6 - 13720833000\, j_2^2\, j_3^4\, j_6 + 1616110650\, j_2^6\, j_4\, j_6 \\
&- 41880037500\, j_2^3\, j_3\, j_4\, j_6 - 198098460000\, j_3^3\, j_4\, j_6 - 50739668460\, j_2^2\, j_4^2\, j_6 - 30550786200\, j_2\, j_3^2\, j_4^2\, j_6 + 343850637600\, j_2^3\, j_4^3\, j_6 \\
&- 316389265920\, j_4^4\, j_6 + 9708967800\, j_2^4\, j_3\, j_5\, j_6 - 291269034000\, j_2\, j_3^3\, j_5\, j_6 - 248928167250\, j_2^2\, j_3\, j_4\, j_5\, j_6 - 2166618034800\, j_3\, j_4^2\, j_5\, j_6 \\
&+ 71491116375\, j_2^3\, j_5^2\, j_6 + 4288020282000\, j_3^3\, j_5^2\, j_6 - 1124706303000\, j_2\, j_4\, j_5^2\, j_6 - 3118641750\, j_2^5\, j_6^2 + 93559252500\, j_2^2\, j_3^2\, j_6^2 + 6336225000\, j_2^3\, j_4\, j_6^2 \\
&- 4275803700000\, j_3^2\, j_4\, j_6^2 + 640073664000\, j_2\, j_4^2\, j_6^2 - 308629440000\, j_2\, j_3\, j_5\, j_6^2 + 100018800\, j_2^6\, j_3\, j_7 - 6001128000\, j_2^3\, j_3^3\, j_7 \\
&+ 90016920000\, j_3^5\, j_7 - 7372814400\, j_2^4\, j_3\, j_4\, j_7 + 221184432000\, j_2\, j_3^3\, j_4\, j_7 - 305977155750\, j_2^2\, j_3\, j_4^2\, j_7 + 1008391582800\, j_3\, j_4^3\, j_7 \\
&+ 2470107150\, j_2^5\, j_5\, j_7 - 74103214500\, j_2^2\, j_3^2\, j_5\, j_7 - 91538251875\, j_3^2\, j_4\, j_5\, j_7 - 4395489462000\, j_2^2\, j_4\, j_5\, j_7 + 308767221000\, j_2\, j_4^2\, j_5\, j_7 \\
&+ 1484314713000\, j_2\, j_3\, j_5^2\, j_7 - 1406192886000\, j_5^3\, j_7 - 83587140000\, j_2^3\, j_3\, j_6\, j_7 + 2507614200000\, j_3^3\, j_6\, j_7 + 1928934000000\, j_2\, j_3\, j_4\, j_6\, j_7 \\
&+ 450888322500\, j_2^2\, j_5\, j_6\, j_7 + 1512835380000\, j_4\, j_5\, j_6\, j_7 + 121451400\, j_2^7\, j_8 - 7287084000\, j_2^4\, j_3^2\, j_8 + 109306260000\, j_2\, j_3^4\, j_8 - 23588362350\, j_2^5\, j_4\, j_8 \\
&+ 707650870500\, j_2^2\, j_3^2\, j_4\, j_8 + 1093636687500\, j_3^2\, j_4^2\, j_8 + 4713488010000\, j_3^3\, j_4\, j_8 - 1533337192800\, j_2\, j_4^3\, j_8 + 28934010000\, j_2^2\, j_3\, j_5\, j_8 \\
&- 868020300000\, j_3^3\, j_5\, j_8 - 1961725878000\, j_2\, j_3\, j_4\, j_5\, j_8 - 1304200500750\, j_2^2\, j_5^2\, j_8 + 6138143550000\, j_4\, j_5^2\, j_8 - 6429780000\, j_2^4\, j_6\, j_8 \\
&+ 1928934000000\, j_2\, j_3^2\, j_6\, j_8 - 932318100000\, j_2^2\, j_4\, j_6\, j_8 - 6811892640000\, j_4^2\, j_6\, j_8 - 15624365400000\, j_3\, j_5\, j_6\, j_8 + 1696747500\, j_2^5\, j_3\, j_9 \\
&- 50902425000\, j_2^2\, j_3^3\, j_9 - 120664007100\, j_2^3\, j_3\, j_4\, j_9 + 1329311088000\, j_3^3\, j_4\, j_9 + 1750920948000\, j_2\, j_3\, j_4^2\, j_9 - 8342639550\, j_2^4\, j_5\, j_9 \\
&+ 937461924000\, j_2\, j_3^2\, j_5\, j_9 + 1888150824000\, j_2^2\, j_4\, j_5\, j_9 + 273798403200\, j_4^2\, j_5\, j_9 - 11874517704000\, j_3\, j_5^2\, j_9 + 1527072750000\, j_2^2\, j_3\, j_6\, j_9 \\
&+ 12201885360000\, j_3\, j_4\, j_6\, j_9 - 1153788300\, j_2^6\, j_{10} + 18324873000\, j_2^3\, j_3^2\, j_{10} + 488663280000\, j_3^4\, j_{10} + 51993956700\, j_2^4\, j_4\, j_{10} \\
&+ 730790424000\, j_2\, j_3^2\, j_4\, j_{10} + 2438723700000\, j_2^2\, j_4^2\, j_{10} - 2095152998400\, j_4^3\, j_{10} - 687182737500\, j_2^2\, j_3\, j_5\, j_{10} + 12008440836000\, j_3\, j_4\, j_5\, j_{10} \\
&- 3749847696000\, j_2\, j_5^2\, j_{10} - 819796950000\, j_2^3\, j_6\, j_{10} - 21218274000000\, j_3^2\, j_6\, j_{10} + 1686439440000\, j_2\, j_4\, j_6\, j_{10} + 9258883200000\, j_6^2\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = &-392\, j_2^{11} + 35280\, j_2^8\, j_3^2 - 1058400\, j_2^5\, j_3^4 + 10584000\, j_2^2\, j_3^6 + 40740\, j_2^9\, j_4 - 2856000\, j_2^6\, j_3^2\, j_4 + 61362000\, j_2^3\, j_3^4\, j_4 - 370440000\, j_3^6\, j_4 - 124140\, j_2^7\, j_4^2 \\
&+ 5936400\, j_2^4\, j_3^2\, j_4^2 - 66366000\, j_2\, j_3^4\, j_4^2 - 76735620\, j_2^5\, j_4^3 + 1800273600\, j_2^2\, j_3^2\, j_4^3 + 2536560360\, j_2^3\, j_4^4 + 19717846200\, j_3^2\, j_4^4 - 4249568448\, j_2\, j_4^5 \\
&- 476280\, j_2^7\, j_3\, j_5 + 28576800\, j_2^4\, j_3^3\, j_5 - 428652000\, j_2\, j_3^5\, j_5 + 21886200\, j_2^5\, j_3\, j_4\, j_5 - 656586000\, j_2^2\, j_3^3\, j_4\, j_5 + 417133800\, j_2^3\, j_3\, j_4^2\, j_5 \\
&- 13126374000\, j_3^3\, j_4^2\, j_5 - 15611462100\, j_2\, j_3\, j_4^3\, j_5 - 6429780\, j_2^6\, j_5^2 + 192893400\, j_2^3\, j_3^2\, j_5^2 + 378897750\, j_2^4\, j_4\, j_5^2 - 6934977000\, j_2\, j_3^2\, j_4\, j_5^2 \\
&- 17324320500\, j_2^2\, j_4^2\, j_5^2 + 17620221600\, j_4^3\, j_5^2 - 868020300\, j_2^2\, j_3\, j_5^3 + 45881073000\, j_3\, j_4\, j_5^3 + 23436548100\, j_2\, j_5^4 - 580230\, j_2^8\, j_6 + 34813800\, j_2^5\, j_3^2\, j_6 \\
&- 522207000\, j_2^2\, j_3^4\, j_6 + 60975720\, j_2^6\, j_4\, j_6 - 1609459200\, j_2^3\, j_3^2\, j_4\, j_6 - 6594372000\, j_3^4\, j_4\, j_6 - 1956375720\, j_2^4\, j_4^2\, j_6 + 1731464100\, j_2\, j_3^2\, j_4^2\, j_6 \\
&+ 14895705600\, j_2^2\, j_4^3\, j_6 - 17473605120\, j_4^4\, j_6 + 314174700\, j_2^4\, j_3\, j_5\, j_6 - 9425241000\, j_2\, j_3^3\, j_5\, j_6 - 7624611000\, j_2^2\, j_3\, j_4\, j_5\, j_6 - 83295831600\, j_3\, j_4^2\, j_5\, j_6 \\
&+ 2483247375\, j_2^3\, j_5^2\, j_6 + 147915558000\, j_3^3\, j_5^2\, j_6 - 47125038600\, j_2\, j_4\, j_5^2\, j_6 - 138348000\, j_2^2\, j_6^2 + 4150440000\, j_2^2\, j_3^2\, j_6^2 + 1659204000\, j_2^3\, j_4\, j_6^2 \\
&- 161876880000\, j_3^2\, j_4\, j_6^2 + 10552032000\, j_2\, j_4^2\, j_6^2 - 25412940000\, j_2\, j_3\, j_5\, j_6^2 + 40415760000\, j_5^2\, j_6^2 + 3704400\, j_2^6\, j_3\, j_7 - 222264000\, j_2^3\, j_3^3\, j_7 \\
&+ 3333960000\, j_3^5\, j_7 - 195785100\, j_2^4\, j_3\, j_4\, j_7 + 5873553000\, j_2\, j_3^3\, j_4\, j_7 - 13915881000\, j_2^2\, j_3\, j_4^2\, j_7 + 49257363600\, j_3\, j_4^3\, j_7 + 122472000\, j_2^2\, j_5\, j_7 \\
&- 3674160000\, j_2^3\, j_3^2\, j_5\, j_7 - 3784384800\, j_2^3\, j_4\, j_5\, j_7 - 147976794000\, j_2^2\, j_4\, j_5\, j_7 + 13998549600\, j_2\, j_4^2\, j_5\, j_7 + 67581580500\, j_2\, j_3\, j_5^2\, j_7 \\
&- 66961566000\, j_5^3\, j_7 - 2789640000\, j_2^2\, j_3\, j_6\, j_7 + 83689200000\, j_3^3\, j_6\, j_7 + 79552125000\, j_2\, j_3\, j_4\, j_6\, j_7 + 18255982500\, j_2^2\, j_5\, j_6\, j_7 \\
&- 41334300000\, j_4\, j_5\, j_6\, j_7 + 76545000\, j_2^4\, j_7^2 - 2296350000\, j_2\, j_3^2\, j_7^2 - 1148175000\, j_2^2\, j_4\, j_7^2 - 16533720000\, j_4^2\, j_7^2 + 5499900\, j_2^7\, j_8 - 329994000\, j_2^4\, j_3^2\, j_8 \\
&+ 4949910000\, j_2\, j_3^4\, j_8 - 1024682400\, j_2^2\, j_4\, j_8 + 30740472000\, j_2^2\, j_3^2\, j_4\, j_8 + 45892299600\, j_3^2\, j_4^2\, j_8 + 129177342000\, j_2^3\, j_4^2\, j_8 - 70948029600\, j_2\, j_4^3\, j_8 \\
&+ 949158000\, j_2^3\, j_3\, j_5\, j_8 - 28474740000\, j_3^3\, j_5\, j_8 - 105333574500\, j_2\, j_3\, j_4\, j_5\, j_8 - 55491297750\, j_2^2\, j_5^2\, j_8 + 265366206000\, j_4\, j_5^2\, j_8 - 178605000\, j_2^4\, j_6\, j_8 \\
&+ 5358150000\, j_2\, j_3^2\, j_6\, j_8 - 44549190000\, j_2^2\, j_4\, j_6\, j_8 + 95528160000\, j_4^2\, j_6\, j_8 - 537345900000\, j_3\, j_5\, j_6\, j_8 + 13778100000\, j_3\, j_4\, j_7\, j_8 + 62937000\, j_2^2\, j_3\, j_9 \\
&- 1888110000\, j_2^2\, j_3^3\, j_9 - 4599844200\, j_2^3\, j_3\, j_4\, j_9 + 53030376000\, j_3^3\, j_4\, j_9 + 46294416000\, j_2\, j_3\, j_4^2\, j_9 - 549593100\, j_2^4\, j_5\, j_9 + 41977278000\, j_2\, j_3^2\, j_5\, j_9 \\
&+ 11867536800\, j_2^2\, j_4\, j_5\, j_9 + 31744742400\, j_4^2\, j_5\, j_9 - 461290788000\, j_3\, j_5^2\, j_9 + 53887680000\, j_2^2\, j_3\, j_6\, j_9 + 448247520000\, j_3\, j_4\, j_6\, j_9 - 765450000\, j_2^2\, j_7\, j_9 \\
&- 18370800000\, j_3^2\, j_7\, j_9 + 220449600000\, j_6\, j_7\, j_9 - 46607400\, j_2^6\, j_{10} + 908334000\, j_2^3\, j_3^2\, j_{10} + 14696640000\, j_3^4\, j_{10} + 2088147600\, j_2^4\, j_4\, j_{10} \\
&+ 22320522000\, j_2\, j_3^2\, j_4\, j_{10} + 107652888000\, j_2^2\, j_4^2\, j_{10} - 218686003200\, j_4^3\, j_{10} - 25489485000\, j_2^2\, j_3\, j_5\, j_{10} + 447512688000\, j_3\, j_4\, j_5\, j_{10} \\
&- 163683828000\, j_2\, j_5^2\, j_{10} - 31842720000\, j_2^3\, j_6\, j_{10} - 661348800000\, j_3^2\, j_6\, j_{10} + 88179840000\, j_2\, j_4\, j_6\, j_{10} + 41334300000\, j_2\, j_3\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 5096\, j_2^{11} - 458640\, j_2^8\, j_3^2 + 13759200\, j_2^5\, j_3^4 - 137592000\, j_2^2\, j_3^6 - 443520\, j_2^9\, j_4 + 28627200\, j_2^6\, j_3^2\, j_4 - 520128000\, j_2^3\, j_3^4\, j_4 + 1814400000\, j_3^6\, j_4 \\
& - 7755720\, j_2^7\, j_4^2 + 552661200\, j_2^4\, j_3^2\, j_4^2 - 9599688000\, j_2\, j_3^4\, j_4^2 + 1264264740\, j_2^5\, j_4^3 - 32927002200\, j_2^2\, j_3^2\, j_4^3 - 37554396120\, j_2^3\, j_4^4 \\
& - 90975263400\, j_3^2\, j_4^4 + 26493213024\, j_2\, j_4^5 + 6191640\, j_2^7\, j_3\, j_5 - 371498400\, j_2^4\, j_3^3\, j_5 + 5572476000\, j_2\, j_3^5\, j_5 - 304819200\, j_2^5\, j_3\, j_4\, j_5 \\
& + 9144576000\, j_2^2\, j_3^3\, j_4\, j_5 - 836357400\, j_2^3\, j_3\, j_4^2\, j_5 + 60454512000\, j_3^3\, j_4^2\, j_5 + 15308343900\, j_2\, j_3\, j_4^3\, j_5 + 83587140\, j_2^6\, j_5^2 - 2507614200\, j_2^3\, j_3^2\, j_5^2 \\
& - 4675368600\, j_2^4\, j_4\, j_5^2 + 78535170000\, j_2\, j_3^2\, j_4\, j_5^2 + 234850338900\, j_2^2\, j_4^2\, j_5^2 - 78582409200\, j_3^2\, j_5^2 + 11284263900\, j_2^2\, j_3\, j_5^3 - 193444524000\, j_3\, j_4\, j_5^3 \\
& - 304675125300\, j_2\, j_5^4 + 10104570\, j_2^8\, j_6 - 606274200\, j_2^5\, j_3^2\, j_6 + 9094113000\, j_2^2\, j_3^4\, j_6 - 1065657060\, j_2^6\, j_4\, j_6 + 31054897800\, j_2^3\, j_3^2\, j_4\, j_6 \\
& + 27444420000\, j_3^4\, j_4\, j_6 + 32724232200\, j_2^4\, j_4^2\, j_6 - 115775932500\, j_2\, j_3^2\, j_4^2\, j_6 - 202754242800\, j_2^2\, j_3^4\, j_6 + 121545411840\, j_4^4\, j_6 - 1047645900\, j_2^4\, j_3\, j_5\, j_6 \\
& + 31429377000\, j_2\, j_3^3\, j_5\, j_6 - 28485456300\, j_2^2\, j_3\, j_4\, j_5\, j_6 + 438804491400\, j_3\, j_4^2\, j_5\, j_6 - 47277636525\, j_2^3\, j_5^2\, j_6 - 663369588000\, j_3^2\, j_5^2\, j_6 \\
& + 630393345900\, j_2\, j_4\, j_5^2\, j_6 + 2134188000\, j_2^5\, j_6^2 - 64025640000\, j_2^2\, j_3^2\, j_6^2 - 5592888000\, j_2^3\, j_4\, j_6^2 + 829981080000\, j_3^2\, j_4\, j_6^2 - 272957040000\, j_2\, j_4^2\, j_6^2 \\
& + 524486340000\, j_2\, j_3\, j_5\, j_6^2 - 303118200000\, j_5^2\, j_6^2 - 18144000\, j_2^6\, j_3\, j_7 + 1088640000\, j_2^3\, j_3^3\, j_7 - 16329600000\, j_3^5\, j_7 - 1674294300\, j_2^4\, j_3\, j_4\, j_7 \\
& + 50228829000\, j_2\, j_3^3\, j_4\, j_7 + 289794777300\, j_2^2\, j_3\, j_4^2\, j_7 - 382003727400\, j_3\, j_4^3\, j_7 - 1241559900\, j_2^5\, j_5\, j_7 + 37246797000\, j_2^2\, j_3^2\, j_5\, j_7 \\
& + 46060953750\, j_2^3\, j_4\, j_5\, j_7 + 700386750000\, j_2^2\, j_4\, j_5\, j_7 + 33356780100\, j_2\, j_4^2\, j_5\, j_7 - 863680198500\, j_2\, j_3\, j_5^2\, j_7 + 468730962000\, j_3^3\, j_7 \\
& + 14441490000\, j_2^3\, j_3\, j_6\, j_7 - 433244700000\, j_3^3\, j_6\, j_7 - 468160155000\, j_2\, j_3\, j_4\, j_6\, j_7 - 294506887500\, j_2^2\, j_5\, j_6\, j_7 + 106091370000\, j_4\, j_5\, j_6\, j_7 \\
& - 1377810000\, j_2^4\, j_7^2 + 41334300000\, j_2\, j_3^2\, j_7^2 + 2066715000\, j_2^2\, j_4\, j_7^2 + 97430850000\, j_4^2\, j_7^2 - 74220300\, j_7^2\, j_8 + 4453218000\, j_2^4\, j_3^2\, j_8 \\
& - 66798270000\, j_2\, j_3^4\, j_8 + 14603255100\, j_2^3\, j_4\, j_8 - 438097653000\, j_2^2\, j_3^2\, j_4\, j_8 - 671522140800\, j_2^3\, j_4^2\, j_8 - 391144950000\, j_3^2\, j_4^2\, j_8 \\
& + 122992506000\, j_2\, j_4^3\, j_8 - 3490452000\, j_2^2\, j_3^3\, j_5\, j_8 + 104713560000\, j_3^3\, j_5\, j_8 + 1620511231500\, j_2\, j_3\, j_4\, j_5\, j_8 + 818729147250\, j_2^2\, j_5^2\, j_8 \\
& - 2372175477000\, j_4\, j_5^2\, j_8 + 4618215000\, j_2^4\, j_6\, j_8 - 138546450000\, j_2\, j_3^2\, j_6\, j_8 + 708653610000\, j_2^2\, j_4\, j_6\, j_8 - 789944400000\, j_4^2\, j_6\, j_8 \\
& + 2335387950000\, j_3\, j_5\, j_6\, j_8 + 310007250000\, j_3\, j_4\, j_7\, j_8 + 1860043500000\, j_7^2\, j_8 - 757285200\, j_2^5\, j_3\, j_9 + 22718556000\, j_2^2\, j_3^3\, j_9 \\
& + 43385706000\, j_2^3\, j_3\, j_4\, j_9 - 279236160000\, j_3^3\, j_4\, j_9 + 330453950400\, j_2\, j_3\, j_4^2\, j_9 + 2571912000\, j_2^4\, j_5\, j_9 - 383857866000\, j_2\, j_3^2\, j_5\, j_9 \\
& - 61009426800\, j_2^2\, j_4\, j_5\, j_9 - 513206668800\, j_4^2\, j_5\, j_9 + 2462697594000\, j_3\, j_5^2\, j_9 - 631955520000\, j_2^2\, j_3\, j_6\, j_9 - 2006091360000\, j_3\, j_4\, j_6\, j_9 \\
& + 24800580000\, j_2^3\, j_7\, j_9 - 347208120000\, j_2\, j_4\, j_7\, j_9 + 713059200\, j_2^6\, j_{10} - 20064996000\, j_2^3\, j_3^2\, j_{10} - 39803400000\, j_3^4\, j_{10} - 28627830000\, j_2^4\, j_4\, j_{10} \\
& - 163500120000\, j_2\, j_3^2\, j_4\, j_{10} - 1750039149600\, j_2^2\, j_4^2\, j_{10} + 1724797670400\, j_4^3\, j_{10} + 306700506000\, j_2^2\, j_3\, j_5\, j_{10} - 2787585192000\, j_3\, j_4\, j_5\, j_{10} \\
& + 2403176202000\, j_2\, j_5^2\, j_{10} + 531528480000\, j_2^3\, j_6\, j_{10} + 3012811200000\, j_3^2\, j_6\, j_{10} - 837708480000\, j_2\, j_4\, j_6\, j_{10} - 744017400000\, j_2\, j_3\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 5488\, j_2^{10}\, j_3 - 493920\, j_2^7\, j_3^3 + 14817600\, j_2^4\, j_3^5 - 148176000\, j_2\, j_3^7 + 79870\, j_2^8\, j_3\, j_4 - 4792200\, j_2^5\, j_3^3\, j_4 + 71883000\, j_2^2\, j_3^5\, j_4 - 59795820\, j_2^6\, j_3\, j_4^2 \\
& + 2371912200\, j_2^3\, j_3^3\, j_4^2 - 17341128000\, j_3^5\, j_4^2 + 2488273830\, j_2^4\, j_3\, j_4^3 - 39790472400\, j_2\, j_3^3\, j_4^3 - 70269330600\, j_2^2\, j_3\, j_4^4 - 86770852128\, j_3\, j_4^5 \\
& + 6667920\, j_2^6\, j_3^2\, j_5 - 400075200\, j_2^3\, j_3^4\, j_5 + 6001128000\, j_3^6\, j_5 + 9887220\, j_2^7\, j_4\, j_5 - 528538500\, j_2^4\, j_3^2\, j_4\, j_5 + 6957657000\, j_2\, j_3^4\, j_4\, j_5 \\
& - 993245085\, j_2^5\, j_4^2\, j_5 + 13383553050\, j_2^2\, j_3^2\, j_4^2\, j_5 + 23895147420\, j_2^3\, j_4^3\, j_5 - 49969888200\, j_3^2\, j_4^3\, j_5 + 68630684400\, j_2\, j_4^4\, j_5 + 90016920\, j_2^5\, j_3\, j_5^2 \\
& - 2700507600\, j_2^2\, j_3^3\, j_5^2 - 2420990775\, j_2^2\, j_3\, j_4\, j_5^2 + 85730400000\, j_3^3\, j_4\, j_5^2 + 28687841280\, j_2\, j_3\, j_4^2\, j_5^2 + 12152284200\, j_2\, j_3^2\, j_5^3 \\
& - 56180202750\, j_2^2\, j_4\, j_5^3 - 122762871000\, j_4^2\, j_5^3 - 328111673400\, j_3\, j_5^4 + 16321410\, j_2^7\, j_3\, j_6 - 979284600\, j_2^4\, j_3^3\, j_6 + 14689269000\, j_2\, j_3^5\, j_6 \\
& - 1954956150\, j_2^5\, j_3\, j_4\, j_6 + 58648684500\, j_2^2\, j_3^3\, j_4\, j_6 + 58880448720\, j_2^3\, j_3\, j_4^2\, j_6 - 118748316600\, j_3^3\, j_4^2\, j_6 - 142003951200\, j_2\, j_3\, j_4^3\, j_6 \\
& - 398249460\, j_2^6\, j_5\, j_6 + 12973867200\, j_2^3\, j_3^2\, j_5\, j_6 - 30791502000\, j_3^4\, j_5\, j_6 + 14393005830\, j_2^4\, j_4\, j_5\, j_6 - 137182503150\, j_2\, j_3^2\, j_4\, j_5\, j_6 \\
& + 65446908120\, j_2^2\, j_4^2\, j_5\, j_6 + 203172649920\, j_3^2\, j_4\, j_5\, j_6 - 236672164575\, j_2^2\, j_3\, j_5^2\, j_6 + 43915397400\, j_3\, j_4\, j_5^2\, j_6 + 150746192100\, j_2\, j_5^3\, j_6 \\
& + 2635416000\, j_2^4\, j_3\, j_6^2 - 79062480000\, j_2\, j_3^3\, j_6^2 + 190968435000\, j_2^2\, j_3\, j_4\, j_6^2 + 78253776000\, j_3\, j_4^2\, j_6^2 - 19735852500\, j_3^2\, j_5\, j_6^2 \\
& + 788719680000\, j_3^2\, j_5\, j_6^2 - 238861224000\, j_2\, j_4\, j_5\, j_6^2 - 3360420\, j_2^8\, j_7 + 201625200\, j_2^5\, j_3^2\, j_7 - 3024378000\, j_2^2\, j_3^4\, j_7 + 489145230\, j_2^6\, j_4\, j_7 \\
& - 18940805100\, j_2^3\, j_3^2\, j_4\, j_7 + 127993446000\, j_3^4\, j_4\, j_7 - 6300367920\, j_2^4\, j_4^2\, j_7 + 388146682350\, j_2\, j_3^2\, j_4^2\, j_7 - 31090741920\, j_2^2\, j_4^3\, j_7 \\
& - 69044814720\, j_4^4\, j_7 - 619759350\, j_2^4\, j_3\, j_5\, j_7 + 18592780500\, j_2\, j_3^3\, j_5\, j_7 + 196807528575\, j_2^2\, j_3\, j_4\, j_5\, j_7 + 912744012600\, j_3\, j_4^2\, j_5\, j_7 \\
& - 707275800\, j_2^3\, j_5^2\, j_7 - 890203041000\, j_2^2\, j_3^2\, j_5^2\, j_7 - 118918781100\, j_2\, j_4\, j_5^2\, j_7 + 3083913000\, j_5^3\, j_6\, j_7 - 92517390000\, j_2^2\, j_3^2\, j_6\, j_7 \\
& - 64430478000\, j_2^3\, j_4\, j_6\, j_7 - 617105790000\, j_3^2\, j_4\, j_6\, j_7 - 98100072000\, j_2\, j_4^2\, j_6\, j_7 - 333544837500\, j_2\, j_3\, j_5\, j_6\, j_7 - 202538070000\, j_5^2\, j_6\, j_7 \\
& - 3214890000\, j_2^3\, j_3\, j_7^2 + 96446700000\, j_3^3\, j_7^2 + 2411167500\, j_2\, j_3\, j_4\, j_7^2 - 83349000\, j_2^6\, j_3\, j_8 + 5000940000\, j_2^3\, j_3^3\, j_8 - 75014100000\, j_3^5\, j_8 \\
& + 18237356550\, j_2^4\, j_3\, j_4\, j_8 - 547120696500\, j_2\, j_3^3\, j_4\, j_8 - 1115245341000\, j_2^2\, j_3\, j_4^2\, j_8 - 2316980408400\, j_3\, j_4^3\, j_8 - 1360970100\, j_2^5\, j_5\, j_8 \\
& + 40829103000\, j_2^2\, j_3^2\, j_5\, j_8 - 19876440150\, j_2^3\, j_4\, j_5\, j_8 + 2254694211000\, j_3^2\, j_4\, j_5\, j_8 + 323344450800\, j_2\, j_4^2\, j_5\, j_8 + 1116129435750\, j_2\, j_3\, j_5^2\, j_8 \\
& - 286446699000\, j_5^3\, j_8 + 5358150000\, j_2^3\, j_3\, j_6\, j_8 - 160744500000\, j_3^3\, j_6\, j_8 + 962859555000\, j_2\, j_3\, j_4\, j_6\, j_8 + 417131977500\, j_2^2\, j_5\, j_6\, j_8 \\
& + 2615083380000\, j_4\, j_5\, j_6\, j_8 - 803722500\, j_2^4\, j_7\, j_8 + 24111675000\, j_2\, j_3^2\, j_7\, j_8 + 33756345000\, j_2^2\, j_4\, j_7\, j_8 - 3472081200000\, j_4^2\, j_7\, j_8 \\
& - 29569050\, j_2^7\, j_9 - 220279500\, j_2^4\, j_3^2\, j_9 + 33220530000\, j_2\, j_3^4\, j_9 + 2362518900\, j_2^5\, j_4\, j_9 + 18873445500\, j_2^2\, j_3^2\, j_4\, j_9 - 87591974400\, j_3^2\, j_4^2\, j_9 \\
& + 1071201348000\, j_3^3\, j_4^2\, j_9 - 237424219200\, j_2\, j_4^3\, j_9 - 15994077750\, j_2^3\, j_3\, j_5\, j_9 - 327918780000\, j_3^3\, j_5\, j_9 - 137367657000\, j_2\, j_3\, j_4\, j_5\, j_9 \\
& + 337081216500\, j_2^2\, j_5^2\, j_9 + 957302388000\, j_4\, j_5^2\, j_9 - 30148524000\, j_2^4\, j_6\, j_9 - 745854480000\, j_2\, j_3^2\, j_6\, j_9 - 354985092000\, j_2^2\, j_4\, j_6\, j_9 \\
& - 890616384000\, j_4^2\, j_6\, j_9 + 72335025000\, j_2^2\, j_3\, j_7\, j_9 - 810152280000\, j_3\, j_4\, j_7\, j_9 + 1994422500\, j_2^5\, j_3\, j_{10} - 59832675000\, j_2^2\, j_3^3\, j_{10} \\
& - 77458947300\, j_2^3\, j_3\, j_4\, j_{10} - 368701956000\, j_3^3\, j_4\, j_{10} - 2576504700000\, j_2\, j_3\, j_4^2\, j_{10} + 12854201850\, j_2^4\, j_5\, j_{10} + 422115057000\, j_2\, j_3^2\, j_5\, j_{10} \\
& - 151393762800\, j_2^2\, j_4\, j_5\, j_{10} - 1301534438400\, j_4^2\, j_5\, j_{10} + 2742944148000\, j_3\, j_5^2\, j_{10} + 1650310200000\, j_2^2\, j_3\, j_6\, j_{10} + 3538216080000\, j_3\, j_4\, j_6\, j_{10} \\
& - 1234517760000\, j_2\, j_5\, j_6\, j_{10} - 27326565000\, j_2^3\, j_7\, j_{10} - 1350253800000\, j_3^2\, j_7\, j_{10} + 462944160000\, j_2\, j_4\, j_7\, j_{10} + 2314720800000\, j_6\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & 2296\, j_2^{10}\, j_3 - 206640\, j_2^7\, j_3^3 + 6199200\, j_2^4\, j_3^5 - 61992000\, j_2\, j_3^7 + 146790\, j_2^8\, j_3\, j_4 - 8807400\, j_2^5\, j_3^3\, j_4 + 132111000\, j_2^2\, j_3^5\, j_4 - 35779140\, j_2^6\, j_3\, j_4^2 \\
& + 1331969400\, j_2^3\, j_3^3\, j_4^2 - 7757856000\, j_3^5\, j_4^2 + 1278861210\, j_2^4\, j_3\, j_4^3 - 16894963800\, j_2\, j_3^3\, j_4^3 - 35787922200\, j_2^2\, j_3\, j_4^4 + 13355466624\, j_3\, j_4^5 \\
& + 2789640\, j_2^6\, j_3^2\, j_5 - 167378400\, j_2^3\, j_3^4\, j_5 + 2510676000\, j_3^6\, j_5 + 5556600\, j_2^7\, j_4\, j_5 - 214496100\, j_2^4\, j_3^2\, j_4\, j_5 + 1433943000\, j_2\, j_3^4\, j_4\, j_5 - 584809875\, j_2^5\, j_4^2\, j_5 \\
& + 6544779750\, j_2^2\, j_3^2\, j_4^2\, j_5 + 15707821320\, j_2^3\, j_4^3\, j_5 - 56479056300\, j_3^2\, j_4^3\, j_5 - 18236693160\, j_2\, j_4^4\, j_5 + 37660140\, j_2^5\, j_3\, j_5^2 - 1129804200\, j_2^2\, j_3^3\, j_5^2 \\
& - 428269275\, j_2^3\, j_3\, j_4\, j_5^2 + 36925308000\, j_3^3\, j_4\, j_5^2 + 122126125950\, j_2\, j_3\, j_4^2\, j_5^2 + 5084118900\, j_2\, j_3^2\, j_5^3 - 31744742400\, j_2^2\, j_4\, j_5^3 + 54897855300\, j_4^2\, j_5^3 \\
& - 137271210300\, j_3\, j_5^4 + 8906940\, j_2^7\, j_3\, j_6 - 534416400\, j_2^4\, j_3^3\, j_6 + 8016246000\, j_2\, j_3^5\, j_6 - 1079717850\, j_2^5\, j_3\, j_4\, j_6 + 32391535500\, j_2^2\, j_3^3\, j_4\, j_6 \\
& + 32992395120\, j_2^3\, j_3\, j_4^2\, j_6 - 73249361100\, j_3^3\, j_4^2\, j_6 - 94510476000\, j_2\, j_3\, j_4^3\, j_6 - 227775240\, j_2^6\, j_5\, j_6 + 7217853300\, j_2^3\, j_3^2\, j_5\, j_6 - 11537883000\, j_3^4\, j_5\, j_6 \\
& + 9344669490\, j_2^4\, j_4\, j_5\, j_6 - 102778101450\, j_2\, j_3^2\, j_4\, j_5\, j_6 - 5497768080\, j_2^2\, j_4^2\, j_5\, j_6 - 44830000800\, j_4^3\, j_5\, j_6 - 135755619300\, j_2^2\, j_3\, j_5^2\, j_6 \\
& + 425249246700\, j_3\, j_4\, j_5^2\, j_6 + 92816170650\, j_2\, j_5^3\, j_6 + 1242297000\, j_2^4\, j_3\, j_6^2 - 37268910000\, j_2^2\, j_3^3\, j_6^2 + 118349505000\, j_2^2\, j_3\, j_4\, j_6^2 \\
& - 285039000000\, j_3\, j_4^3\, j_6^2 - 12310987500\, j_2^3\, j_5\, j_6^2 + 355781160000\, j_3^3\, j_5\, j_6^2 - 46933020000\, j_2\, j_4\, j_5\, j_6^2 - 1893780\, j_2^8\, j_7 + 113626800\, j_2^5\, j_3^2\, j_7 \\
& - 1704402000\, j_2^2\, j_3^4\, j_7 + 283936590\, j_2^6\, j_4\, j_7 - 10424748600\, j_2^3\, j_3^2\, j_4\, j_7 + 57199527000\, j_3^4\, j_4\, j_7 - 4568759640\, j_2^4\, j_4^2\, j_7 + 218617659450\, j_2\, j_3^2\, j_4^2\, j_7 \\
& - 3463683120\, j_2^2\, j_4^3\, j_7 + 8077903200\, j_4^4\, j_7 - 43630650\, j_2^4\, j_3\, j_5\, j_7 + 1308919500\, j_2\, j_3^3\, j_5\, j_7 + 109249999425\, j_2^2\, j_3\, j_4\, j_5\, j_7 - 95854241700\, j_3\, j_4^2\, j_5\, j_7 \\
& - 454677300\, j_2^3\, j_5^2\, j_7 - 367668598500\, j_3^3\, j_5^2\, j_7 - 72355692150\, j_2\, j_4\, j_5^2\, j_7 + 1801359000\, j_2^5\, j_6\, j_7 - 54040770000\, j_2^2\, j_3^2\, j_6\, j_7 \\
& - 43461522000\, j_2^3\, j_4\, j_6\, j_7 - 43226055000\, j_2^2\, j_4\, j_6\, j_7 + 80450982000\, j_2\, j_4^2\, j_6\, j_7 - 155003625000\, j_2\, j_3\, j_5\, j_6\, j_7 + 6200145000\, j_5^2\, j_6\, j_7 \\
& - 1607445000\, j_2^2\, j_3\, j_7^2 + 48223350000\, j_3^3\, j_7^2 + 17567077500\, j_2\, j_3\, j_4\, j_7^2 - 34870500\, j_2^6\, j_3\, j_8 + 2092230000\, j_2^3\, j_3^3\, j_8 - 31383450000\, j_3^5\, j_8 \\
& + 8471745450\, j_2^4\, j_3\, j_4\, j_8 - 254152363500\, j_2\, j_3^3\, j_4\, j_8 - 574638624000\, j_2^2\, j_3\, j_4^2\, j_8 - 292701956400\, j_3\, j_4^3\, j_8 - 766980900\, j_2^5\, j_5\, j_8 \\
& + 23009427000\, j_2^2\, j_3^2\, j_5\, j_8 - 9587261250\, j_2^3\, j_4\, j_5\, j_8 + 871258153500\, j_3^2\, j_4\, j_5\, j_8 + 379035531000\, j_2\, j_4^2\, j_5\, j_8 + 554292963000\, j_2\, j_3\, j_5^2\, j_8 \\
& - 184144306500\, j_5^3\, j_8 + 3291435000\, j_2^3\, j_3\, j_6\, j_8 - 98743050000\, j_3^3\, j_6\, j_8 + 433780515000\, j_2\, j_3\, j_4\, j_6\, j_8 + 226305292500\, j_2^2\, j_5\, j_6\, j_8 \\
& - 682015950000\, j_4\, j_5\, j_6\, j_8 - 114817500\, j_2^4\, j_7\, j_8 + 3444525000\, j_2\, j_3^2\, j_7\, j_8 - 2066715000\, j_2^2\, j_4\, j_7\, j_8 - 235605510000\, j_4^2\, j_7\, j_8 \\
& - 16176510\, j_2^7\, j_9 - 139311900\, j_2^4\, j_3^2\, j_9 + 18738216000\, j_2\, j_3^4\, j_9 + 1306674180\, j_2^5\, j_4\, j_9 + 10745387100\, j_2^2\, j_3^2\, j_4\, j_9 - 53108758080\, j_3^2\, j_4^2\, j_9 \\
& + 492668114400\, j_3^3\, j_4^2\, j_9 + 99665264160\, j_2\, j_4^3\, j_9 - 10124607150\, j_2^3\, j_3\, j_5\, j_9 - 145772298000\, j_3^3\, j_5\, j_9 - 68284263600\, j_2\, j_3\, j_4\, j_5\, j_9 \\
& + 190468454400\, j_2^2\, j_5^2\, j_9 - 447898474800\, j_4\, j_5^2\, j_9 - 17819676000\, j_2^4\, j_6\, j_9 - 402320520000\, j_2\, j_3^2\, j_6\, j_9 - 108938844000\, j_2^2\, j_4\, j_6\, j_9 \\
& + 242494560000\, j_4^2\, j_6\, j_9 + 31000725000\, j_2^2\, j_3\, j_7\, j_9 + 90935460000\, j_3\, j_4\, j_7\, j_9 + 744017400000\, j_7^2\, j_9 + 1109902500\, j_2^5\, j_3\, j_{10} \\
& - 33297075000\, j_2^2\, j_3^3\, j_{10} - 47211425100\, j_2^2\, j_3\, j_4\, j_{10} - 82025622000\, j_3^3\, j_4\, j_{10} - 1071660618000\, j_2\, j_3\, j_4^2\, j_{10} + 7065868950\, j_2^4\, j_5\, j_{10} \\
& + 237534444000\, j_2\, j_3^2\, j_5\, j_{10} - 93498186600\, j_2^2\, j_4\, j_5\, j_{10} + 571405363200\, j_4^2\, j_5\, j_{10} + 1071385056000\, j_3\, j_5^2\, j_{10} + 936910800000\, j_2^2\, j_3\, j_6\, j_{10} \\
& - 705438720000\, j_3\, j_4\, j_6\, j_{10} - 744017400000\, j_2\, j_5\, j_6\, j_{10} - 13089195000\, j_2^3\, j_7\, j_{10} - 537345900000\, j_3^2\, j_7\, j_{10} - 140536620000\, j_2\, j_4\, j_7\, j_{10},
\end{aligned}
$$

$$
\begin{aligned}
0 = {} & -3136\, j_2^{12} + 282240\, j_2^9 j_3^2 - 8467200\, j_2^6 j_3^4 + 84672000\, j_2^3 j_3^6 + 281232\, j_2^{10} j_4 - 23319030\, j_2^7 j_3^2 j_4 + 639815400\, j_2^4 j_3^4 j_4 - 5800599000\, j_2 j_3^6 j_4 \\
& + 2770530\, j_2^8 j_4^2 + 226037700\, j_2^5 j_3^2 j_4^2 - 9274608000\, j_2^2 j_3^4 j_4^2 - 640259640\, j_2^6 j_4^3 + 6366498750\, j_2^3 j_3^2 j_4^3 - 8138799000\, j_3^4 j_4^3 + 20176761420\, j_2^2 j_4^4 \\
& + 363254211900\, j_2 j_3^2 j_4^4 - 4691010024\, j_2^3 j_4^5 - 21019554432\, j_4^6 - 3810240\, j_2^8 j_3 j_5 + 228614400\, j_2^5 j_3^3 j_5 - 3429216000\, j_2^2 j_3^5 j_5 \\
& + 78287580\, j_2^6 j_3 j_4 j_5 - 3083856300\, j_2^3 j_3^3 j_4 j_5 + 22056867000\, j_3^5 j_4 j_5 + 11236644405\, j_2^4 j_3^2 j_4^2 j_5 - 178230184650\, j_2 j_3^4 j_4^2 j_5 \\
& - 277610344200\, j_2^2 j_3^3 j_4^3 j_5 + 181783133820\, j_3^2 j_4^4 j_5 - 51438240\, j_2^7 j_5^2 + 1543147200\, j_2^4 j_3^2 j_5^2 + 2728267920\, j_2^5 j_4 j_5^2 - 70957597725\, j_2^2 j_3^2 j_4 j_5^2 \\
& - 136382752485\, j_2^3 j_4^2 j_5^2 - 186183465300\, j_3^3 j_4 j_5^2 - 163758820230\, j_2 j_4^3 j_5^2 - 6944162400\, j_2^5 j_3 j_5^3 + 674679111750\, j_2 j_3 j_4 j_5^3 + 187492384800\, j_2^2 j_5^4 \\
& + 380564900100\, j_4 j_5^4 - 6131664\, j_2^9 j_6 + 329995890\, j_2^6 j_3^2 j_6 - 3244260600\, j_2^3 j_3^4 j_6 - 34113555000\, j_3^6 j_6 + 636743646\, j_2^7 j_4 j_6 \\
& - 11369425410\, j_2^4 j_3^2 j_4 j_6 - 231986519100\, j_2 j_3^4 j_4 j_6 - 18847621944\, j_2^5 j_4^2 j_6 - 192900955680\, j_2^2 j_3^2 j_4^2 j_6 + 92875822776\, j_2^3 j_4^3 j_6 \\
& + 122638780620\, j_3^2 j_4^3 j_6 + 44658819936\, j_2 j_4^4 j_6 + 5036422860\, j_2^5 j_3 j_5 j_6 - 151092685800\, j_2^2 j_3^3 j_5 j_6 - 146045495610\, j_2^3 j_3 j_4 j_5 j_6 \\
& + 363873752550\, j_3^3 j_4 j_5 j_6 - 797037628320\, j_2 j_3 j_4^2 j_5 j_6 + 30644331480\, j_2^4 j_5^2 j_6 + 2333407348125\, j_2 j_3^2 j_5^2 j_6 - 392227832115\, j_2^2 j_4 j_5^2 j_6 \\
& - 691147287360\, j_4^2 j_5^2 j_6 - 105898476600\, j_3 j_5^3 j_6 - 1263729600\, j_2^6 j_6^2 + 37233189000\, j_2^3 j_3^2 j_6^2 + 20360970000\, j_3^4 j_6^2 + 2303834400\, j_2^4 j_4 j_6^2 \\
& - 2508605883000\, j_2 j_3^2 j_4 j_6^2 + 82826841600\, j_2^2 j_4^2 j_6^2 + 422227468800\, j_4^3 j_6^2 - 68209249500\, j_2^2 j_3 j_5 j_6^2 - 419956488000\, j_3 j_4 j_5 j_6^2 \\
& + 459086292000\, j_2 j_5^2 j_6^2 + 48977460\, j_2^7 j_3 j_7 - 2938647600\, j_2^4 j_3^3 j_7 + 44079714000\, j_2 j_3^5 j_7 - 4568596830\, j_2^5 j_3 j_4 j_7 + 137057904900\, j_2^2 j_3^3 j_4 j_7 \\
& - 89936802900\, j_3^5 j_4 j_7 - 581967807750\, j_3^2 j_4^3 j_7 + 218629053720\, j_2 j_3 j_4^4 j_7 + 838252800\, j_2^6 j_5 j_7 - 29500187850\, j_2^3 j_3^2 j_5 j_7 \\
& + 130578115500\, j_3^4 j_5 j_7 - 34356457800\, j_2^4 j_4 j_5 j_7 - 2152360817775\, j_2 j_3^2 j_4 j_5 j_7 + 145612472040\, j_2^2 j_4^2 j_5 j_7 + 159765336360\, j_4^3 j_5 j_7 \\
& + 539908626600\, j_2^2 j_3 j_5^2 j_7 + 598810004100\, j_3 j_4 j_5^2 j_7 - 515604058200\, j_2 j_5^3 j_7 - 42615153000\, j_2^4 j_3 j_6 j_7 + 1278454590000\, j_2 j_3^3 j_6 j_7 \\
& + 1009261134000\, j_2^2 j_3 j_4 j_6 j_7 - 301261437000\, j_3 j_4^2 j_6 j_7 + 162244782000\, j_2^3 j_5 j_6 j_7 + 618062602500\, j_3^2 j_5 j_6 j_7 + 507240751500\, j_2 j_4 j_5 j_6 j_7 \\
& + 428652000\, j_2^5 j_7^2 - 12859560000\, j_2^2 j_3^2 j_7^2 + 8588349000\, j_2^3 j_4 j_7^2 - 221482957500\, j_2^2 j_3 j_4 j_7^2 - 140949963000\, j_2 j_4^2 j_7^2 + 44135280\, j_2^8 j_8 \\
& - 2648116800\, j_2^5 j_3 j_8 + 39721752000\, j_2^2 j_3 j_8 - 8768008620\, j_2^3 j_4 j_8 + 246336863850\, j_2^3 j_3^2 j_4 j_8 + 501101842500\, j_3^3 j_4 j_8 + 406754822880\, j_2^4 j_4^2 j_8 \\
& + 4176898374600\, j_2 j_3^2 j_4^2 j_8 - 67832341920\, j_2^2 j_4^3 j_8 - 262820013120\, j_4^4 j_8 + 15592216500\, j_2^2 j_3 j_5 j_8 - 467766495000\, j_2 j_3^3 j_5 j_8 \\
& - 687230960850\, j_2^3 j_3 j_4 j_5 j_8 - 2776052922300\, j_3 j_4^2 j_5 j_8 - 499015225800\, j_2^3 j_5^2 j_8 - 1762804559250\, j_3^2 j_5^2 j_8 + 1428947418150\, j_2 j_4 j_5^2 j_8 \\
& - 3143448000\, j_2^5 j_6 j_8 + 94303440000\, j_2^2 j_3^2 j_6 j_8 - 396610263000\, j_2^3 j_4 j_6 j_8 - 200930625000\, j_3^3 j_4 j_6 j_8 + 324520182000\, j_2 j_4^2 j_6 j_8 \\
& - 6907994887500\, j_2 j_3 j_5 j_6 j_8 - 3211675110000\, j_5^2 j_6 j_8 + 4018612500\, j_2^2 j_3 j_7 j_8 - 120558375000\, j_3^2 j_7 j_8 - 188071065000\, j_2 j_3 j_4 j_7 j_8 \\
& + 773359650\, j_2^6 j_3 j_9 - 20378830500\, j_2^3 j_3^3 j_9 - 84658770000\, j_3^5 j_9 - 53333800380\, j_2^4 j_3 j_4 j_9 + 428990328900\, j_2 j_3^3 j_4 j_9 + 900564172200\, j_2^2 j_3 j_4^2 j_9 \\
& - 894518341920\, j_3 j_4^3 j_9 - 2537619840\, j_2^3 j_5 j_9 + 427435699950\, j_2^2 j_3^2 j_5 j_9 + 63644718060\, j_2^3 j_4 j_5 j_9 + 753083389800\, j_3^3 j_4 j_5 j_9 \\
& + 797818124880\, j_2 j_4^2 j_5 j_9 - 5519741087700\, j_2 j_3 j_5^2 j_9 - 1218700501200\, j_5^3 j_9 + 716277492000\, j_2^3 j_3 j_6 j_9 + 1324534680000\, j_3^2 j_6 j_9 \\
& + 5401566324000\, j_2 j_3 j_4 j_6 j_9 - 6429780000\, j_2^4 j_7 j_9 - 110913705000\, j_2 j_3^2 j_7 j_9 - 58143582000\, j_2^2 j_4 j_7 j_9 + 585293688000\, j_4^2 j_7 j_9 \\
& - 469135800\, j_2^7 j_{10} + 2125399500\, j_2^4 j_3^2 j_{10} + 358460235000\, j_2 j_3^4 j_{10} + 21321354600\, j_2^5 j_4 j_{10} + 531383044500\, j_2^2 j_3^2 j_4 j_{10} + 95120946000\, j_2^3 j_4^2 j_{10} \\
& + 1236538548000\, j_2^2 j_4^3 j_{10} - 673032628800\, j_2 j_4^4 j_{10} - 325105751250\, j_3^3 j_5 j_{10} - 786040605000\, j_3^2 j_5^2 j_{10} + 4705910055000\, j_2 j_3 j_4 j_5 j_{10} \\
& - 1332411160500\, j_2^3 j_5^2 j_{10} - 133923132000\, j_4 j_5^2 j_{10} - 336063168000\, j_2 j_6 j_{10} - 12730964400000\, j_2 j_3^2 j_6 j_{10} + 263069856000\, j_2^2 j_4 j_6 j_{10} \\
& + 1807686720000\, j_4^2 j_6 j_{10} + 2893401000000\, j_3 j_5 j_6 j_{10} + 303807105000\, j_2^2 j_3 j_7 j_{10} + 3480348060000\, j_3 j_4 j_7 j_{10} + 5208121800000\, j_7^2 j_{10},
\end{aligned}
$$

DGA MI, La Roche Marguerite, 35174 Bruz, France.

IRMAR, Université de Rennes 1, Campus de Beaulieu, 35042 Rennes, France.
E-mail address: reynald.lercier@m4x.org

Institut de Mathématiques de Luminy, UMR 6206 du CNRS, Luminy, Case 907, 13288 Marseille, France.
E-mail address: ritzenth@iml.univ-mrs.fr